

Katsuyuki Tamai

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9141422/katsuyuki-tamai-publications-by-year.pdf>

Version: 2024-04-17

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

60
papers

11,179
citations

34
h-index

62
g-index

62
ext. papers

11,789
ext. citations

9.5
avg, IF

5.05
L-index

#	Paper	IF	Citations
60	Phosphorylation of MCM4 by Cdc7 kinase facilitates its interaction with Cdc45 on the chromatin. <i>Journal of Biological Chemistry</i> , 2006 , 281, 39249-61	5.4	138
59	Different requirements for the association of ATR-ATRIP and 9-1-1 to the stalled replication forks. <i>Gene</i> , 2006 , 377, 88-95	3.8	20
58	Isolation and characterization of the TIGA genes, whose transcripts are induced by growth arrest. <i>Nucleic Acids Research</i> , 2006 , 34, 4878-92	20.1	20
57	DNA polymerase lambda directly binds to proliferating cell nuclear antigen through its confined C-terminal region. <i>Genes To Cells</i> , 2005 , 10, 705-15	2.3	23
56	Microtubule bundle formation and cell death induced by the human CLASP/Orbit N-terminal fragment. <i>Cell Structure and Function</i> , 2005 , 30, 7-13	2.2	5
55	Two functional coding single nucleotide polymorphisms in STK15 (Aurora-A) coordinately increase esophageal cancer risk. <i>Cancer Research</i> , 2005 , 65, 3548-54	10.1	59
54	The centrosomal protein Lats2 is a phosphorylation target of Aurora-A kinase. <i>Genes To Cells</i> , 2004 , 9, 383-97	2.3	125
53	Regulation of checkpoint kinases through dynamic interaction with Crb2. <i>EMBO Journal</i> , 2004 , 23, 418-28	3.3	47
52	Amino-terminal domain of ATRIP contributes to intranuclear relocation of the ATR-ATRIP complex following DNA damage. <i>FEBS Letters</i> , 2004 , 577, 289-93	3.8	19
51	Generation and application of phospho-specific antibodies for p53 and pRB. <i>Methods in Molecular Biology</i> , 2003 , 223, 17-26	1.4	6
50	Terminal deoxynucleotidyltransferase forms a ternary complex with a novel chromatin remodeling protein with 82 kDa and core histone. <i>Genes To Cells</i> , 2003 , 8, 559-71	2.3	13
49	NBS1 localizes to gamma-H2AX foci through interaction with the FHA/BRCT domain. <i>Current Biology</i> , 2002 , 12, 1846-51	6.3	250
48	Enzyme-linked immunosorbent assay for distinct cyclin-dependent kinase activities using phosphorylation-site-specific anti-pRB monoclonal antibodies. <i>Analytical Biochemistry</i> , 2002 , 301, 65-74	3.1	5
47	Over-expression of human DNA polymerase lambda in E. coli and characterization of the recombinant enzyme. <i>Genes To Cells</i> , 2002 , 7, 639-51	2.3	96
46	Activation of cyclin D1-kinase in murine fibroblasts lacking both p21(Cip1) and p27(Kip1). <i>Oncogene</i> , 2002 , 21, 8067-74	9.2	65
45	Identification of X-linked inhibitor of apoptosis-associated factor-1 as an interferon-stimulated gene that augments TRAIL Apo2L-induced apoptosis. <i>Journal of Biological Chemistry</i> , 2002 , 277, 28504-11	5.4	129
44	Terminal deoxynucleotidyltransferase directly interacts with a novel nuclear protein that is homologous to p65. <i>Genes To Cells</i> , 2001 , 6, 641-52	2.3	24

43	Terminal deoxynucleotidyltransferase is negatively regulated by direct interaction with proliferating cell nuclear antigen. <i>Genes To Cells</i> , 2001 , 6, 815-24	2.3	26
42	Ectopic expression of BEAF32A in the Drosophila eye imaginal disc inhibits differentiation of photoreceptor cells and induces apoptosis. <i>Chromosoma</i> , 2001 , 110, 313-21	2.8	11
41	Identification of XAF1 as an antagonist of XIAP anti-Caspase activity. <i>Nature Cell Biology</i> , 2001 , 3, 128-33	3.4	346
40	Stimulation of DNA polymerase alpha activity by cdk2-phosphorylated Rb protein. <i>Biochemical and Biophysical Research Communications</i> , 2001 , 282, 984-90	3.4	9
39	Replication of the rat aldolase B locus differs between aldolase B-expressing and non-expressing cells. <i>FEBS Letters</i> , 2001 , 505, 332-6	3.8	5
38	Pim-1 translocates sorting nexin 6/TRAF4-associated factor 2 from cytoplasm to nucleus. <i>FEBS Letters</i> , 2001 , 506, 33-8	3.8	41
37	ORC1 interacts with c-Myc to inhibit E-box-dependent transcription by abrogating c-Myc-SNF5/INI1 interaction. <i>Genes To Cells</i> , 2000 , 5, 481-90	2.3	34
36	PAP-1, a novel target protein of phosphorylation by pim-1 kinase. <i>FEBS Journal</i> , 2000 , 267, 5168-78		50
35	c-Myb acetylation at the carboxyl-terminal conserved domain by transcriptional co-activator p300. <i>Oncogene</i> , 2000 , 19, 444-51	9.2	109
34	Ataxia telangiectasia-mutated phosphorylates Chk2 in vivo and in vitro. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000 , 97, 10389-94	11.5	692
33	Human Cdc7-related kinase complex. In vitro phosphorylation of MCM by concerted actions of Cdks and Cdc7 and that of a critical threonine residue of Cdc7 by Cdk. <i>Journal of Biological Chemistry</i> , 2000 , 275, 29042-52	5.4	125
32	Multiple Ras-dependent phosphorylation pathways regulate Myc protein stability. <i>Genes and Development</i> , 2000 , 14, 2501-14	12.6	892
31	p53AIP1, a potential mediator of p53-dependent apoptosis, and its regulation by Ser-46-phosphorylated p53. <i>Cell</i> , 2000 , 102, 849-62	56.2	1005
30	Chk1 is an essential kinase that is regulated by Atr and required for the G2/M DNA damage checkpoint. <i>Genes and Development</i> , 2000 , 14, 1448-1459	12.6	769
29	The human homologs of checkpoint kinases Chk1 and Cds1 (Chk2) phosphorylate p53 at multiple DNA damage-inducible sites. <i>Genes and Development</i> , 2000 , 14, 289-300	12.6	470
28	Establishment of and recovery from damage checkpoint requires sequential interactions of Crb2 with protein kinases Rad3, Chk1, and Cdc2. <i>Cold Spring Harbor Symposia on Quantitative Biology</i> , 2000 , 65, 443-9	3.9	7
27	Cell cycle-dependent switch of up-and down-regulation of human hsp70 gene expression by interaction between c-Myc and CBF/NF-Y. <i>Journal of Biological Chemistry</i> , 1999 , 274, 24270-9	5.4	57
26	XIAP, a cellular member of the inhibitor of apoptosis protein family, links the receptors to TAB1-TAK1 in the BMP signaling pathway. <i>EMBO Journal</i> , 1999 , 18, 179-87	13	291

25	Immuno-histochemical detection of human telomerase catalytic component, hTERT, in human colorectal tumor and non-tumor tissue sections. <i>Oncogene</i> , 1999 , 18, 1561-7	9.2	145
24	Requirement of ATM in phosphorylation of the human p53 protein at serine 15 following DNA double-strand breaks. <i>Molecular and Cellular Biology</i> , 1999 , 19, 2828-34	4.8	110
23	Assembly of the Epstein-Barr virus BBLF4, BSLF1 and BBLF2/3 proteins and their interactive properties. <i>Journal of General Virology</i> , 1999 , 80 (Pt 11), 2879-2887	4.9	26
22	AMY-1, a novel C-MYC binding protein that stimulates transcription activity of C-MYC. <i>Genes To Cells</i> , 1998 , 3, 549-65	2.3	64
21	Activation of the ATM kinase by ionizing radiation and phosphorylation of p53. <i>Science</i> , 1998 , 281, 1677-9	9.3	1551
20	The second-largest subunit of the mouse DNA polymerase alpha-primase complex facilitates both production and nuclear translocation of the catalytic subunit of DNA polymerase alpha. <i>Molecular and Cellular Biology</i> , 1998 , 18, 3552-62	4.8	39
19	The inhibitory effect of novel triterpenoid compounds, fomitelic acids, on DNA polymerase beta. <i>Biochemical Journal</i> , 1998 , 330 (Pt 3), 1325-32	3.8	54
18	Overexpression of 3' untranslated region of the myotonic dystrophy kinase cDNA inhibits myoblast differentiation in vitro. <i>Journal of Biological Chemistry</i> , 1997 , 272, 29626-35	5.4	34
17	p53 is phosphorylated by CDK7-cyclin H in a p36MAT1-dependent manner. <i>Molecular and Cellular Biology</i> , 1997 , 17, 7220-9	4.8	150
16	DJ-1, a novel oncogene which transforms mouse NIH3T3 cells in cooperation with ras. <i>Biochemical and Biophysical Research Communications</i> , 1997 , 231, 509-13	3.4	633
15	Elevation of neuronal expression of NAIP reduces ischemic damage in the rat hippocampus. <i>Nature Medicine</i> , 1997 , 3, 997-1004	50.5	243
14	Cdk4-Cyclin D1 and Cdk2-Cyclin E/A Phosphorylate Different Sites in the RB Protein 1997 , 229-231		1
13	Epstein-Barr virus single-stranded DNA-binding protein: purification, characterization, and action on DNA synthesis by the viral DNA polymerase. <i>Virology</i> , 1996 , 222, 352-64	3.6	29
12	Suppression of apoptosis in mammalian cells by NAIP and a related family of IAP genes. <i>Nature</i> , 1996 , 379, 349-53	50.4	917
11	Investigation of myotonic dystrophy kinase isoform translocation and membrane association. <i>Journal of Biological Chemistry</i> , 1996 , 271, 15187-93	5.4	27
10	Characterization of myotonic dystrophy kinase (DMK) protein in human and rodent muscle and central nervous tissue. <i>Human Molecular Genetics</i> , 1995 , 4, 1063-72	5.6	60
9	The gene for neuronal apoptosis inhibitory protein is partially deleted in individuals with spinal muscular atrophy. <i>Cell</i> , 1995 , 80, 167-78	56.2	884
8	Molecular cloning of a human protein that binds to the retinoblastoma protein and chromosomal mapping. <i>Genomics</i> , 1995 , 27, 511-9	4.3	31

7	Expression patterns of DNA replication enzymes and the regulatory factor DREF during <i>Drosophila</i> development analyzed with specific antibodies. <i>Biology of the Cell</i> , 1995 , 85, 147-55	3-5	25
6	Intermediate form of mucopolysaccharidosis type II (Hunter disease): a C1327 to T substitution in the iduronate sulfatase gene. <i>Biochemical and Biophysical Research Communications</i> , 1992 , 183, 809-13	3-4	28
5	Efficient purification of a full length and biochemically active p110Rb, the retinoblastoma gene product. <i>Biochemical and Biophysical Research Communications</i> , 1992 , 187, 697-702	3-4	10
4	Defect of a fiber cell-specific 94-kDa protein in the lens of inherited microphthalmic mutant mouse Elo. <i>Biochemical and Biophysical Research Communications</i> , 1991 , 179, 1175-80	3-4	15
3	A novel stimulating protein of mammalian DNA polymerase alpha. <i>Journal of Biochemistry</i> , 1989 , 106, 389-95	3-1	7
2	Structural study of immunoaffinity-purified DNA polymerase alpha-DNA primase complex from calf thymus. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1988 , 950, 263-73		108
1	Characterization of a Mr = 56,000 polypeptide associated with 10S DNA polymerase alpha purified from calf thymus using monoclonal antibody. <i>Nucleic Acids Research</i> , 1985 , 13, 6635-49	20-1	4