

# An Verrijken

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9143567/publications.pdf>

Version: 2024-02-01

18  
papers

1,799  
citations

687363

13  
h-index

839539

18  
g-index

19  
all docs

19  
docs citations

19  
times ranked

3393  
citing authors

#	ARTICLE	IF	CITATIONS
1	Association of Non-alcoholic Fatty Liver Disease with Chronic Kidney Disease: A Systematic Review and Meta-analysis. <i>PLoS Medicine</i> , 2014, 11, e1001680.	8.4	507
2	PPAR $\alpha$ gene expression correlates with severity and histological treatment response in patients with non-alcoholic steatohepatitis. <i>Journal of Hepatology</i> , 2015, 63, 164-173.	3.7	270
3	Association of Adipose Tissue Inflammation With Histologic Severity of Nonalcoholic Fatty Liver Disease. <i>Gastroenterology</i> , 2015, 149, 635-648.e14.	1.3	249
4	Prothrombotic factors in histologically proven nonalcoholic fatty liver disease and nonalcoholic steatohepatitis. <i>Hepatology</i> , 2014, 59, 121-129.	7.3	141
5	Noninvasive Detection of Nonalcoholic Steatohepatitis Using Clinical Markers and Circulating Levels of Lipids and Metabolites. <i>Clinical Gastroenterology and Hepatology</i> , 2016, 14, 1463-1472.e6.	4.4	120
6	Transcriptional network analysis implicates altered hepatic immune function in NASH development and resolution. <i>Nature Metabolism</i> , 2019, 1, 604-614.	11.9	102
7	C-reactive protein levels in relation to various features of non-alcoholic fatty liver disease among obese patients. <i>Journal of Hepatology</i> , 2011, 55, 660-665.	3.7	98
8	Bile Acid Alterations Are Associated With Insulin Resistance, but Not With NASH, in Obese Subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, 3783-3794.	3.6	78
9	Muscle fat content is strongly associated with NASH: A longitudinal study in patients with morbid obesity. <i>Journal of Hepatology</i> , 2021, 75, 292-301.	3.7	68
10	Pro-Inflammatory Cytokines but Not Endotoxin-Related Parameters Associate with Disease Severity in Patients with NAFLD. <i>PLoS ONE</i> , 2016, 11, e0166048.	2.5	52
11	Plasma BCAA Changes in Patients With NAFLD Are Sex Dependent. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, 2311-2321.	3.6	39
12	NASH-related increases in plasma bile acid levels depend on insulin resistance. <i>JHEP Reports</i> , 2021, 3, 100222.	4.9	24
13	Malnutrition and its relation with diabetic foot ulcer severity and outcome: a review. <i>Acta Clinica Belgica</i> , 2022, 77, 79-85.	1.2	18
14	Malnutrition according to the 2018 GLIM criteria is highly prevalent in people with a diabetic foot ulcer but does not affect outcome. <i>Clinical Nutrition ESPEN</i> , 2021, 43, 335-341.	1.2	10
15	Posttranscriptional Regulation of the Human LDL Receptor by the U2-Spliceosome. <i>Circulation Research</i> , 2022, 130, 80-95.	4.5	9
16	Coronary artery calcifications and diastolic dysfunction versus visceral fat area in type 1 diabetes: VISCERA study. <i>Journal of Diabetes and Its Complications</i> , 2018, 32, 271-278.	2.3	8
17	DNA sequencing and copy number variation analysis of MCHR2 in a cohort of Prader Willi like (PWL) patients. <i>Obesity Research and Clinical Practice</i> , 2018, 12, 158-166.	1.8	2
18	Reply. <i>Hepatology</i> , 2014, 60, 1451-1451.	7.3	0