

# Xia Liu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9144053/publications.pdf>

Version: 2024-02-01

19  
papers

662  
citations

567281

15  
h-index

752698

20  
g-index

20  
all docs

20  
docs citations

20  
times ranked

1056  
citing authors

#	ARTICLE	IF	CITATIONS
1	Expression analysis of histone acetyltransferases in rice under drought stress. <i>Biochemical and Biophysical Research Communications</i> , 2014, 443, 400-405.	2.1	88
2	Wnt3a: functions and implications in cancer. <i>Chinese Journal of Cancer</i> , 2015, 34, 554-62.	4.9	72
3	Upregulation of NEK2 is associated with drug resistance in ovarian cancer. <i>Oncology Reports</i> , 2014, 31, 745-754.	2.6	62
4	Litchi seed extracts diminish prostate cancer progression via induction of apoptosis and attenuation of EMT through Akt/GSK-3 $\beta$ signaling. <i>Scientific Reports</i> , 2017, 7, 41656.	3.3	58
5	Tumor suppressor genes associated with drug resistance in ovarian cancer (Review). <i>Oncology Reports</i> , 2013, 30, 3-10.	2.6	50
6	Oncogenes associated with drug resistance in ovarian cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2015, 141, 381-395.	2.5	41
7	Cross-validation of genes potentially associated with overall survival and drug resistance in ovarian cancer. <i>Oncology Reports</i> , 2017, 37, 3084-3092.	2.6	35
8	Discovery of microarray-identified genes associated with ovarian cancer progression. <i>International Journal of Oncology</i> , 2015, 46, 2467-2478.	3.3	34
9	Microarray-based identification of genes associated with cancer progression and prognosis in hepatocellular carcinoma. <i>Journal of Experimental and Clinical Cancer Research</i> , 2016, 35, 127.	8.6	33
10	Upregulation of E2F transcription factor 3 is associated with poor prognosis in hepatocellular carcinoma. <i>Oncology Reports</i> , 2014, 31, 1139-1146.	2.6	30
11	Microarray-based identification of genes associated with prognosis and drug resistance in ovarian cancer. <i>Journal of Cellular Biochemistry</i> , 2019, 120, 6057-6070.	2.6	26
12	Downregulation of tumor suppressor gene ribonuclease T2 and gametogenetin binding protein 2 is associated with drug resistance in ovarian cancer. <i>Oncology Reports</i> , 2014, 32, 362-372.	2.6	22
13	Downregulation of transient receptor potential cation channel, subfamily C, member 1 contributes to drug resistance and high histological grade in ovarian cancer. <i>International Journal of Oncology</i> , 2016, 48, 243-252.	3.3	21
14	Associations of tumor suppressor SPARCL1 with cancer progression and prognosis. <i>Oncology Letters</i> , 2017, 14, 2603-2610.	1.8	21
15	Downregulation of NEK11 is associated with drug resistance in ovarian cancer. <i>International Journal of Oncology</i> , 2014, 45, 1266-1274.	3.3	20
16	NCALD affects drug resistance and prognosis by acting as a ceRNA of CX3CL1 in ovarian cancer. <i>Journal of Cellular Biochemistry</i> , 2020, 121, 4470-4483.	2.6	16
17	Low expression of KCNN3 may affect drug resistance in ovarian cancer. <i>Molecular Medicine Reports</i> , 2018, 18, 1377-1386.	2.4	13
18	Big Data-Based Identification of Multi-Gene Prognostic Signatures in Liver Cancer. <i>Frontiers in Oncology</i> , 2020, 10, 847.	2.8	10

#	ARTICLE	IF	CITATIONS
19	Low Expression of SLC7A11 Confers Drug Resistance and Worse Survival in Ovarian Cancer via Inhibition of Cell Autophagy as a Competing Endogenous RNA. <i>Frontiers in Oncology</i> , 2021, 11, 744940.	2.8	9