

Xia Liu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9144053/publications.pdf>

Version: 2024-02-01

19
papers

662
citations

567281

15
h-index

752698

20
g-index

20
all docs

20
docs citations

20
times ranked

1056
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Low Expression of SLC7A11 Confers Drug Resistance and Worse Survival in Ovarian Cancer via Inhibition of Cell Autophagy as a Competing Endogenous RNA. <i>Frontiers in Oncology</i> , 2021, 11, 744940. | 2.8 | 9 |
| 2 | Big Data-Based Identification of Multi-Gene Prognostic Signatures in Liver Cancer. <i>Frontiers in Oncology</i> , 2020, 10, 847. | 2.8 | 10 |
| 3 | NCALD affects drug resistance and prognosis by acting as a ceRNA of CX3CL1 in ovarian cancer. <i>Journal of Cellular Biochemistry</i> , 2020, 121, 4470-4483. | 2.6 | 16 |
| 4 | Microarray-based identification of genes associated with prognosis and drug resistance in ovarian cancer. <i>Journal of Cellular Biochemistry</i> , 2019, 120, 6057-6070. | 2.6 | 26 |
| 5 | Low expression of KCNN3 may affect drug resistance in ovarian cancer. <i>Molecular Medicine Reports</i> , 2018, 18, 1377-1386. | 2.4 | 13 |
| 6 | Litchi seed extracts diminish prostate cancer progression via induction of apoptosis and attenuation of EMT through Akt/GSK-3 β signaling. <i>Scientific Reports</i> , 2017, 7, 41656. | 3.3 | 58 |
| 7 | Cross-validation of genes potentially associated with overall survival and drug resistance in ovarian cancer. <i>Oncology Reports</i> , 2017, 37, 3084-3092. | 2.6 | 35 |
| 8 | Associations of tumor suppressor SPARCL1 with cancer progression and prognosis. <i>Oncology Letters</i> , 2017, 14, 2603-2610. | 1.8 | 21 |
| 9 | Microarray-based identification of genes associated with cancer progression and prognosis in hepatocellular carcinoma. <i>Journal of Experimental and Clinical Cancer Research</i> , 2016, 35, 127. | 8.6 | 33 |
| 10 | Downregulation of transient receptor potential cation channel, subfamily C, member 1 contributes to drug resistance and high histological grade in ovarian cancer. <i>International Journal of Oncology</i> , 2016, 48, 243-252. | 3.3 | 21 |
| 11 | Wnt3a: functions and implications in cancer. <i>Chinese Journal of Cancer</i> , 2015, 34, 554-62. | 4.9 | 72 |
| 12 | Discovery of microarray-identified genes associated with ovarian cancer progression. <i>International Journal of Oncology</i> , 2015, 46, 2467-2478. | 3.3 | 34 |
| 13 | Oncogenes associated with drug resistance in ovarian cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2015, 141, 381-395. | 2.5 | 41 |
| 14 | Expression analysis of histone acetyltransferases in rice under drought stress. <i>Biochemical and Biophysical Research Communications</i> , 2014, 443, 400-405. | 2.1 | 88 |
| 15 | Upregulation of NEK2 is associated with drug resistance in ovarian cancer. <i>Oncology Reports</i> , 2014, 31, 745-754. | 2.6 | 62 |
| 16 | Upregulation of E2F transcription factor 3 is associated with poor prognosis in hepatocellular carcinoma. <i>Oncology Reports</i> , 2014, 31, 1139-1146. | 2.6 | 30 |
| 17 | Downregulation of tumor suppressor gene ribonuclease T2 and gametogenetin binding protein 2 is associated with drug resistance in ovarian cancer. <i>Oncology Reports</i> , 2014, 32, 362-372. | 2.6 | 22 |
| 18 | Downregulation of NEK11 is associated with drug resistance in ovarian cancer. <i>International Journal of Oncology</i> , 2014, 45, 1266-1274. | 3.3 | 20 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Tumor suppressor genes associated with drug resistance in ovarian cancer (Review). <i>Oncology Reports</i> , 2013, 30, 3-10. | 2.6 | 50 |