

Howard J Hoffman

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9146638/howard-j-hoffman-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

87
papers

10,160
citations

37
h-index

92
g-index

92
ext. papers

11,686
ext. citations

5.5
avg, IF

5.33
L-index

#	Paper	IF	Citations
87	Gap detection responses modelled using the Hill equation in adults with well-controlled HIV.. <i>International Journal of Audiology</i> , 2022 , 1-10	2.6	
86	Hearing loss prevalence and years lived with disability, 1990-2019: findings from the Global Burden of Disease Study 2019. <i>Lancet, The</i> , 2021 , 397, 996-1009	4.0	82
85	Prenatal Exposure to Tobacco and Alcohol Alters Development of the Neonatal Auditory System. <i>Developmental Neuroscience</i> , 2021 , 43, 358-375	2.2	
84	Association of Prenatal Exposure to Maternal Drinking and Smoking With the Risk of Stillbirth. <i>JAMA Network Open</i> , 2021 , 4, e2121726	10.4	3
83	Associations between chronic cigarette smoking and taste function: Results from the 2013-2014 national health and nutrition examination survey. <i>Physiology and Behavior</i> , 2021 , 240, 113554	3.5	1
82	Concurrent prenatal drinking and smoking increases risk for SIDS: Safe Passage Study report. <i>EClinicalMedicine</i> , 2020 , 19, 100247	11.3	24
81	Words-in-Noise Test Performance in Young Adults Perinatally HIV Infected and Exposed, Uninfected. <i>American Journal of Audiology</i> , 2020 , 29, 68-78	1.8	0
80	Vestibular and oculomotor abnormalities among HIV-infected and HIV-uninfected men and women: A pilot study. <i>Journal of Vestibular Research: Equilibrium and Orientation</i> , 2020 , 30, 329-334	2.5	0
79	Birth Prevalence of Congenital Cytomegalovirus Infection in HIV-Exposed Uninfected Children in the Era of Combination Antiretroviral Therapy. <i>Journal of Pediatrics</i> , 2020 , 216, 82-87.e2	3.6	3
78	Associations of olfactory dysfunction with anthropometric and cardiometabolic measures: Findings from the 2013-2014 national health and nutrition examination survey (NHANES). <i>Physiology and Behavior</i> , 2020 , 215, 112702	3.5	15
77	Association between vertigo, cognitive and psychiatric conditions in US children: 2012 National Health Interview Survey. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2020 , 130, 109802	1.7	7
76	Alcohol and Taste Intensity. <i>Chemosensory Perception</i> , 2019 , 12, 90-99	1.2	3
75	Poorer outcomes of all low birth weight groups at age 10: Missouri statewide case-control study. <i>Early Human Development</i> , 2019 , 136, 60-69	2.2	3
74	Genome-wide association meta-analysis identifies five novel loci for age-related hearing impairment. <i>Scientific Reports</i> , 2019 , 9, 15192	4.9	14
73	Kids Nowadays Hear Better Than We Did: Declining Prevalence of Hearing Loss in US Youth, 1966-2010. <i>Laryngoscope</i> , 2019 , 129, 1922-1939	3.6	9
72	Chronic Cigarette Smoking Associates Directly and Indirectly with Self-Reported Olfactory Alterations: Analysis of the 2011-2014 National Health and Nutrition Examination Survey. <i>Nicotine and Tobacco Research</i> , 2019 , 21, 818-827	4.9	64
71	Risk for Speech and Language Impairments in Preschool Age HIV-exposed Uninfected Children With In Utero Combination Antiretroviral Exposure. <i>Pediatric Infectious Disease Journal</i> , 2018 , 37, 678-685	3.4	12

70	Hearing Disability Prevalence and Risk Factors in Two Recent National Surveys. <i>American Journal of Preventive Medicine</i> , 2018 , 55, 326-335	6.1	7
69	Childhood hearing loss and established risk factors: What is the contribution of tobacco exposure prenatally or after birth?. <i>Paediatric and Perinatal Epidemiology</i> , 2018 , 32, 439-441	2.7	
68	Transient Otoacoustic Emissions and Auditory Brainstem Responses in Low-Risk Cohort of Newborn and One-Month-Old Infants: Assessment of Infant Auditory System Physiology in the Prenatal Alcohol in SIDS and Stillbirth Network Safe Passage Study. <i>Journal of the American Academy of Audiology</i> , 2018 , 29, 748-763	1.3	2
67	Structural equation modeling of associations among taste-related risk factors, taste functioning, and adiposity. <i>Obesity</i> , 2017 , 25, 781-787	8	17
66	The Reduction in the Age-Adjusted Prevalence of Hearing Impairment in the United States-Reply. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2017 , 143, 957-958	3.9	1
65	Declining Prevalence of Hearing Loss in US Adults Aged 20 to 69 Years. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2017 , 143, 274-285	3.9	149
64	Vital Signs: Noise-Induced Hearing Loss Among Adults - United States 2011-2012. <i>Morbidity and Mortality Weekly Report</i> , 2017 , 66, 139-144	31.7	63
63	Drinking and smoking patterns during pregnancy: Development of group-based trajectories in the Safe Passage Study. <i>Alcohol</i> , 2017 , 62, 49-60	2.7	30
62	Occupational noise exposure, hearing loss, and notched audiograms in the HUNT Nord-Trøndelag hearing loss study, 1996-1998. <i>Laryngoscope</i> , 2017 , 127, 1442-1450	3.6	14
61	Racial and Ethnic Differences in Receipt of Pressure Equalization Tubes Among US Children, 2014. <i>Academic Pediatrics</i> , 2017 , 17, 88-94	2.7	10
60	Childhood otitis media is associated with dizziness in adulthood: the HUNT cohort study. <i>European Archives of Oto-Rhino-Laryngology</i> , 2016 , 273, 2047-54	3.5	16
59	Prevalence and Risk Factors of Self-Reported Smell and Taste Alterations: Results from the 2011-2012 US National Health and Nutrition Examination Survey (NHANES). <i>Chemical Senses</i> , 2016 , 41, 69-76	4.8	99
58	Vestibular vertigo and comorbid cognitive and psychiatric impairment: the 2008 National Health Interview Survey. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2016 , 87, 367-72	5.5	69
57	Sleep Apnea Is Associated with Hearing Impairment: The Hispanic Community Health Study/Study of Latinos. <i>Journal of Clinical Sleep Medicine</i> , 2016 , 12, 719-26	3.1	18
56	New chemosensory component in the U.S. National Health and Nutrition Examination Survey (NHANES): first-year results for measured olfactory dysfunction. <i>Reviews in Endocrine and Metabolic Disorders</i> , 2016 , 17, 221-40	10.5	118
55	Epidemiology of Dizziness and Balance Problems in Children in the United States: A Population-Based Study. <i>Journal of Pediatrics</i> , 2016 , 171, 240-7.e1-3	3.6	54
54	Hearing in older adults with exfoliation syndrome/exfoliation glaucoma or primary open-angle glaucoma. <i>Acta Ophthalmologica</i> , 2016 , 94, 140-6	3.7	6
53	Longitudinal Evaluation of Language Impairment in Youth With Perinatally Acquired Human Immunodeficiency Virus (HIV) and Youth With Perinatal HIV Exposure. <i>Journal of the Pediatric Infectious Diseases Society</i> , 2016 , 5, S33-S40	4.8	11

52	Impact of Sensory Impairments on Functional Disability in Adults With Arthritis. <i>American Journal of Preventive Medicine</i> , 2016 , 50, 454-462	6.1	4
51	Risk Factors for Hearing Impairment among Adults with Diabetes: The Hispanic Community Health Study/Study of Latinos (HCHS/SOL). <i>Journal of Clinical and Translational Endocrinology</i> , 2016 , 6, 15-22	2.4	5
50	Speech audiometry findings from HIV+ and HIV- adults in the MACS and WIHS longitudinal cohort studies. <i>Journal of Communication Disorders</i> , 2016 , 64, 103-109	1.9	4
49	Sex-specific predictors of hearing-aid use in older persons: The age, gene/environment susceptibility - Reykjavik study. <i>International Journal of Audiology</i> , 2015 , 54, 634-41	2.6	20
48	Hearing loss among HIV-seropositive and HIV-seronegative men and women. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2015 , 141, 202-10	3.9	24
47	Distortion product otoacoustic emission data in perinatally HIV-infected and HIV-exposed but uninfected children and adolescents in the Pediatric HIV/AIDS Cohort Study. <i>Pediatric Infectious Disease Journal</i> , 2015 , 34, 276-8	3.4	6
46	Hearing Impairment Prevalence and Associated Risk Factors in the Hispanic Community Health Study/Study of Latinos. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2015 , 141, 641-8	3.9	49
45	The Taste and Smell Protocol in the 2011-2014 US National Health and Nutrition Examination Survey (NHANES): Test-Retest Reliability and Validity Testing. <i>Chemosensory Perception</i> , 2015 , 8, 138-148	1.2	73
44	Association Between Childhood Hearing Disorders and Tinnitus in Adulthood. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2015 , 141, 983-9	3.9	8
43	Statistical Modeling Approaches with Taste Functioning and Central Adiposity in Women: Implications for Mining the New National Health and Nutrition Examination Survey (NHANES) Taste and Smell Protocol. <i>FASEB Journal</i> , 2015 , 29, 602.4	0.9	
42	Self-reported Hearing Trouble in Adults Aged 18 and Over: United States, 2014. <i>NCHS Data Brief</i> , 2015 , 1-8	14.7	9
41	Sensitivity and Specificity of Self-Reported Olfactory Function in a Home-Based Study of Independent-Living, Healthy Older Women. <i>Chemosensory Perception</i> , 2014 , 7, 108-116	1.2	29
40	Breastfeeding and risk of infections at 6 years. <i>Pediatrics</i> , 2014 , 134 Suppl 1, S13-20	7.4	48
39	Hearing impairment associated with depression in US adults, National Health and Nutrition Examination Survey 2005-2010. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2014 , 140, 293-302	3.9	186
38	Cochlear function among HIV-seropositive and HIV-seronegative men and women. <i>Ear and Hearing</i> , 2014 , 35, 56-62	3.4	10
37	Otoacoustic emissions, pure-tone audiometry, and self-reported hearing. <i>International Journal of Audiology</i> , 2013 , 52, 74-82	2.6	18
36	Prevalence and impact of bilateral vestibular hypofunction: results from the 2008 US National Health Interview Survey. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2013 , 139, 803-10	3.9	117
35	Gustation assessment using the NIH Toolbox. <i>Neurology</i> , 2013 , 80, S20-4	6.5	135

34	Olfactory assessment using the NIH Toolbox. <i>Neurology</i> , 2013 , 80, S32-6	6.5	53
33	Panel 1: Epidemiology, natural history, and risk factors. <i>Otolaryngology - Head and Neck Surgery</i> , 2013 , 148, E1-E25	5.5	28
32	Evaluation of risk for late language emergence after in utero antiretroviral drug exposure in HIV-exposed uninfected infants. <i>Pediatric Infectious Disease Journal</i> , 2013 , 32, e406-13	3.4	41
31	Common values in assessing health outcomes from disease and injury: disability weights measurement study for the Global Burden of Disease Study 2010. <i>Lancet, The</i> , 2012 , 380, 2129-43	4.0	842
30	Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. <i>Lancet, The</i> , 2012 , 380, 2197-223	4.0	5768
29	Taste damage (otitis media, tonsillectomy and head and neck cancer), oral sensations and BMI. <i>Physiology and Behavior</i> , 2012 , 107, 516-26	3.5	67
28	Prevalence of abnormalities in vestibular function and balance among HIV-seropositive and HIV-seronegative women and men. <i>PLoS ONE</i> , 2012 , 7, e38419	3.7	18
27	Hearing threshold levels at age 70 years (65-74 years) in the unscreened older adult population of the United States, 1959-1962 and 1999-2006. <i>Ear and Hearing</i> , 2012 , 33, 437-40	3.4	32
26	Occupation and the risk of bothersome tinnitus: results from a prospective cohort study (HUNT). <i>BMJ Open</i> , 2012 , 2, e000512	3	26
25	Hearing loss in perinatally HIV-infected and HIV-exposed but uninfected children and adolescents. <i>Pediatric Infectious Disease Journal</i> , 2012 , 31, 835-41	3.4	37
24	Language impairment in children perinatally infected with HIV compared to children who were HIV-exposed and uninfected. <i>Journal of Developmental and Behavioral Pediatrics</i> , 2012 , 33, 112-23	2.4	33
23	The modified Romberg Balance Test: normative data in U.S. adults. <i>Otology and Neurotology</i> , 2011 , 32, 1309-11	2.6	76
22	Risk factors for hearing impairment among U.S. adults with diabetes: National Health and Nutrition Examination Survey 1999-2004. <i>Diabetes Care</i> , 2011 , 34, 1540-5	14.6	57
21	Low heritability of tinnitus: results from the second Nord-Trøndelag health study. <i>JAMA Otolaryngology</i> , 2010 , 136, 178-82		37
20	Americans hear as well or better today compared with 40 years ago: hearing threshold levels in the unscreened adult population of the United States, 1959-1962 and 1999-2004. <i>Ear and Hearing</i> , 2010 , 31, 725-34	3.4	63
19	Perspectives on population-based epidemiological studies of olfactory and taste impairment. <i>Annals of the New York Academy of Sciences</i> , 2009 , 1170, 514-30	6.5	54
18	Symposium overview: Epidemiological studies of taste and smell. <i>Annals of the New York Academy of Sciences</i> , 2009 , 1170, 511-3	6.5	1
17	Diabetes and hearing impairment in the United States: audiometric evidence from the National Health and Nutrition Examination Survey, 1999 to 2004. <i>Annals of Internal Medicine</i> , 2008 , 149, 1-10	8	239

16	Use of a dummy (pacifier) during sleep and risk of sudden infant death syndrome (SIDS): population based case-control study. <i>BMJ, The</i> , 2006 , 332, 18-22	5.9	75
15	Hearing loss induced by occupational and impulse noise: results on threshold shifts by frequencies, age and gender from the Nord-Trøndelag Hearing Loss Study. <i>International Journal of Audiology</i> , 2006 , 45, 309-17	2.6	45
14	Screened and unscreened hearing threshold levels for the adult population: results from the Nord-Trøndelag Hearing Loss Study. <i>International Journal of Audiology</i> , 2005 , 44, 213-30	2.6	77
13	Recent advances in otitis media. 1. Epidemiology, natural history, and risk factors. <i>Annals of Otolaryngology, Rhinology and Laryngology</i> , 2005 , 114, 8-15	2.1	29
12	Otoacoustic emissions in the general adult population of Nord-Trøndelag, Norway: III. Relationships with pure-tone hearing thresholds. <i>International Journal of Audiology</i> , 2005 , 44, 15-23	2.6	6
11	From psychophysics to the clinic: missteps and advances. <i>Food Quality and Preference</i> , 2004 , 15, 617-632	5.8	56
10	Racial/ethnic disparities in the diagnosis of otitis media in infancy. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2004 , 68, 795-804	1.7	44
9	Hearing loss associated with ear infections in Nord-Trøndelag, Norway. <i>Ear and Hearing</i> , 2004 , 25, 388-96	3.4	14
8	Hearing loss induced by noise, ear infections, and head injuries: results from the Nord-Trøndelag Hearing Loss Study. <i>International Journal of Audiology</i> , 2003 , 42, 89-105	2.6	51
7	Genetic and environmental influences on self-reported reduced hearing in the old and oldest old. <i>Journal of the American Geriatrics Society</i> , 2001 , 49, 1512-7	5.6	95
6	Age-related changes in the prevalence of smell/taste problems among the United States adult population. Results of the 1994 disability supplement to the National Health Interview Survey (NHIS). <i>Annals of the New York Academy of Sciences</i> , 1998 , 855, 716-22	6.5	187
5	Home apnea monitoring and sudden infant death syndrome. <i>Preventive Medicine</i> , 1996 , 25, 645-9	4.3	14
4	Exercise, employment, other daily activities, and adverse pregnancy outcomes. <i>American Journal of Epidemiology</i> , 1996 , 143, 211-8	3.8	54
3	Pre-pregnancy risk factors of small-for-gestational age births among parous women in Scandinavia. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1993 , 72, 273-9	3.8	118
2	The tendency to repeat gestational age and birth weight in successive births, related to perinatal survival. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1983 , 62, 385-92	3.8	44
1	Obstetric service and perinatal mortality in Norway. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 1978 , 77, 3-19	3.8	29