

Shoichiro Tsugane

List of Publications by Citations

Source: <https://exaly.com/author-pdf/9146738/shoichiro-tsugane-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

917
papers

32,208
citations

85
h-index

128
g-index

983
ext. papers

37,427
ext. citations

5.2
avg, IF

6.98
L-index

#	Paper	IF	Citations
917	Association analysis identifies 65 new breast cancer risk loci. <i>Nature</i> , 2017 , 551, 92-94	50.4	643
916	Association between body-mass index and risk of death in more than 1 million Asians. <i>New England Journal of Medicine</i> , 2011 , 364, 719-29	59.2	616
915	Intake of fish and n3 fatty acids and risk of coronary heart disease among Japanese: the Japan Public Health Center-Based (JPHC) Study Cohort I. <i>Circulation</i> , 2006 , 113, 195-202	16.7	422
914	Diabetes mellitus and the risk of cancer: results from a large-scale population-based cohort study in Japan. <i>Archives of Internal Medicine</i> , 2006 , 166, 1871-7		396
913	Soy, isoflavones, and breast cancer risk in Japan. <i>Journal of the National Cancer Institute</i> , 2003 , 95, 906-13	7.7	315
912	Diet and the risk of gastric cancer: review of epidemiological evidence. <i>Gastric Cancer</i> , 2007 , 10, 75-83	7.6	310
911	What has made the population of Japan healthy?. <i>Lancet, The</i> , 2011 , 378, 1094-105	4.0	284
910	Decreased serum free testosterone in workers exposed to high levels of di-n-butyl phthalate (DBP) and di-2-ethylhexyl phthalate (DEHP): a cross-sectional study in China. <i>Environmental Health Perspectives</i> , 2006 , 114, 1643-8	8.4	272
909	Epidemiology of gastric cancer in Japan. <i>Postgraduate Medical Journal</i> , 2005 , 81, 419-24	2	245
908	Correlation between long-term survival in breast cancer patients and amplification of two putative oncogene-coamplification units: hst-1/int-2 and c-erbB-2/ear-1. <i>Cancer Research</i> , 1989 , 49, 3104-8	10.1	239
907	The JPHC study: design and some findings on the typical Japanese diet. <i>Japanese Journal of Clinical Oncology</i> , 2014 , 44, 777-82	2.8	237
906	Salt, salted food intake, and risk of gastric cancer: epidemiologic evidence. <i>Cancer Science</i> , 2005 , 96, 1-6	6.9	227
905	A randomized trial of surgery with and without chemotherapy for localized squamous carcinoma of the thoracic esophagus: the Japan Clinical Oncology Group Study. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 1997 , 114, 205-9	1.5	223
904	Genome-wide association study identifies 112 new loci for body mass index in the Japanese population. <i>Nature Genetics</i> , 2017 , 49, 1458-1467	36.3	214
903	Green tea consumption and prostate cancer risk in Japanese men: a prospective study. <i>American Journal of Epidemiology</i> , 2008 , 167, 71-7	3.8	205
902	Salt and salted food intake and subsequent risk of gastric cancer among middle-aged Japanese men and women. <i>British Journal of Cancer</i> , 2004 , 90, 128-34	8.7	205
901	Association of dietary intake of soy, beans, and isoflavones with risk of cerebral and myocardial infarctions in Japanese populations: the Japan Public Health Center-based (JPHC) study cohort I. <i>Circulation</i> , 2007 , 116, 2553-62	16.7	201

900	Baseline survey of JPHC study--design and participation rate. Japan Public Health Center-based Prospective Study on Cancer and Cardiovascular Diseases. <i>Journal of Epidemiology</i> , 2001 , 11, S24-9	3.4	196
899	Association between body mass index and cardiovascular disease mortality in east Asians and south Asians: pooled analysis of prospective data from the Asia Cohort Consortium. <i>BMJ, The</i> , 2013 , 347, f5446	5.9	188
898	Tumor thickness predicts cervical metastasis in patients with stage I/II carcinoma of the tongue. <i>Cancer</i> , 1998 , 82, 1443-8	6.4	185
897	Multi-step carcinogenesis model for adult T-cell leukemia. <i>Japanese Journal of Cancer Research</i> , 1989 , 80, 191-5		181
896	Soy product and isoflavone consumption in relation to prostate cancer in Japanese men. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2007 , 16, 538-45	4	165
895	Alcohol consumption and all-cause and cancer mortality among middle-aged Japanese men: seven-year follow-up of the JPHC study Cohort I. Japan Public Health Center. <i>American Journal of Epidemiology</i> , 1999 , 150, 1201-7	3.8	155
894	Rice intake and type 2 diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2010 , 92, 1468-77	7	154
893	Alcohol consumption and risk of stroke among middle-aged men: the JPHC Study Cohort I. <i>Stroke</i> , 2004 , 35, 1124-9	6.7	153
892	Gastric cancer screening and subsequent risk of gastric cancer: a large-scale population-based cohort study, with a 13-year follow-up in Japan. <i>International Journal of Cancer</i> , 2006 , 118, 2315-21	7.5	152
891	Influence of coffee drinking on subsequent risk of hepatocellular carcinoma: a prospective study in Japan. <i>Journal of the National Cancer Institute</i> , 2005 , 97, 293-300	9.7	152
890	Adult mortality attributable to preventable risk factors for non-communicable diseases and injuries in Japan: a comparative risk assessment. <i>PLoS Medicine</i> , 2012 , 9, e1001160	11.6	144
889	Validity and reproducibility of a self-administered food-frequency questionnaire to assess isoflavone intake in a Japanese population in comparison with dietary records and blood and urine isoflavones. <i>Journal of Nutrition</i> , 2001 , 131, 2741-7	4.1	135
888	Consumption of n-3 fatty acids and fish reduces risk of hepatocellular carcinoma. <i>Gastroenterology</i> , 2012 , 142, 1468-75	13.3	134
887	Plasma isoflavone level and subsequent risk of breast cancer among Japanese women: a nested case-control study from the Japan Public Health Center-based prospective study group. <i>Journal of Clinical Oncology</i> , 2008 , 26, 1677-83	2.2	133
886	Cigarette smoking and subsequent risk of lung cancer by histologic type in middle-aged Japanese men and women: the JPHC study. <i>International Journal of Cancer</i> , 2002 , 99, 245-51	7.5	130
885	Alcohol consumption and other risk factors for self-reported diabetes among middle-aged Japanese: a population-based prospective study in the JPHC study cohort I. <i>Diabetic Medicine</i> , 2005 , 22, 323-31	3.5	130
884	Insulin resistance and cancer: epidemiological evidence. <i>Cancer Science</i> , 2010 , 101, 1073-9	6.9	129
883	Correlation between histologic grade of malignancy and copy number of c-erbB-2 gene in breast carcinoma. A retrospective analysis of 176 cases. <i>Cancer</i> , 1990 , 65, 1794-800	6.4	129

882	Cigarette smoking, alcohol consumption and subsequent gastric cancer risk by subsite and histologic type. <i>International Journal of Cancer</i> , 2002 , 101, 560-6	7.5	127
881	Vegetables, fruit and risk of gastric cancer in Japan: a 10-year follow-up of the JPHC Study Cohort I. <i>International Journal of Cancer</i> , 2002 , 102, 39-44	7.5	127
880	Fruit and vegetable intake and risk of total cancer and cardiovascular disease: Japan Public Health Center-Based Prospective Study. <i>American Journal of Epidemiology</i> , 2008 , 167, 59-70	3.8	125
879	Major prognostic factors of adult patients with advanced T-cell lymphoma/leukemia. <i>Journal of Clinical Oncology</i> , 1988 , 6, 1088-97	2.2	125
878	Body mass index and diabetes in Asia: a cross-sectional pooled analysis of 900,000 individuals in the Asia cohort consortium. <i>PLoS ONE</i> , 2011 , 6, e19930	3.7	124
877	Daily total physical activity level and premature death in men and women: results from a large-scale population-based cohort study in Japan (JPHC study). <i>Annals of Epidemiology</i> , 2008 , 18, 522-30	6.4	123
876	Under- and overweight impact on mortality among middle-aged Japanese men and women: a 10-y follow-up of JPHC study cohort I. <i>International Journal of Obesity</i> , 2002 , 26, 529-37	5.5	122
875	Validity and reproducibility of a self-administered food frequency questionnaire in the JPHC Study Cohort II: study design, participant profile and results in comparison with Cohort I. <i>Journal of Epidemiology</i> , 2003 , 13, S134-47	3.4	121
874	Genome-wide association study in east Asians identifies novel susceptibility loci for breast cancer. <i>PLoS Genetics</i> , 2012 , 8, e1002532	6	118
873	Urinary bisphenol A and plasma hormone concentrations in male workers exposed to bisphenol A diglycidyl ether and mixed organic solvents. <i>Occupational and Environmental Medicine</i> , 2002 , 59, 625-8	2.1	118
872	Attributable causes of cancer in Japan in 2005--systematic assessment to estimate current burden of cancer attributable to known preventable risk factors in Japan. <i>Annals of Oncology</i> , 2012 , 23, 1362-1369	18.3	117
871	Daily total physical activity level and total cancer risk in men and women: results from a large-scale population-based cohort study in Japan. <i>American Journal of Epidemiology</i> , 2008 , 168, 391-403	3.8	116
870	Helicobacter pylori, dietary factors, and atrophic gastritis in five Japanese populations with different gastric cancer mortality. <i>Cancer Causes and Control</i> , 1993 , 4, 297-305	2.8	114
869	Cigarette smoking and risk of stroke and its subtypes among middle-aged Japanese men and women: the JPHC Study Cohort I. <i>Stroke</i> , 2004 , 35, 1248-53	6.7	111
868	Genome-wide association analysis in East Asians identifies breast cancer susceptibility loci at 1q32.1, 5q14.3 and 15q26.1. <i>Nature Genetics</i> , 2014 , 46, 886-90	36.3	110
867	Population attributable fraction of mortality associated with tobacco smoking in Japan: a pooled analysis of three large-scale cohort studies. <i>Journal of Epidemiology</i> , 2008 , 18, 251-64	3.4	109
866	Reproductive factors, hormone use and the risk of lung cancer among middle-aged never-smoking Japanese women: a large-scale population-based cohort study. <i>International Journal of Cancer</i> , 2005 , 117, 662-6	7.5	109
865	Genome-wide association study of colorectal cancer identifies six new susceptibility loci. <i>Nature Communications</i> , 2015 , 6, 7138	17.4	106

864	The impact of green tea and coffee consumption on the reduced risk of stroke incidence in Japanese population: the Japan public health center-based study cohort. <i>Stroke</i> , 2013 , 44, 1369-74	6.7	106
863	Effect of alcohol consumption, cigarette smoking and flushing response on esophageal cancer risk: a population-based cohort study (JPHC study). <i>Cancer Letters</i> , 2009 , 275, 240-6	9.9	106
862	Association of Helicobacter pylori infection and environmental factors in non-cardia gastric cancer in Japan. <i>Gastric Cancer</i> , 2004 , 7, 46-53	7.6	106
861	Impact of metabolic factors on subsequent cancer risk: results from a large-scale population-based cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2009 , 18, 240-7	2	104
860	Validity and reproducibility of a self-administered food frequency questionnaire used in the baseline survey of the JPHC Study Cohort I. <i>Journal of Epidemiology</i> , 2003 , 13, S125-33	3.4	104
859	Salty food intake and risk of Helicobacter pylori infection. <i>Japanese Journal of Cancer Research</i> , 1994 , 85, 474-8		104
858	Effect of Helicobacter pylori infection combined with CagA and pepsinogen status on gastric cancer development among Japanese men and women: a nested case-control study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2006 , 15, 1341-7	4	101
857	Validity of Short and Long Self-Administered Food Frequency Questionnaires in Ranking Dietary Intake in Middle-Aged and Elderly Japanese in the Japan Public Health Center-Based Prospective Study for the Next Generation (JPHC-NEXT) Protocol Area. <i>Journal of Epidemiology</i> , 2016 , 26, 420-32	3.4	99
856	Lack of long-term effect of vitamin C supplementation on blood pressure. <i>Hypertension</i> , 2002 , 40, 797-803	8.5	99
855	Calcium, vitamin D and dairy intake in relation to type 2 diabetes risk in a Japanese cohort. <i>Diabetologia</i> , 2009 , 52, 2542-50	10.3	98
854	Interaction between adiponectin and leptin influences the risk of colorectal adenoma. <i>Cancer Research</i> , 2010 , 70, 5430-7	10.1	96
853	Quality of diet and mortality among Japanese men and women: Japan Public Health Center based prospective study. <i>BMJ, The</i> , 2016 , 352, i1209	5.9	95
852	Serum cholesterol levels in relation to the incidence of cancer: the JPHC study cohorts. <i>International Journal of Cancer</i> , 2009 , 125, 2679-86	7.5	94
851	Identification of a functional genetic variant at 16q12.1 for breast cancer risk: results from the Asia Breast Cancer Consortium. <i>PLoS Genetics</i> , 2010 , 6, e1001002	6	93
850	A novel allelic variant of serum amyloid A, SAA1 gamma: genomic evidence, evolution, frequency, and implication as a risk factor for reactive systemic AA-amyloidosis. <i>Human Molecular Genetics</i> , 1995 , 4, 1083-7	5.6	92
849	A data-based approach for designing a semiquantitative food frequency questionnaire for a population-based prospective study in Japan. <i>Journal of Epidemiology</i> , 1996 , 6, 45-53	3.4	92
848	Soy product and isoflavone intakes are associated with a lower risk of type 2 diabetes in overweight Japanese women. <i>Journal of Nutrition</i> , 2010 , 140, 580-6	4.1	91
847	Blood pressure and the risk of stroke, cardiovascular disease, and all-cause mortality among Japanese: the JPHC Study. <i>American Journal of Hypertension</i> , 2009 , 22, 273-80	2.3	90

846	Psychological factors, coffee and risk of diabetes mellitus among middle-aged Japanese: a population-based prospective study in the JPHC study cohort. <i>Endocrine Journal</i> , 2009 , 56, 459-68	2.9	90
845	Alcohol drinking and colorectal cancer in Japanese: a pooled analysis of results from five cohort studies. <i>American Journal of Epidemiology</i> , 2008 , 167, 1397-406	3.8	90
844	Identification of six new genetic loci associated with atrial fibrillation in the Japanese population. <i>Nature Genetics</i> , 2017 , 49, 953-958	36.3	89
843	Dietary cadmium intake and breast cancer risk in Japanese women: a case-control study. <i>International Journal of Hygiene and Environmental Health</i> , 2014 , 217, 70-7	6.9	89
842	Effect of coffee and green tea consumption on the risk of liver cancer: cohort analysis by hepatitis virus infection status. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2009 , 18, 1746-53	4	88
841	Consumption of sodium and salted foods in relation to cancer and cardiovascular disease: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2010 , 91, 456-64	7	87
840	Dietary calcium intake and risks of stroke, its subtypes, and coronary heart disease in Japanese: the JPHC Study Cohort I. <i>Stroke</i> , 2008 , 39, 2449-56	6.7	87
839	Body mass index, physical activity and the risk of pancreatic cancer in relation to smoking status and history of diabetes: a large-scale population-based cohort study in Japan--the JPHC study. <i>Cancer Causes and Control</i> , 2007 , 18, 603-12	2.8	87
838	Single measurement of serum phospholipid fatty acid as a biomarker of specific fatty acid intake in middle-aged Japanese men. <i>European Journal of Clinical Nutrition</i> , 2001 , 55, 643-50	5.2	87
837	Association between type 2 diabetes and risk of cancer mortality: a pooled analysis of over 771,000 individuals in the Asia Cohort Consortium. <i>Diabetologia</i> , 2017 , 60, 1022-1032	10.3	86
836	Plasma isoflavones and subsequent risk of prostate cancer in a nested case-control study: the Japan Public Health Center. <i>Journal of Clinical Oncology</i> , 2008 , 26, 5923-9	2.2	86
835	Egg consumption, serum total cholesterol concentrations and coronary heart disease incidence: Japan Public Health Center-based prospective study. <i>British Journal of Nutrition</i> , 2006 , 96, 921-8	3.6	86
834	Study design and organization of the JPHC study. Japan Public Health Center-based Prospective Study on Cancer and Cardiovascular Diseases. <i>Journal of Epidemiology</i> , 2001 , 11, S3-7	3.4	86
833	Large-scale genome-wide association study in a Japanese population identifies novel susceptibility loci across different diseases. <i>Nature Genetics</i> , 2020 , 52, 669-679	36.3	85
832	Diabetes mellitus and cancer risk: pooled analysis of eight cohort studies in Japan. <i>Cancer Science</i> , 2013 , 104, 1499-507	6.9	85
831	Tobacco smoking and gastric cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2006 , 36, 800-7	2.8	85
830	Plasma vitamin D and risk of colorectal cancer: the Japan Public Health Center-Based Prospective Study. <i>British Journal of Cancer</i> , 2007 , 97, 446-51	8.7	85
829	Prospective study of three major dietary patterns and risk of gastric cancer in Japan. <i>International Journal of Cancer</i> , 2004 , 110, 435-42	7.5	85

828	Carotenoid intakes and risk of breast cancer defined by estrogen receptor and progesterone receptor status: a pooled analysis of 18 prospective cohort studies. <i>American Journal of Clinical Nutrition</i> , 2012 , 95, 713-25	7	84
827	Meat intake and cause-specific mortality: a pooled analysis of Asian prospective cohort studies. <i>American Journal of Clinical Nutrition</i> , 2013 , 98, 1032-41	7	83
826	Identification of 28 new susceptibility loci for type 2 diabetes in the Japanese population. <i>Nature Genetics</i> , 2019 , 51, 379-386	36.3	83
825	Cruciferous vegetables, mushrooms, and gastrointestinal cancer risks in a multicenter, hospital-based case-control study in Japan. <i>Nutrition and Cancer</i> , 2003 , 46, 138-47	2.8	82
824	Dietary patterns and subsequent colorectal cancer risk by subsite: a prospective cohort study. <i>International Journal of Cancer</i> , 2005 , 115, 790-8	7.5	82
823	The impact of the metabolic syndrome and its components on the incidence of ischemic heart disease and stroke: the Japan public health center-based study. <i>Hypertension Research</i> , 2009 , 32, 289-98 ^{4.7}	4.7	81
822	Folate, vitamin B6, vitamin B12, and vitamin B2 intake, genetic polymorphisms of related enzymes, and risk of colorectal cancer in a hospital-based case-control study in Japan. <i>Nutrition and Cancer</i> , 2005 , 53, 42-50	2.8	81
821	Fish intake and type 2 diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2011 , 94, 884-91	7	79
820	Genome-wide association study identifies breast cancer risk variant at 10q21.2: results from the Asia Breast Cancer Consortium. <i>Human Molecular Genetics</i> , 2011 , 20, 4991-9	5.6	79
819	Plasma C-reactive protein and risk of colorectal cancer in a nested case-control study: Japan Public Health Center-based prospective study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2006 , 15, 690-5 ⁴	5.4	79
818	Prognostic significance of the overexpression of c-erbB-2 protein in adenocarcinoma of the uterine cervix. <i>Cancer</i> , 1994 , 73, 148-53	6.4	79
817	Physical activity and risk of colorectal cancer in Japanese men and women: the Japan Public Health Center-based prospective study. <i>Cancer Causes and Control</i> , 2007 , 18, 199-209	2.8	78
816	Validity of a self-administered food frequency questionnaire used in the 5-year follow-up survey of the JPHC Study Cohort I: comparison with dietary records for food groups. <i>Journal of Epidemiology</i> , 2003 , 13, S57-63	3.4	78
815	Self-administered food frequency questionnaire used in the 5-year follow-up survey of the JPHC Study: questionnaire structure, computation algorithms, and area-based mean intake. <i>Journal of Epidemiology</i> , 2003 , 13, S13-22	3.4	77
814	Burden of total and cause-specific mortality related to tobacco smoking among adults aged ≥45 years in Asia: a pooled analysis of 21 cohorts. <i>PLoS Medicine</i> , 2014 , 11, e1001631	11.6	76
813	Body mass index and mortality from all causes and major causes in Japanese: results of a pooled analysis of 7 large-scale cohort studies. <i>Journal of Epidemiology</i> , 2011 , 21, 417-30	3.4	76
812	Impact of population aging on trends in diabetes prevalence: A meta-regression analysis of 160,000 Japanese adults. <i>Journal of Diabetes Investigation</i> , 2015 , 6, 533-42	3.9	75
811	Dietary intake of folate, vitamin B6, and vitamin B12, genetic polymorphism of related enzymes, and risk of breast cancer: a case-control study in Brazilian women. <i>BMC Cancer</i> , 2009 , 9, 122	4.8	75

810	Clinicopathologic characteristics of adenosquamous carcinoma of the lung. <i>Cancer</i> , 1991 , 67, 649-54	6.4	75
809	Soft drink, 100% fruit juice, and vegetable juice intakes and risk of diabetes mellitus. <i>Clinical Nutrition</i> , 2013 , 32, 300-8	5.9	74
808	Reproducibility and validity of dietary patterns assessed by a food frequency questionnaire used in the 5-year follow-up survey of the Japan Public Health Center-Based Prospective Study. <i>Journal of Epidemiology</i> , 2012 , 22, 205-15	3.4	74
807	Overweight and obesity trends among Japanese adults: a 10-year follow-up of the JPHC Study. <i>International Journal of Obesity</i> , 2008 , 32, 1861-7	5.5	74
806	Immunohistochemical study on overexpression of c-erbB-2 protein in human breast cancer: its correlation with gene amplification and long-term survival of patients. <i>Japanese Journal of Cancer Research</i> , 1990 , 81, 327-32		74
805	Allele frequencies of single nucleotide polymorphisms (SNPs) in 40 candidate genes for gene-environment studies on cancer: data from population-based Japanese random samples. <i>Journal of Human Genetics</i> , 2003 , 48, 654-658	4.3	73
804	Genome-wide association study identifies seven novel susceptibility loci for primary open-angle glaucoma. <i>Human Molecular Genetics</i> , 2018 , 27, 1486-1496	5.6	72
803	Impact of the revision of a nutrient database on the validity of a self-administered food frequency questionnaire (FFQ). <i>Journal of Epidemiology</i> , 2006 , 16, 107-16	3.4	72
802	Demographics, lifestyles, health characteristics, and dietary intake among dietary supplement users in Japan. <i>International Journal of Epidemiology</i> , 2003 , 32, 546-53	7.8	72
801	Tobacco smoking and lung cancer risk: an evaluation based on a systematic review of epidemiological evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2006 , 36, 309-24	2.8	71
800	Green tea consumption and subsequent risk of gastric cancer by subsite: the JPHC Study. <i>Cancer Causes and Control</i> , 2004 , 15, 483-91	2.8	71
799	Dietary patterns and all-cause, cancer, and cardiovascular disease mortality in Japanese men and women: The Japan public health center-based prospective study. <i>PLoS ONE</i> , 2017 , 12, e0174848	3.7	71
798	Urinary phthalate monoesters and endometriosis in infertile Japanese women. <i>Science of the Total Environment</i> , 2009 , 408, 37-42	10.2	70
797	Coffee consumption and risk of colorectal cancer in a population-based prospective cohort of Japanese men and women. <i>International Journal of Cancer</i> , 2007 , 121, 1312-8	7.5	70
796	Fruit and vegetable consumption and squamous cell carcinoma of the esophagus in Japan: the JPHC study. <i>International Journal of Cancer</i> , 2008 , 123, 1935-40	7.5	70
795	Intake of n-3 and n-6 polyunsaturated fatty acids and development of colorectal cancer by subsite: Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2011 , 129, 1718-29	7.5	69
794	Isoflavone intake and risk of lung cancer: a prospective cohort study in Japan. <i>American Journal of Clinical Nutrition</i> , 2010 , 91, 722-8	7	68
793	Calvasculin, as a factor affecting the microfilament assemblies in rat fibroblasts transfected by src gene. <i>FEBS Letters</i> , 1993 , 324, 51-5	3.8	68

792	Novel Common Genetic Susceptibility Loci for Colorectal Cancer. <i>Journal of the National Cancer Institute</i> , 2019 , 111, 146-157	9.7	67
791	Insulin resistance and cancer: epidemiological evidence. <i>Endocrine-Related Cancer</i> , 2012 , 19, F1-8	5.7	67
790	Soy intake and breast cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2014 , 44, 282-95	2.8	65
789	Heterocyclic amines content of meat and fish cooked by Brazilian methods. <i>Journal of Food Composition and Analysis</i> , 2010 , 23, 61-69	4.1	65
788	Plasma C-peptide, insulin-like growth factor-I, insulin-like growth factor binding proteins and risk of colorectal cancer in a nested case-control study: the Japan public health center-based prospective study. <i>International Journal of Cancer</i> , 2007 , 120, 2007-12	7.5	65
787	Cancer incidence rates among Japanese immigrants in the city of Sã Paulo, Brazil, 1969-78. <i>Cancer Causes and Control</i> , 1990 , 1, 189-93	2.8	65
786	Daily omega-3 fatty acid intake and depression in Japanese patients with newly diagnosed lung cancer. <i>British Journal of Cancer</i> , 2004 , 90, 787-93	8.7	64
785	Metabolic syndrome and all-cause and cardiovascular disease mortality: Japan Public Health Center-based Prospective (JPHC) Study. <i>Circulation Journal</i> , 2009 , 73, 878-84	2.9	63
784	Passive smoking and lung cancer in Japanese non-smoking women: a prospective study. <i>International Journal of Cancer</i> , 2008 , 122, 653-7	7.5	63
783	Green tea and coffee intake and risk of pancreatic cancer in a large-scale, population-based cohort study in Japan (JPHC study). <i>European Journal of Cancer Prevention</i> , 2007 , 16, 542-8	2	63
782	Trans-ethnic genome-wide association study of colorectal cancer identifies a new susceptibility locus in VTI1A. <i>Nature Communications</i> , 2014 , 5, 4613	17.4	62
781	The effects of smoking and smoking cessation on mortality from cardiovascular disease among Japanese: pooled analysis of three large-scale cohort studies in Japan. <i>Tobacco Control</i> , 2010 , 19, 50-7	5.3	62
780	Diabetes mellitus and risk of stroke and its subtypes among Japanese: the Japan public health center study. <i>Stroke</i> , 2011 , 42, 2611-4	6.7	62
779	Green tea consumption and gastric cancer in Japanese: a pooled analysis of six cohort studies. <i>Gut</i> , 2009 , 58, 1323-32	19.2	62
778	Dietary calcium, vitamin D, and the risk of colorectal cancer. <i>American Journal of Clinical Nutrition</i> , 2008 , 88, 1576-83	7	62
777	Validity of the self-administered food frequency questionnaire used in the 5-year follow-up survey of the JPHC Study Cohort I: comparison with dietary records for main nutrients. <i>Journal of Epidemiology</i> , 2003 , 13, S51-6	3.4	62
776	Dietary intake of folate, vitamin B2, vitamin B6, vitamin B12, genetic polymorphism of related enzymes, and risk of breast cancer: a case-control study in Japan. <i>Nutrition and Cancer</i> , 2009 , 61, 447-56	2.8	61
775	Association between body mass index and the colorectal cancer risk in Japan: pooled analysis of population-based cohort studies in Japan. <i>Annals of Oncology</i> , 2012 , 23, 479-90	10.3	61

774	Fresh and pickled vegetable consumption and gastric cancer in Japanese and Korean populations: a meta-analysis of observational studies. <i>Cancer Science</i> , 2010 , 101, 508-16	6.9	60
773	Replication and functional genomic analyses of the breast cancer susceptibility locus at 6q25.1 generalize its importance in women of Chinese, Japanese, and European ancestry. <i>Cancer Research</i> , 2011 , 71, 1344-55	10.1	60
772	Colorectal cancer screening using fecal occult blood test and subsequent risk of colorectal cancer: a prospective cohort study in Japan. <i>Cancer Detection and Prevention</i> , 2007 , 31, 3-11		60
771	Association of Animal and Plant Protein Intake With All-Cause and Cause-Specific Mortality in a Japanese Cohort. <i>JAMA Internal Medicine</i> , 2019 , 179, 1509-1518	11.5	59
770	Serum organochlorines and breast cancer risk in Japanese women: a case-control study. <i>Cancer Causes and Control</i> , 2009 , 20, 567-80	2.8	59
769	Social support and stroke and coronary heart disease: the JPHC study cohorts II. <i>Stroke</i> , 2008 , 39, 768-756.7		59
768	Reproducibility of a self-administered food frequency questionnaire used in the 5-year follow-up survey of the JPHC Study Cohort I to assess food and nutrient intake. <i>Journal of Epidemiology</i> , 2003 , 13, S115-24	3.4	59
767	hOGG1 Ser326Cys polymorphism, interaction with environmental exposures, and gastric cancer risk in Japanese populations. <i>Cancer Science</i> , 2004 , 95, 977-83	6.9	59
766	Clinical and radiologic assessments of the results of hepatectomy for small hepatocellular carcinoma and therapeutic arterial embolization for postoperative recurrence. <i>Cancer</i> , 1989 , 64, 1848-52.6.4		59
765	Body weight at age 20 years, subsequent weight change and breast cancer risk defined by estrogen and progesterone receptor status--the Japan public health center-based prospective study. <i>International Journal of Cancer</i> , 2011 , 129, 1214-24	7.5	58
764	Food and nutrient intakes assessed with dietary records for the validation study of a self-administered food frequency questionnaire in JPHC Study Cohort I. <i>Journal of Epidemiology</i> , 2003 , 13, S23-50	3.4	58
763	Active and passive smoking and breast cancer risk in middle-aged Japanese women. <i>International Journal of Cancer</i> , 2005 , 114, 317-22	7.5	58
762	Soft drink intake in relation to incident ischemic heart disease, stroke, and stroke subtypes in Japanese men and women: the Japan Public Health Centre-based study cohort I. <i>American Journal of Clinical Nutrition</i> , 2012 , 96, 1390-7	7	57
761	Perceived level of life enjoyment and risks of cardiovascular disease incidence and mortality: the Japan public health center-based study. <i>Circulation</i> , 2009 , 120, 956-63	16.7	57
760	Alcohol drinking and esophageal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 677-92	2.8	57
759	Cigarette smoking and completed suicide among middle-aged men: a population-based cohort study in Japan. <i>Annals of Epidemiology</i> , 2005 , 15, 286-92	6.4	57
758	Fish, long-chain n-3 polyunsaturated fatty acids, and risk of colorectal cancer in middle-aged Japanese: the JPHC study. <i>Nutrition and Cancer</i> , 2004 , 49, 32-40	2.8	57
757	Preoperative radioactive therapy for esophageal carcinoma. Randomized evaluation trial in eight institutions. <i>Chest</i> , 1988 , 93, 1054-8	5.3	57

756	Report of the Japan Diabetes Society/Japanese Cancer Association Joint Committee on Diabetes and Cancer. <i>Cancer Science</i> , 2013 , 104, 965-76	6.9	56
755	Body mass index and weight change since 20 years of age and risk of coronary heart disease among Japanese: the Japan Public Health Center-Based Study. <i>International Journal of Obesity</i> , 2008 , 32, 144-51	5.5	56
754	Tobacco smoking and breast cancer risk: an evaluation based on a systematic review of epidemiological evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2006 , 36, 387-94	2.8	56
753	Blood pressure change in a free-living population-based dietary modification study in Japan. <i>Journal of Hypertension</i> , 2006 , 24, 451-8	1.9	56
752	Impact of alcohol drinking on total cancer risk: data from a large-scale population-based cohort study in Japan. <i>British Journal of Cancer</i> , 2005 , 92, 182-7	8.7	56
751	Prediction of the 10-year probability of gastric cancer occurrence in the Japanese population: the JPHC study cohort II. <i>International Journal of Cancer</i> , 2016 , 138, 320-31	7.5	56
750	Seaweed consumption and the risk of thyroid cancer in women: the Japan Public Health Center-based Prospective Study. <i>European Journal of Cancer Prevention</i> , 2012 , 21, 254-60	2	55
749	Association of green tea consumption with mortality due to all causes and major causes of death in a Japanese population: the Japan Public Health Center-based Prospective Study (JPHC Study). <i>Annals of Epidemiology</i> , 2015 , 25, 512-518.e3	6.4	54
748	Coffee consumption and risk of endometrial cancer: a prospective study in Japan. <i>International Journal of Cancer</i> , 2008 , 123, 2406-10	7.5	54
747	Associations between serum levels of selected organochlorine compounds and endometriosis in infertile Japanese women. <i>Environmental Research</i> , 2005 , 99, 118-25	7.9	54
746	Cross-sectional study with multiple measurements of biological markers for assessing stomach cancer risks at the population level. <i>Environmental Health Perspectives</i> , 1992 , 98, 207-10	8.4	54
745	Dietary soy and isoflavone intake and risk of colorectal cancer in the Japan public health center-based prospective study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2008 , 17, 2128-35	4	53
744	Role and impact of menstrual and reproductive factors on breast cancer risk in Japan. <i>European Journal of Cancer Prevention</i> , 2007 , 16, 116-23	2	53
743	Association between lung cancer incidence and family history of lung cancer: data from a large-scale population-based cohort study, the JPHC study. <i>Chest</i> , 2006 , 130, 968-75	5.3	53
742	Impact of body mass index on the risk of total cancer incidence and mortality among middle-aged Japanese: data from a large-scale population-based cohort study--the JPHC study. <i>Cancer Causes and Control</i> , 2004 , 15, 671-80	2.8	53
741	Meat consumption and colorectal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2014 , 44, 641-50	2.8	52
740	Social support and cancer incidence and mortality: the JPHC study cohort II. <i>Cancer Causes and Control</i> , 2013 , 24, 847-60	2.8	52
739	Fish intake and serum levels of organochlorines among Japanese women. <i>Science of the Total Environment</i> , 2006 , 359, 90-100	10.2	52

738	Characterizing rare and low-frequency height-associated variants in the Japanese population. <i>Nature Communications</i> , 2019 , 10, 4393	17.4	51
737	Effect of soy isoflavones on endometriosis: interaction with estrogen receptor 2 gene polymorphism. <i>Epidemiology</i> , 2007 , 18, 402-8	3.1	51
736	hOGG1 exon7 polymorphism and gastric cancer in case-control studies of Japanese Brazilians and non-Japanese Brazilians. <i>Cancer Letters</i> , 2001 , 170, 53-61	9.9	51
735	Population-specific and trans-ancestry genome-wide analyses identify distinct and shared genetic risk loci for coronary artery disease. <i>Nature Genetics</i> , 2020 , 52, 1169-1177	36.3	51
734	Association of coffee intake with total and cause-specific mortality in a Japanese population: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2015 , 101, 1029-37	7	50
733	Genetic and epigenetic alterations in normal tissues have differential impacts on cancer risk among tissues. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, 1328-1333	11.5	50
732	Low-carbohydrate diet and type 2 diabetes risk in Japanese men and women: the Japan Public Health Center-Based Prospective Study. <i>PLoS ONE</i> , 2015 , 10, e0118377	3.7	50
731	Atrophic gastritis, <i>Helicobacter pylori</i> , and colorectal cancer risk: a case-control study. <i>Helicobacter</i> , 2007 , 12, 328-32	4.9	50
730	Patterns of alcohol drinking and all-cause mortality: results from a large-scale population-based cohort study in Japan. <i>American Journal of Epidemiology</i> , 2007 , 165, 1039-46	3.8	50
729	Impact of demographic, environmental, and lifestyle factors on vitamin D sufficiency in 9084 Japanese adults. <i>Bone</i> , 2015 , 74, 10-7	4.7	49
728	Plasma biomarker discovery and validation for colorectal cancer by quantitative shotgun mass spectrometry and protein microarray. <i>Cancer Science</i> , 2011 , 102, 630-8	6.9	49
727	Plasma levels of carotenoids, retinol and tocopherol and the risk of gastric cancer in Japan: a nested case-control study. <i>Carcinogenesis</i> , 2008 , 29, 1042-8	4.6	49
726	Validity of a self-administered food frequency questionnaire used in the 5-year follow-up survey of the JPHC Study Cohort I to assess fatty acid intake: comparison with dietary records and serum phospholipid level. <i>Journal of Epidemiology</i> , 2003 , 13, S64-81	3.4	49
725	CYP1A1 and CYP2E1 polymorphism and lung cancer, case-control study in Rio de Janeiro, Brazil. <i>Pharmacogenetics and Genomics</i> , 1995 , 5 Spec No, S145-8		49
724	Urinary salt excretion and stomach cancer mortality among four Japanese populations. <i>Cancer Causes and Control</i> , 1991 , 2, 165-8	2.8	49
723	Weight change and all-cause, cancer and cardiovascular disease mortality in Japanese men and women: the Japan Public Health Center-Based Prospective Study. <i>International Journal of Obesity</i> , 2010 , 34, 348-56	5.5	48
722	Associations between unprocessed red and processed meat, poultry, seafood and egg intake and the risk of prostate cancer: A pooled analysis of 15 prospective cohort studies. <i>International Journal of Cancer</i> , 2016 , 138, 2368-82	7.5	48
721	Dietary intake of saturated fatty acids and incident stroke and coronary heart disease in Japanese communities: the JPHC Study. <i>European Heart Journal</i> , 2013 , 34, 1225-32	9.5	47

720	Visceral fat volume and the prevalence of colorectal adenoma. <i>American Journal of Epidemiology</i> , 2009 , 170, 1502-11	3.8	47
719	Long-term dietary cadmium intake and cancer incidence. <i>Epidemiology</i> , 2012 , 23, 368-76	3.1	47
718	Impact of alcohol intake on total mortality and mortality from major causes in Japan: a pooled analysis of six large-scale cohort studies. <i>Journal of Epidemiology and Community Health</i> , 2012 , 66, 448-56 ^{5.1}	5.1	47
717	Intake of folate, vitamin B6 and vitamin B12 and the risk of CHD: the Japan Public Health Center-Based Prospective Study Cohort I. <i>Journal of the American College of Nutrition</i> , 2008 , 27, 127-36	3.5	47
716	Risk factors of biliary tract cancer in a large-scale population-based cohort study in Japan (JPHC study); with special focus on cholelithiasis, body mass index, and their effect modification. <i>Cancer Causes and Control</i> , 2008 , 19, 33-41	2.8	47
715	Coffee consumption and risk of type 2 diabetes mellitus. <i>Lancet, The</i> , 2003 , 361, 703-4	4.0	47
714	Development of substituted fatty acid food composition table for the use in nutritional epidemiologic studies for Japanese populations: its methodological backgrounds and the evaluation. <i>Journal of Epidemiology</i> , 1999 , 9, 190-207	3.4	47
713	Genetic polymorphisms of ADH1B, ADH1C and ALDH2, alcohol consumption, and the risk of gastric cancer: the Japan Public Health Center-based prospective study. <i>Carcinogenesis</i> , 2015 , 36, 223-31	4.6	46
712	Association of Breakfast Intake With Incident Stroke and Coronary Heart Disease: The Japan Public Health Center-Based Study. <i>Stroke</i> , 2016 , 47, 477-81	6.7	46
711	Body mass index and breast cancer risk in Japan: a pooled analysis of eight population-based cohort studies. <i>Annals of Oncology</i> , 2014 , 25, 519-24	10.3	46
710	Green tea drinking and subsequent risk of breast cancer in a population-based cohort of Japanese women. <i>Breast Cancer Research</i> , 2010 , 12, R88	8.3	46
709	Metabolic factors and subsequent risk of hepatocellular carcinoma by hepatitis virus infection status: a large-scale population-based cohort study of Japanese men and women (JPHC Study Cohort II). <i>Cancer Causes and Control</i> , 2009 , 20, 741-50	2.8	46
708	Coffee, green tea, and caffeine consumption and subsequent risk of bladder cancer in relation to smoking status: a prospective study in Japan. <i>Cancer Science</i> , 2009 , 100, 294-91	6.9	46
707	Prospective study showing that dietary vitamin C reduced the risk of age-related cataracts in a middle-aged Japanese population. <i>European Journal of Nutrition</i> , 2007 , 46, 118-24	5.2	46
706	Cigarette smoking and risk of coronary heart disease incidence among middle-aged Japanese men and women: the JPHC Study Cohort I. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2006 , 13, 207-13		46
705	Body size and risk for breast cancer in relation to estrogen and progesterone receptor status in Japan. <i>Annals of Epidemiology</i> , 2007 , 17, 304-12	6.4	46
704	Estimation of dietary HCA intakes in a large-scale population-based prospective study in Japan. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2002 , 506-507, 233-41	3.3	46
703	Association between endometriosis and genetic polymorphisms of the estradiol-synthesizing enzyme genes HSD17B1 and CYP19. <i>Human Reproduction</i> , 2005 , 20, 974-8	5.7	46

702	Genetic Predisposition to Ischemic Stroke: A Polygenic Risk Score. <i>Stroke</i> , 2017 , 48, 253-258	6.7	45
701	Plasma isoflavones and the risk of lung cancer in women: a nested case-control study in Japan. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2011 , 20, 419-27	4	45
700	Tobacco smoking and colorectal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2006 , 36, 25-39	2.8	45
699	Background characteristics of basic health examination participants: the JPHC Study Baseline Survey. <i>Journal of Epidemiology</i> , 2003 , 13, 216-25	3.4	45
698	Role of potassium channels in regulation of brain arteriolar tone: comparison of cerebrum versus brain stem. <i>Stroke</i> , 2001 , 32, 218-24	6.7	45
697	Oxidative DNA damage induced by silica in vivo. <i>Environmental Research</i> , 1995 , 69, 102-7	7.9	45
696	Smoking cessation increases short-term risk of type 2 diabetes irrespective of weight gain: the Japan Public Health Center-Based Prospective Study. <i>PLoS ONE</i> , 2012 , 7, e17061	3.7	45
695	Plasma tea polyphenol levels and subsequent risk of breast cancer among Japanese women: a nested case-control study. <i>Breast Cancer Research and Treatment</i> , 2010 , 124, 827-34	4.4	44
694	A novel splice-site variant of the base excision repair gene MYH is associated with production of an aberrant mRNA transcript encoding a truncated MYH protein not localized in the nucleus. <i>Carcinogenesis</i> , 2004 , 25, 1859-66	4.6	44
693	Fermented and non-fermented soy food consumption and gastric cancer in Japanese and Korean populations: a meta-analysis of observational studies. <i>Cancer Science</i> , 2011 , 102, 231-44	6.9	43
692	Risk factors for breast cancer: epidemiological evidence from Japanese studies. <i>Cancer Science</i> , 2011 , 102, 1607-14	6.9	43
691	Vegetable, fruit and antioxidant nutrient consumption and subsequent risk of hepatocellular carcinoma: a prospective cohort study in Japan. <i>British Journal of Cancer</i> , 2009 , 100, 181-4	8.7	43
690	Green tea consumption and gastric cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 335-46	2.8	43
689	Serum triglycerides and colorectal adenoma in a case-control study among cancer screening examinees (Japan). <i>Cancer Causes and Control</i> , 2006 , 17, 1245-52	2.8	43
688	Validity and reproducibility of the self-administered food frequency questionnaire in the JPHC Study Cohort I: study design, conduct and participant profiles. <i>Journal of Epidemiology</i> , 2003 , 13, S2-12	3.4	43
687	Alcohol consumption, smoking, and subsequent risk of colorectal cancer in middle-aged and elderly Japanese men and women: Japan Public Health Center-based prospective study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2003 , 12, 1492-500	4	43
686	Red meat intake may increase the risk of colon cancer in Japanese, a population with relatively low red meat consumption. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2011 , 20, 603-12	1	43
685	Tobacco Smoking and Mortality in Asia: A Pooled Meta-analysis. <i>JAMA Network Open</i> , 2019 , 2, e191474	10.4	42

684	Global methylation levels in peripheral blood leukocyte DNA by LUMA and breast cancer: a case-control study in Japanese women. <i>British Journal of Cancer</i> , 2014 , 110, 2765-71	8.7	42
683	Cigarette smoking and liver cancer risk: an evaluation based on a systematic review of epidemiologic evidence among Japanese. <i>Japanese Journal of Clinical Oncology</i> , 2006 , 36, 445-56	2.8	42
682	Low intake of vitamin B-6 is associated with increased risk of colorectal cancer in Japanese men. <i>Journal of Nutrition</i> , 2007 , 137, 1808-14	4.1	42
681	Plasma 25-hydroxyvitamin D concentration and subsequent risk of total and site specific cancers in Japanese population: large case-cohort study within Japan Public Health Center-based Prospective Study cohort. <i>BMJ, The</i> , 2018 , 360, k671	5.9	41
680	Urinary bisphenol-A concentration in infertile Japanese women and its association with endometriosis: A cross-sectional study. <i>Environmental Health and Preventive Medicine</i> , 2007 , 12, 258-64	4.2	41
679	Evaluation based on systematic review of epidemiological evidence among Japanese populations: tobacco smoking and total cancer risk. <i>Japanese Journal of Clinical Oncology</i> , 2005 , 35, 404-11	2.8	41
678	Associations of all-cause mortality with census-based neighbourhood deprivation and population density in Japan: a multilevel survival analysis. <i>PLoS ONE</i> , 2014 , 9, e97802	3.7	41
677	Investigations in the possibility of early detection of colorectal cancer by gas chromatography/triple-quadrupole mass spectrometry. <i>Oncotarget</i> , 2017 , 8, 17115-17126	3.3	41
676	Dietary fish, n-3 polyunsaturated fatty acid consumption, and depression risk in Japan: a population-based prospective cohort study. <i>Translational Psychiatry</i> , 2017 , 7, e1242	8.6	40
675	Fish, n - 3 polyunsaturated fatty acids and n - 6 polyunsaturated fatty acids intake and breast cancer risk: The Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2015 , 137, 2915-26	7.5	40
674	Association of vegetable and fruit intake with gastric cancer risk among Japanese: a pooled analysis of four cohort studies. <i>Annals of Oncology</i> , 2014 , 25, 1228-33	10.3	40
673	High serum total cholesterol levels is a risk factor of ischemic stroke for general Japanese population: the JPHC study. <i>Atherosclerosis</i> , 2012 , 221, 565-9	3.1	40
672	The effect of 5-year vitamin C supplementation on serum pepsinogen level and Helicobacter pylori infection. <i>Cancer Science</i> , 2003 , 94, 378-82	6.9	40
671	Comparison of cause-specific mortality between respondents and nonrespondents in a population-based prospective study: ten-year follow-up of JPHC Study Cohort I. Japan Public Health Center. <i>Journal of Clinical Epidemiology</i> , 2002 , 55, 150-6	5.7	40
670	Analysis of purine- and pyrimidine-induced vascular responses in the isolated rat cerebral arteriole. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2001 , 280, H767-76	5.2	40
669	Association between GWAS-identified lung adenocarcinoma susceptibility loci and EGFR mutations in never-smoking Asian women, and comparison with findings from Western populations. <i>Human Molecular Genetics</i> , 2017 , 26, 454-465	5.6	40
668	Daily Total Physical Activity and Incident Stroke: The Japan Public Health Center-Based Prospective Study. <i>Stroke</i> , 2017 , 48, 1730-1736	6.7	39
667	Association of Diabetes With All-Cause and Cause-Specific Mortality in Asia: A Pooled Analysis of More Than 1 Million Participants. <i>JAMA Network Open</i> , 2019 , 2, e192696	10.4	39

666	Changing trends in the prevalence of H. pylori infection in Japan (1908-2003): a systematic review and meta-regression analysis of 170,752 individuals. <i>Scientific Reports</i> , 2017 , 7, 15491	4.9	39
665	Association of dietary and genetic factors related to one-carbon metabolism with global methylation level of leukocyte DNA. <i>Cancer Science</i> , 2012 , 103, 2159-64	6.9	39
664	Cigarette smoking and esophageal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 63-73	2.8	39
663	Self-reported stroke and myocardial infarction had adequate sensitivity in a population-based prospective study JPHC (Japan Public Health Center)-based Prospective Study. <i>Journal of Clinical Epidemiology</i> , 2009 , 62, 667-73	5.7	39
662	Public awareness of risk factors for cancer among the Japanese general population: a population-based survey. <i>BMC Public Health</i> , 2006 , 6, 2	4.1	39
661	Dietary fiber intake and subsequent risk of colorectal cancer: the Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2006 , 119, 1475-80	7.5	39
660	Analysis of the AhR, ARNT, and AhRR gene polymorphisms: genetic contribution to endometriosis susceptibility and severity. <i>Fertility and Sterility</i> , 2005 , 84, 454-8	4.8	39
659	Association between serum organochlorines and global methylation level of leukocyte DNA among Japanese women: a cross-sectional study. <i>Science of the Total Environment</i> , 2014 , 490, 603-9	10.2	38
658	Death by suicide and other externally caused injuries following a cancer diagnosis: the Japan Public Health Center-based Prospective Study. <i>Psycho-Oncology</i> , 2014 , 23, 1034-41	3.9	38
657	Alcohol and smoking and subsequent risk of prostate cancer in Japanese men: the Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2014 , 134, 971-8	7.5	38
656	Dietary glycemic index, glycemic load and incidence of type 2 diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study. <i>Nutrition Journal</i> , 2013 , 12, 165	4.3	38
655	Dietary fiber intake and risk of cardiovascular disease in the Japanese population: the Japan Public Health Center-based study cohort. <i>European Journal of Clinical Nutrition</i> , 2011 , 65, 1233-41	5.2	38
654	Association of weight change in different periods of adulthood with risk of type 2 diabetes in Japanese men and women: the Japan Public Health Center-Based Prospective Study. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, 1104-10	5.1	38
653	Obesity and liver cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 212-21	2.8	38
652	Associations of body mass index, smoking, and alcohol consumption with prostate cancer mortality in the Asia Cohort Consortium. <i>American Journal of Epidemiology</i> , 2015 , 182, 381-9	3.8	37
651	Overall and Central Obesity and Risk of Lung Cancer: A Pooled Analysis. <i>Journal of the National Cancer Institute</i> , 2018 , 110, 831-842	9.7	37
650	Body mass index, weight change and risk of stroke and stroke subtypes: the Japan Public Health Center-based prospective (JPHC) study. <i>International Journal of Obesity</i> , 2011 , 35, 283-91	5.5	37
649	Adult height and the risk of cardiovascular disease among middle aged men and women in Japan. <i>European Journal of Epidemiology</i> , 2011 , 26, 13-21	12.1	37

648	Isoflavone consumption and subsequent risk of hepatocellular carcinoma in a population-based prospective cohort of Japanese men and women. <i>International Journal of Cancer</i> , 2009 , 124, 1644-9	7.5	37
647	Clinical biopsychosocial risk factors for depression in lung cancer patients: a comprehensive analysis using data from the Lung Cancer Database Project. <i>Annals of Oncology</i> , 2012 , 23, 1973-1979	10.3	37
646	Alcohol consumption-associated breast cancer incidence and potential effect modifiers: the Japan Public Health Center-based Prospective Study. <i>International Journal of Cancer</i> , 2010 , 127, 685-95	7.5	37
645	A prospective study of the significance of gene and chromosome alterations as prognostic indicators of breast cancer patients with lymph node metastases. <i>Breast Cancer Research and Treatment</i> , 1998 , 48, 21-32	4.4	37
644	Plasma tea polyphenols and gastric cancer risk: a case-control study nested in a large population-based prospective study in Japan. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2008 , 17, 343-51	4	37
643	Alcohol consumption and suicide among middle-aged men in Japan. <i>British Journal of Psychiatry</i> , 2006 , 188, 231-6	5.4	37
642	Effect of vitamin C on common cold: randomized controlled trial. <i>European Journal of Clinical Nutrition</i> , 2006 , 60, 9-17	5.2	37
641	Temporal change in the reproducibility of a self-administered food frequency questionnaire. <i>American Journal of Epidemiology</i> , 1995 , 142, 1231-5	3.8	37
640	Calvasculin, an encoded protein from mRNA termed pEL-98, 18A2, 42A, or p9Ka, is secreted by smooth muscle cells in culture and exhibits Ca(2+)-dependent binding to 36-kDa microfibril-associated glycoprotein. <i>Journal of Biological Chemistry</i> , 1992 , 267, 17136-40	5.4	37
639	Calvasculin, an encoded protein from mRNA termed pEL-98, 18A2, 42A, or p9Ka, is secreted by smooth muscle cells in culture and exhibits Ca(2+)-dependent binding to 36-kDa microfibril-associated glycoprotein.. <i>Journal of Biological Chemistry</i> , 1992 , 267, 17136-17140	5.4	37
638	Helicobacter pylori blood biomarker for gastric cancer risk in East Asia. <i>International Journal of Epidemiology</i> , 2016 , 45, 774-81	7.8	37
637	Low Free Testosterone and Prostate Cancer Risk: A Collaborative Analysis of 20 Prospective Studies. <i>European Urology</i> , 2018 , 74, 585-594	10.2	36
636	Validity of a self-administered food frequency questionnaire for middle-aged urban cancer screenees: comparison with 4-day weighed dietary records. <i>Journal of Epidemiology</i> , 2011 , 21, 447-58	3.4	36
635	Living arrangement and coronary heart disease: the JPHC study. <i>Heart</i> , 2009 , 95, 577-83	5.1	36
634	Trends in smoking by birth cohorts born between 1900 and 1977 in Japan. <i>Preventive Medicine</i> , 2006 , 42, 120-7	4.3	36
633	Cross-sectional Epidemiologic Study for Assessing Cancer Risks at the Population Level. <i>Journal of Epidemiology</i> , 1992 , 2, 75-81	3.4	36
632	Quantification of the increase in thyroid cancer prevalence in Fukushima after the nuclear disaster in 2011--a potential overdiagnosis?. <i>Japanese Journal of Clinical Oncology</i> , 2016 , 46, 284-6	2.8	35
631	Fruit and vegetable consumption, Helicobacter pylori antibodies, and gastric cancer risk: A pooled analysis of prospective studies in China, Japan, and Korea. <i>International Journal of Cancer</i> , 2017 , 140, 591-599	7.5	35

630	Vegetable and fruit intake and risk of type 2 diabetes: Japan Public Health Center-based Prospective Study. <i>British Journal of Nutrition</i> , 2013 , 109, 709-17	3.6	35
629	Risk factors for epithelial ovarian cancer in Japan - results from the Japan Public Health Center-based Prospective Study cohort. <i>International Journal of Oncology</i> , 2012 , 40, 21-30	4.4	35
628	Alcohol drinking and colorectal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2006 , 36, 582-97	2.8	35
627	High Dietary Acid Load Score Is Associated with Increased Risk of Type 2 Diabetes in Japanese Men: The Japan Public Health Center-based Prospective Study. <i>Journal of Nutrition</i> , 2016 , 146, 1076-83	4.1	35
626	Development of a prediction model for 10-year risk of hepatocellular carcinoma in middle-aged Japanese: the Japan Public Health Center-based Prospective Study Cohort II. <i>Preventive Medicine</i> , 2012 , 55, 137-43	4.3	34
625	Alcohol drinking and primary liver cancer: a pooled analysis of four Japanese cohort studies. <i>International Journal of Cancer</i> , 2012 , 130, 2645-53	7.5	34
624	Dietary patterns and suicide in Japanese adults: the Japan Public Health Center-based Prospective Study. <i>British Journal of Psychiatry</i> , 2013 , 203, 422-7	5.4	34
623	Alcohol drinking and gastric cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2008 , 38, 8-25	2.8	34
622	Genomic structure and loss of heterozygosity of EPHB2 in colorectal cancer. <i>Cancer Letters</i> , 2001 , 164, 97-104	9.9	34
621	Fruit and vegetable intake and breast cancer risk defined by estrogen and progesterone receptor status: the Japan Public Health Center-based Prospective Study. <i>Cancer Causes and Control</i> , 2013 , 24, 2117-28	2.8	33
620	Fermented Soy Product Intake Is Inversely Associated with the Development of High Blood Pressure: The Japan Public Health Center-Based Prospective Study. <i>Journal of Nutrition</i> , 2017 , 147, 1749-1756	4.1	33
619	Fish, n-3 PUFA consumption, and pancreatic cancer risk in Japanese: a large, population-based, prospective cohort study. <i>American Journal of Clinical Nutrition</i> , 2015 , 102, 1490-7	7	33
618	Cholesterol and egg intakes and the risk of type 2 diabetes: the Japan Public Health Center-based Prospective Study. <i>British Journal of Nutrition</i> , 2014 , 112, 1636-43	3.6	33
617	Red meat consumption is associated with the risk of type 2 diabetes in men but not in women: a Japan Public Health Center-based Prospective Study. <i>British Journal of Nutrition</i> , 2013 , 110, 1910-8	3.6	33
616	Long chain n-3 fatty acids intake, fish consumption and suicide in a cohort of Japanese men and women--the Japan Public Health Center-based (JPHC) prospective study. <i>Journal of Affective Disorders</i> , 2011 , 129, 282-8	6.6	33
615	Plasma organochlorines and subsequent risk of prostate cancer in Japanese men: a nested case-control study. <i>Environmental Health Perspectives</i> , 2010 , 118, 659-65	8.4	33
614	Isoflavone, polymorphisms in estrogen receptor genes and breast cancer risk in case-control studies in Japanese, Japanese Brazilians and non-Japanese Brazilians. <i>Cancer Science</i> , 2009 , 100, 927-33	6.9	33
613	Body mass, tobacco smoking, alcohol drinking and risk of cancer of the small intestine--a pooled analysis of over 500,000 subjects in the Asia Cohort Consortium. <i>Annals of Oncology</i> , 2012 , 23, 1894-8	10.3	33

612	Type A behaviour and risk of coronary heart disease: the JPHC Study. <i>International Journal of Epidemiology</i> , 2008 , 37, 1395-405	7.8	33
611	Cytochrome P450 1B1 mRNA levels in peripheral blood cells and exposure to polycyclic aromatic hydrocarbons in Chinese coke oven workers. <i>Science of the Total Environment</i> , 2002 , 296, 27-33	10.2	33
610	Prevalence of diabetes estimated by plasma glucose criteria combined with standardized measurement of HbA1c among health checkup participants on Miyako Island, Japan. <i>Diabetes Care</i> , 2000 , 23, 1092-6	14.6	33
609	Increased Levels of Branched-Chain Amino Acid Associated With Increased Risk of Pancreatic Cancer in a Prospective Case-Control Study of a Large Cohort. <i>Gastroenterology</i> , 2018 , 155, 1474-1482.e13-3	13.3	33
608	Diet, nutrition, and cancer risk: what do we know and what is the way forward?. <i>BMJ, The</i> , 2020 , 368, m511	5.9	32
607	GWAS identifies two novel colorectal cancer loci at 16q24.1 and 20q13.12. <i>Carcinogenesis</i> , 2018 , 39, 652-660	6.60	32
606	Meta-analysis of genome-wide association studies identifies multiple lung cancer susceptibility loci in never-smoking Asian women. <i>Human Molecular Genetics</i> , 2016 , 25, 620-9	5.6	32
605	Plasma insulin, C-peptide and blood glucose and the risk of gastric cancer: the Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2015 , 136, 1402-10	7.5	32
604	Development of a point-based prediction model for the incidence of total stroke: Japan public health center study. <i>Stroke</i> , 2013 , 44, 1295-302	6.7	32
603	Social support and suicide in Japanese men and women - the Japan Public Health Center (JPHC)-based prospective study. <i>Journal of Psychiatric Research</i> , 2011 , 45, 1545-50	5.2	32
602	Dietary isoflavone intake and breast cancer risk in case-control studies in Japanese, Japanese Brazilians, and non-Japanese Brazilians. <i>Breast Cancer Research and Treatment</i> , 2009 , 116, 401-11	4.4	32
601	Association between plasma 25-hydroxyvitamin D and colorectal adenoma according to dietary calcium intake and vitamin D receptor polymorphism. <i>American Journal of Epidemiology</i> , 2012 , 175, 236-44	2.8	32
600	Isoflavone intake and risk of gastric cancer: a population-based prospective cohort study in Japan. <i>American Journal of Clinical Nutrition</i> , 2012 , 95, 147-54	7	32
599	Serum aminotransferase level and the risk of hepatocellular carcinoma: a population-based cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2009 , 18, 26-32	2	32
598	Generalizability of relative risk estimates from a well-defined population to a general population. <i>European Journal of Epidemiology</i> , 2006 , 21, 253-62	12.1	32
597	Urinary level of 1,N(6)-ethenodeoxyadenosine, a marker of oxidative stress, is associated with salt excretion and omega 6-polyunsaturated fatty acid intake in postmenopausal Japanese women. <i>International Journal of Cancer</i> , 2002 , 100, 71-5	7.5	32
596	Dietary habits among the JPHC study participants at baseline survey. Japan Public Health Center-based Prospective Study on Cancer and Cardiovascular Diseases. <i>Journal of Epidemiology</i> , 2001 , 11, S30-43	3.4	32
595	Risk factors for stomach cancer in Brazil (II): a case-control study among Japanese Brazilians in Sã Paulo. <i>Japanese Journal of Clinical Oncology</i> , 2002 , 32, 284-90	2.8	32

594	Risk factors for stomach cancer in Brazil (I): a case-control study among non-Japanese Brazilians in S Paulo. <i>Japanese Journal of Clinical Oncology</i> , 2002 , 32, 277-83	2.8	32
593	Leisure-time physical activity and breast cancer risk defined by estrogen and progesterone receptor status--the Japan Public Health Center-based Prospective Study. <i>Preventive Medicine</i> , 2011 , 52, 227-33	4.3	31
592	Plasma levels of C-reactive protein and serum amyloid A and gastric cancer in a nested case-control study: Japan Public Health Center-based prospective study. <i>Carcinogenesis</i> , 2010 , 31, 712-8	4.6	31
591	Genetic polymorphisms in estrogen metabolism and breast cancer risk in case-control studies in Japanese, Japanese Brazilians and non-Japanese Brazilians. <i>Journal of Human Genetics</i> , 2009 , 54, 209-15	4.3	31
590	Fruits and vegetables in relation to prostate cancer in Japanese men: the Japan Public Health Center-Based Prospective Study. <i>Nutrition and Cancer</i> , 2010 , 62, 30-9	2.8	31
589	Epstein-Barr virus-associated gastric carcinoma in Japanese Brazilians and non-Japanese Brazilians in S Paulo. <i>Japanese Journal of Cancer Research</i> , 2001 , 92, 911-7		31
588	Rice consumption is not associated with risk of cardiovascular disease morbidity or mortality in Japanese men and women: a large population-based, prospective cohort study. <i>American Journal of Clinical Nutrition</i> , 2014 , 100, 199-207	7	30
587	Association between education level and dentition status in Japanese adults: Japan public health center-based oral health study. <i>Community Dentistry and Oral Epidemiology</i> , 2012 , 40, 481-7	2.8	30
586	Association between dietary heterocyclic amine levels, genetic polymorphisms of NAT2, CYP1A1, and CYP1A2 and risk of colorectal cancer: a hospital-based case-control study in Japan. <i>Scandinavian Journal of Gastroenterology</i> , 2009 , 44, 952-9	2.4	30
585	No association between fruit or vegetable consumption and the risk of colorectal cancer in Japan. <i>British Journal of Cancer</i> , 2005 , 92, 1782-4	8.7	30
584	Plasma antioxidant vitamins and carotenoids in five Japanese populations with varied mortality from gastric cancer. <i>Nutrition and Cancer</i> , 1999 , 34, 56-61	2.8	30
583	Incidence of type 2 diabetes in Japan: a systematic review and meta-analysis. <i>PLoS ONE</i> , 2013 , 8, e74699	3.7	30
582	Combined impact of five lifestyle factors and subsequent risk of cancer: the Japan Public Health Center Study. <i>Preventive Medicine</i> , 2012 , 54, 112-6	4.3	29
581	Impact of neighborhood socioeconomic conditions on the risk of stroke in Japan. <i>Journal of Epidemiology</i> , 2015 , 25, 254-60	3.4	29
580	Validity and applicability of a simple questionnaire for the estimation of total and domain-specific physical activity. <i>Diabetology International</i> , 2011 , 2, 47-54	2.3	29
579	Altered expression of the human base excision repair gene NTH1 in gastric cancer. <i>Carcinogenesis</i> , 2009 , 30, 1345-52	4.6	29
578	Education, social roles, and the risk of cardiovascular disease among middle-aged Japanese women: the JPHC Study Cohort I. <i>Stroke</i> , 2008 , 39, 2886-90	6.7	29
577	Residuals of beta-hexachlorocyclohexane, dichlorodiphenyltrichloroethane, and hexachlorobenzene in serum, and relations with consumption of dietary components in rural residents in Japan. <i>Science of the Total Environment</i> , 2002 , 286, 119-27	10.2	29

576	Food composition and empirical weight methods in predicting nutrient intakes from food frequency questionnaire. <i>Annals of Epidemiology</i> , 2001 , 11, 213-8	6.4	29
575	Observer disagreement in histological classification of ovarian tumors in Japan. <i>Gynecologic Oncology</i> , 1994 , 54, 54-8	4.9	29
574	Cancer mortality among Japanese residents of the city of Sã Paulo, Brazil. <i>International Journal of Cancer</i> , 1990 , 45, 436-9	7.5	29
573	Association of soy and fermented soy product intake with total and cause specific mortality: prospective cohort study. <i>BMJ, The</i> , 2020 , 368, m34	5.9	28
572	Dietary acrylamide intake and risk of breast cancer: The Japan Public Health Center-based Prospective Study. <i>Cancer Science</i> , 2018 , 109, 843-853	6.9	28
571	Coping strategies and risk of cardiovascular disease incidence and mortality: the Japan Public Health Center-based prospective Study. <i>European Heart Journal</i> , 2016 , 37, 890-899	9.5	28
570	Pulse pressure is an independent risk factor for stroke among middle-aged Japanese with normal systolic blood pressure: the JPHC study. <i>Journal of Hypertension</i> , 2011 , 29, 319-24	1.9	28
569	Smoking cessation: predictive factors among middle-aged Japanese. <i>Nicotine and Tobacco Research</i> , 2010 , 12, 1050-4	4.9	28
568	Alcohol consumption, social support, and risk of stroke and coronary heart disease among Japanese men: the JPHC Study. <i>Alcoholism: Clinical and Experimental Research</i> , 2009 , 33, 1025-32	3.7	28
567	Clinical significance of IgG antibody titer against <i>Helicobacter pylori</i> . <i>Helicobacter</i> , 2009 , 14, 231-6	4.9	28
566	Physical activity and colorectal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 2-13	2.8	28
565	Dietary patterns in Japanese migrants to southeastern Brazil and their descendants. <i>Journal of Epidemiology</i> , 1997 , 7, 198-204	3.4	28
564	Food consumption and gastric cancer mortality in five regions of Japan. <i>Nutrition and Cancer</i> , 1997 , 27, 60-4	2.8	28
563	Protective effect of vitamin C on oxidative stress: a randomized controlled trial. <i>International Journal for Vitamin and Nutrition Research</i> , 2008 , 78, 121-8	1.7	28
562	Decrease in risk of lung cancer death in Japanese men after smoking cessation by age at quitting: pooled analysis of three large-scale cohort studies. <i>Cancer Science</i> , 2007 , 98, 584-9	6.9	28
561	Alcohol drinking and breast cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2007 , 37, 568-74	2.8	28
560	Vegetables, fruit consumption and risk of lung cancer among middle-aged Japanese men and women: JPHC study. <i>Cancer Causes and Control</i> , 2004 , 15, 349-57	2.8	28
559	Correlation between Atrophic Gastritis Prevalence and Gastric Cancer Mortality among Middle-aged Men in 5 Areas in Japan. <i>Journal of Epidemiology</i> , 1993 , 3, 35-39	3.4	28

558	Association between green tea/coffee consumption and biliary tract cancer: A population-based cohort study in Japan. <i>Cancer Science</i> , 2016 , 107, 76-83	6.9	28
557	12 new susceptibility loci for prostate cancer identified by genome-wide association study in Japanese population. <i>Nature Communications</i> , 2019 , 10, 4422	17.4	27
556	Impact of Alcohol Intake and Drinking Patterns on Mortality From All Causes and Major Causes of Death in a Japanese Population. <i>Journal of Epidemiology</i> , 2018 , 28, 140-148	3.4	27
555	Development of a Risk Equation for the Incidence of Coronary Artery Disease and Ischemic Stroke for Middle-Aged Japanese - Japan Public Health Center-Based Prospective Study. <i>Circulation Journal</i> , 2016 , 80, 1386-95	2.9	27
554	The combination of Helicobacter pylori- and cytotoxin-associated gene-A seropositivity in relation to the risk of myocardial infarction in middle-aged Japanese: The Japan Public Health Center-based study. <i>Atherosclerosis</i> , 2013 , 230, 67-72	3.1	27
553	Magnesium intake and type II diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study. <i>European Journal of Clinical Nutrition</i> , 2010 , 64, 1244-7	5.2	27
552	Cigarette smoking and pancreas cancer risk: an evaluation based on a systematic review of epidemiologic evidence in the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 1292-302	2.8	27
551	Fasting plasma glucose and 5-year incidence of diabetes in the JPHC diabetes study - suggestion for the threshold for impaired fasting glucose among Japanese. <i>Endocrine Journal</i> , 2010 , 57, 629-37	2.9	27
550	Socioeconomic status and gastric cancer survival in Japan. <i>Gastric Cancer</i> , 2010 , 13, 222-30	7.6	27
549	Green tea intake in relation to serum lipid levels in Middle-aged Japanese men and women. <i>Annals of Epidemiology</i> , 1997 , 7, 280-4	6.4	27
548	Negative psychological aspects and survival in lung cancer patients. <i>Psycho-Oncology</i> , 2008 , 17, 466-73	3.9	27
547	Impact of tobacco smoking on subsequent cancer risk among middle-aged Japanese men and women: data from a large-scale population-based cohort study in Japan--the JPHC study. <i>Preventive Medicine</i> , 2004 , 38, 516-22	4.3	27
546	The impact of C-reactive protein on risk of stroke, stroke subtypes, and ischemic heart disease in middle-aged Japanese: the Japan public health center-based study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012 , 19, 756-66	4	27
545	The association between midlife serum high-density lipoprotein and mild cognitive impairment and dementia after 19 years of follow-up. <i>Translational Psychiatry</i> , 2019 , 9, 26	8.6	26
544	Hepatitis B and C virus infection and risk of lymphoid malignancies: A population-based cohort study (JPHC Study). <i>Cancer Epidemiology</i> , 2015 , 39, 562-6	2.8	26
543	Dietary pattern and breast cancer risk in Japanese women: the Japan Public Health Center-based Prospective Study (JPHC Study). <i>British Journal of Nutrition</i> , 2016 , 115, 1769-79	3.6	26
542	Report of the JDS/JCA Joint Committee on Diabetes and Cancer. <i>Diabetology International</i> , 2013 , 4, 81-96	9.3	26
541	Association of high-density lipoprotein cholesterol concentration with different types of stroke and coronary heart disease: The Japan Public Health Center-based prospective (JPHC) study. <i>Atherosclerosis</i> , 2017 , 265, 147-154	3.1	26

540	Dietary patterns and type 2 diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study. <i>European Journal of Clinical Nutrition</i> , 2013 , 67, 18-24	5.2	26
539	Diabetes and the risk of coronary heart disease in the general Japanese population: the Japan Public Health Center-based prospective (JPHC) study. <i>Atherosclerosis</i> , 2011 , 216, 187-91	3.1	26
538	Plasma testosterone and sex hormone-binding globulin concentrations and the risk of prostate cancer among Japanese men: a nested case-control study. <i>Cancer Science</i> , 2010 , 101, 2652-7	6.9	26
537	A randomized controlled trial for chemoprevention of gastric cancer in high-risk Japanese population; study design, feasibility and protocol modification. <i>Japanese Journal of Cancer Research</i> , 1997 , 88, 344-9		26
536	Screening of 214 single nucleotide polymorphisms in 44 candidate cancer susceptibility genes: a case-control study on gastric and colorectal cancers in the Japanese population. <i>American Journal of Gastroenterology</i> , 2008 , 103, 1476-87	0.7	26
535	Validity of a self-administered food frequency questionnaire in the assessment of heterocyclic amine intake using 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) levels in hair. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2007 , 630, 14-9	3	26
534	Validity of the self-administered food frequency questionnaire used in the 5-year follow-up survey for the JPHC Study to assess folate, vitamin B6 and B12 intake: comparison with dietary records and blood level. <i>Journal of Epidemiology</i> , 2003 , 13, S98-101	3.4	26
533	Smoking and risk of premature death among middle-aged Japanese: ten-year follow-up of the Japan Public Health Center-based prospective study on cancer and cardiovascular diseases (JPHC Study) cohort I. <i>Japanese Journal of Cancer Research</i> , 2002 , 93, 6-14		26
532	Association of habitual smoking and drinking with single nucleotide polymorphism (SNP) in 40 candidate genes: data from random population-based Japanese samples. <i>Journal of Human Genetics</i> , 2005 , 50, 62-68	4.3	26
531	Histopathological modifications of early atherosclerotic lesions by risk factors--findings in PDAY subjects. <i>Atherosclerosis</i> , 2001 , 156, 389-99	3.1	26
530	Serum n-3 fatty acids, fish consumption and cancer mortality in six Japanese populations in Japan and Brazil. <i>Japanese Journal of Cancer Research</i> , 1999 , 90, 914-21		26
529	Smoking, occupation and family history in lung cancer patients under fifty years of age. <i>Japanese Journal of Clinical Oncology</i> , 1987 , 17, 309-17	2.8	26
528	Association of leisure-time physical activity with total and cause-specific mortality: a pooled analysis of nearly a half million adults in the Asia Cohort Consortium. <i>International Journal of Epidemiology</i> , 2018 , 47, 771-779	7.8	25
527	Hepatitis B and C Virus Infection and Risk of Pancreatic Cancer: A Population-Based Cohort Study (JPHC Study Cohort II). <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2016 , 25, 555-7	4	25
526	Green tea consumption and breast cancer risk in Japanese women: a case-control study. <i>Nutrition and Cancer</i> , 2014 , 66, 57-67	2.8	25
525	Dietary arsenic intake and subsequent risk of cancer: the Japan Public Health Center-based (JPHC) Prospective Study. <i>Cancer Causes and Control</i> , 2013 , 24, 1403-15	2.8	25
524	Long-term exposure to particulate matter in relation to mortality and incidence of cardiovascular disease: the JPHC Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013 , 20, 296-309	4	25
523	Methionine synthase A2756G polymorphism interacts with alcohol and folate intake to influence the risk of colorectal adenoma. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2009 , 18, 267-74	4	25

522	Gender difference in the association of insulin and the insulin-like growth factor axis with colorectal neoplasia. <i>International Journal of Obesity</i> , 2012 , 36, 440-7	5.5	25
521	Plasma organochlorine levels and subsequent risk of breast cancer among Japanese women: a nested case-control study. <i>Science of the Total Environment</i> , 2008 , 402, 176-83	10.2	25
520	Interaction between cytochrome P450 gene polymorphisms and serum organochlorine TEQ levels in the risk of endometriosis. <i>Molecular Human Reproduction</i> , 2007 , 13, 399-404	4.4	25
519	Seasonal misclassification error and magnitude of true between-person variation in dietary nutrient intake: a random coefficients analysis and implications for the Japan Public Health Center (JPHC) Cohort Study. <i>Public Health Nutrition</i> , 2003 , 6, 385-91	3.3	25
518	High hemoglobin A1c levels within the non-diabetic range are associated with the risk of all cancers. <i>International Journal of Cancer</i> , 2016 , 138, 1741-53	7.5	25
517	Cigarette smoking and cervical cancer risk: an evaluation based on a systematic review and meta-analysis among Japanese women. <i>Japanese Journal of Clinical Oncology</i> , 2019 , 49, 77-86	2.8	25
516	Why has Japan become the world's most long-lived country: insights from a food and nutrition perspective. <i>European Journal of Clinical Nutrition</i> , 2021 , 75, 921-928	5.2	25
515	Novel epigenetic markers for gastric cancer risk stratification in individuals after <i>Helicobacter pylori</i> eradication. <i>Gastric Cancer</i> , 2018 , 21, 745-755	7.6	24
514	Cigarette smoking and bladder cancer risk: an evaluation based on a systematic review of epidemiologic evidence in the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2016 , 46, 273-283	2.8	24
513	Alcohol consumption and risk of stroke and coronary heart disease among Japanese women: the Japan Public Health Center-based prospective study. <i>Preventive Medicine</i> , 2013 , 57, 505-10	4.3	24
512	Association of gastric cancer risk factors with DNA methylation levels in gastric mucosa of healthy Japanese: a cross-sectional study. <i>Carcinogenesis</i> , 2015 , 36, 1291-8	4.6	24
511	Impact of viral load of hepatitis C on the incidence of hepatocellular carcinoma: A population-based cohort study (JPHC Study). <i>Cancer Letters</i> , 2011 , 300, 173-9	9.9	24
510	Association of alcohol intake with the risk of malignant lymphoma and plasma cell myeloma in Japanese: a population-based cohort study (Japan Public Health Center-based Prospective Study). <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2010 , 19, 429-34	4	24
509	Gender difference in the association between social support and metabolic syndrome in Japan: the 'enkai' effect?. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, 71-7	5.1	24
508	Impact of weight change on specific-cause mortality among middle-aged Japanese individuals. <i>Journal of Epidemiology and Community Health</i> , 2009 , 63, 447-54	5.1	24
507	Association between dietary heterocyclic amine levels, genetic polymorphisms of NAT2, CYP1A1, and CYP1A2 and risk of stomach cancer: a hospital-based case-control study in Japan. <i>Gastric Cancer</i> , 2009 , 12, 198-205	7.6	24
506	Dietary isoflavone and the risk of colorectal adenoma: a case-control study in Japan. <i>British Journal of Cancer</i> , 2009 , 100, 1812-6	8.7	24
505	Education in relation to incidence of and mortality from cancer and cardiovascular disease in Japan. <i>European Journal of Public Health</i> , 2008 , 18, 466-72	2.1	24

504	Female reproductive factors and the risk of gastric cancer in a large-scale population-based cohort study in Japan (JPHC study). <i>European Journal of Cancer Prevention</i> , 2008 , 17, 345-53	2	24
503	Analysis of mutagenic heterocyclic amines in cooked food samples by gas chromatography with nitrogen-phosphorus detector. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2002 , 69, 682-9	2.7	24
502	Association between mortality and incidence rates of coronary heart disease and stroke: The Japan Public Health Center-based prospective (JPHC) study. <i>International Journal of Cardiology</i> , 2016 , 222, 281-286	3.2	24
501	Cruciferous Vegetable Intake Is Inversely Associated with Lung Cancer Risk among Current Nonsmoking Men in the Japan Public Health Center (JPHC) Study. <i>Journal of Nutrition</i> , 2017 , 147, 841-849 ¹	4.1	23
500	Perceived stress level and risk of cancer incidence in a Japanese population: the Japan Public Health Center (JPHC)-based Prospective Study. <i>Scientific Reports</i> , 2017 , 7, 12964	4.9	23
499	Circulating sex hormones in relation to anthropometric, sociodemographic and behavioural factors in an international dataset of 12,300 men. <i>PLoS ONE</i> , 2017 , 12, e0187741	3.7	23
498	Impact of Moderate-Intensity and Vigorous-Intensity Physical Activity on Mortality. <i>Medicine and Science in Sports and Exercise</i> , 2018 , 50, 715-721	1.2	23
497	Self-reported taste preference can be a proxy for daily sodium intake in middle-aged Japanese adults. <i>Journal of the Academy of Nutrition and Dietetics</i> , 2014 , 114, 781-7	3.9	23
496	Diabetes mellitus and liver cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2014 , 44, 986-99	2.8	23
495	Fragment c gamma receptor gene polymorphisms and breast cancer risk in case-control studies in Japanese, Japanese Brazilians, and non-Japanese Brazilians. <i>Breast Cancer Research and Treatment</i> , 2011 , 126, 497-505	4.4	23
494	High dietary intake of magnesium may decrease risk of colorectal cancer in Japanese men. <i>Journal of Nutrition</i> , 2010 , 140, 779-85	4.1	23
493	Secular trends in cancer mortality among Japanese immigrants in the state of Sã Paulo, Brazil, 1979-2001. <i>European Journal of Cancer Prevention</i> , 2008 , 17, 1-8	2	23
492	A population-based dietary intervention trial in a high-risk area for stomach cancer and stroke: changes in intakes and related biomarkers. <i>Preventive Medicine</i> , 2003 , 37, 432-41	4.3	23
491	Vasodilation by intrathecal lipopolysaccharide of the cerebral arteries after subarachnoid haemorrhage in dogs. <i>Acta Neurochirurgica</i> , 1996 , 138, 330-7	3	23
490	Cross-sectional Epidemiologic Study for Assessing Cancer Risks at the Population Level. <i>Journal of Epidemiology</i> , 1992 , 2, 83-89	3.4	23
489	Alcohol consumption and risk of esophageal cancer in Japan: a case-control study in seven hospitals. <i>Japanese Journal of Clinical Oncology</i> , 1994 , 24, 241-6	2.8	23
488	Genome-wide association study identified new susceptible genetic variants in HLA class I region for hepatitis B virus-related hepatocellular carcinoma. <i>Scientific Reports</i> , 2018 , 8, 7958	4.9	23
487	Dietary acid load and mortality among Japanese men and women: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2017 , 106, 146-154	7	22

486	Dietary heterocyclic amine intake, NAT2 genetic polymorphism, and colorectal adenoma risk: the colorectal adenoma study in Tokyo. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2015 , 24, 613-20	4	22
485	Vegetable consumption and colorectal cancer risk: an evaluation based on a systematic review and meta-analysis among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2015 , 45, 973-9	2.8	22
484	Genome-wide association study in East Asians identifies two novel breast cancer susceptibility loci. <i>Human Molecular Genetics</i> , 2016 , 25, 3361-3371	5.6	22
483	Death by suicide and other externally caused injuries after stroke in Japan (1990-2010): the Japan Public Health Center-based prospective study. <i>Psychosomatic Medicine</i> , 2014 , 76, 452-9	3.7	22
482	Prediagnostic plasma antibody levels to periodontopathic bacteria and risk of coronary heart disease. <i>International Heart Journal</i> , 2012 , 53, 209-14	1.8	22
481	Taste preferences and body weight change in Japanese adults: the JPHC Study. <i>International Journal of Obesity</i> , 2009 , 33, 1191-7	5.5	22
480	Relationship of smoking and smoking cessation with number of teeth present: JPHC Oral Health Study*. <i>Oral Diseases</i> , 2009 , 15, 69-75	3.5	22
479	Lung cancer risk and consumption of vegetables and fruit: an evaluation based on a systematic review of epidemiological evidence from Japan. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 693-708 ^{2.8}	2.8	22
478	Histologic grade and p53 immunoreaction as indicators of early recurrence of node-negative breast cancer. <i>Japanese Journal of Clinical Oncology</i> , 1997 , 27, 6-12	2.8	22
477	Alcohol drinking and liver cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2008 , 38, 816-38	2.8	22
476	Susceptibility to lung cancer and genetic polymorphisms in the alcohol metabolite-related enzymes alcohol dehydrogenase 3, aldehyde dehydrogenase 2, and cytochrome P450 2E1 in the Japanese population. <i>Cancer</i> , 2007 , 110, 353-62	6.4	22
475	Marital status and non-small cell lung cancer survival: the Lung Cancer Database Project in Japan. <i>Psycho-Oncology</i> , 2008 , 17, 869-76	3.9	22
474	Association between CDH1 haplotypes and gastric cancer risk in a Japanese population. <i>Scandinavian Journal of Gastroenterology</i> , 2007 , 42, 1479-85	2.4	22
473	2-Amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) level in human hair as biomarkers for dietary grilled/stir-fried meat and fish intake. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2005 , 588, 136-42	3	22
472	Association of body mass index and height with risk of prostate cancer among middle-aged Japanese men. <i>British Journal of Cancer</i> , 2006 , 94, 740-2	8.7	22
471	Validity of a self-administered food frequency questionnaire used in the 5-year follow-up survey of the JPHC Study Cohort I to assess dietary fiber intake: comparison with dietary records. <i>Journal of Epidemiology</i> , 2003 , 13, S106-14	3.4	22
470	Prediction of the gastric cancer mortality in 2000 in Japan. <i>Japanese Journal of Cancer Research</i> , 1988 , 79, 439-44		22
469	Genome-wide association study identifies gastric cancer susceptibility loci at 12q24.11-12 and 20q11.21. <i>Cancer Science</i> , 2018 , 109, 4015-4024	6.9	22

468	Body-Mass Index and Pancreatic Cancer Incidence: A Pooled Analysis of Nine Population-Based Cohort Studies With More Than 340,000 Japanese Subjects. <i>Journal of Epidemiology</i> , 2018 , 28, 245-252	3.4	22
467	Association of Ile462Val (Exon 7) polymorphism of cytochrome P450 IA1 with lung cancer in the Asian population: further evidence from a case-control study in Okinawa. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 1998 , 7, 413-7	4	22
466	A Pooled Analysis of 15 Prospective Cohort Studies on the Association between Fruit, Vegetable, and Mature Bean Consumption and Risk of Prostate Cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2017 , 26, 1276-1287	4	21
465	Seaweed intake and risk of cardiovascular disease: the Japan Public Health Center-based Prospective (JPHC) Study. <i>American Journal of Clinical Nutrition</i> , 2019 , 110, 1449-1455	7	21
464	Personality traits and coping styles explain anxiety in lung cancer patients to a greater extent than other factors. <i>Japanese Journal of Clinical Oncology</i> , 2015 , 45, 456-63	2.8	21
463	Association between Chlamydomphila pneumoniae infection and risk of coronary heart disease for Japanese: the JPHC study. <i>Atherosclerosis</i> , 2014 , 233, 338-342	3.1	21
462	Green tea and coffee consumption and its association with thyroid cancer risk: a population-based cohort study in Japan. <i>Cancer Causes and Control</i> , 2011 , 22, 985-93	2.8	21
461	Body mass index and subsequent risk of kidney cancer: a prospective cohort study in Japan. <i>Annals of Epidemiology</i> , 2010 , 20, 466-72	6.4	21
460	Nutrient consumption and gastric cancer mortality in five regions of Japan. <i>Nutrition and Cancer</i> , 1997 , 27, 310-5	2.8	21
459	Plasma folate and risk of colorectal cancer in a nested case-control study: the Japan Public Health Center-based prospective study. <i>Cancer Causes and Control</i> , 2008 , 19, 67-74	2.8	21
458	Analytical method of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine in human hair by column-switching liquid chromatography-mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2004 , 803, 209-13	3.2	21
457	Food/nutrient intake and risk of atrophic gastritis among the Helicobacter pylori-infected population of northeastern Japan. <i>Cancer Science</i> , 2003 , 94, 372-7	6.9	21
456	Validity and reproducibility of a self-administered questionnaire to determine dietary supplement users among Japanese. <i>European Journal of Clinical Nutrition</i> , 2001 , 55, 360-5	5.2	21
455	Differential effects of cigarette smoking and alcohol consumption on the plasma levels of carotenoids in middle-aged Japanese men. <i>Japanese Journal of Cancer Research</i> , 1996 , 87, 563-9		21
454	Coffee drinking and colorectal cancer risk: an evaluation based on a systematic review and meta-analysis among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2016 , 46, 781-7	2.8	21
453	Physical inactivity, prolonged sedentary behaviors, and use of visual display terminals as potential risk factors for dry eye disease: JPHC-NEXT study. <i>Ocular Surface</i> , 2020 , 18, 56-63	6.5	21
452	Plasma cytokine levels and the presence of colorectal cancer. <i>PLoS ONE</i> , 2019 , 14, e0213602	3.7	20
451	Fiber intake and risk of subsequent prostate cancer in Japanese men. <i>American Journal of Clinical Nutrition</i> , 2015 , 101, 118-25	7	20

450	Validity of a Self-Administered Food-Frequency Questionnaire for Assessing Amino Acid Intake in Japan: Comparison With Intake From 4-Day Weighed Dietary Records and Plasma Levels. <i>Journal of Epidemiology</i> , 2016 , 26, 36-44	3.4	20
449	Hemoglobin a1c levels and the risk of cardiovascular disease in people without known diabetes: a population-based cohort study in Japan. <i>Medicine (United States)</i> , 2015 , 94, e785	1.8	20
448	Diagnosed diabetes and premature death among middle-aged Japanese: results from a large-scale population-based cohort study in Japan (JPHC study). <i>BMJ Open</i> , 2015 , 5, e007736	3	20
447	Soy food and isoflavone intake and endometrial cancer risk: the Japan Public Health Center-based prospective study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2015 , 122, 304-11	3.7	20
446	Association of body mass index and risk of death from pancreas cancer in Asians: findings from the Asia Cohort Consortium. <i>European Journal of Cancer Prevention</i> , 2013 , 22, 244-50	2	20
445	A population-based cohort study for the risk factors of HCC among hepatitis B virus mono-infected subjects in Japan. <i>Journal of Gastroenterology</i> , 2011 , 46, 117-24	6.9	20
444	Leisure-time physical activity and breast cancer risk by hormone receptor status: effective life periods and exercise intensity. <i>Cancer Causes and Control</i> , 2010 , 21, 1787-98	2.8	20
443	Reduced life expectancy due to smoking in large-scale cohort studies in Japan. <i>Journal of Epidemiology</i> , 2008 , 18, 111-8	3.4	20
442	Reproductive factors, exogenous female hormone use and colorectal cancer risk: the Japan Public Health Center-based Prospective Study. <i>European Journal of Cancer Prevention</i> , 2008 , 17, 515-24	2	20
441	Validity of a self-administered food frequency questionnaire used in the 5-year follow-up survey of the JPHC Study Cohort I to assess carotenoids and vitamin C intake: comparison with dietary records and blood level. <i>Journal of Epidemiology</i> , 2003 , 13, S82-91	3.4	20
440	Cancer mortality among Japanese immigrants and their descendants in the state of So Paulo, Brazil, 1999-2001. <i>Japanese Journal of Clinical Oncology</i> , 2004 , 34, 673-80	2.8	20
439	Validating the dietary inflammatory index using inflammatory biomarkers in a Japanese population: A cross-sectional study of the JPHC-FFQ validation study. <i>Nutrition</i> , 2020 , 69, 110569	4.8	20
438	Changes in the Employment Status and Risk of Stroke and Stroke Types. <i>Stroke</i> , 2017 , 48, 1176-1182	6.7	19
437	Coffee drinking and colorectal cancer and its subsites: A pooled analysis of 8 cohort studies in Japan. <i>International Journal of Cancer</i> , 2018 , 143, 307-316	7.5	19
436	Coffee and green tea consumption in relation to brain tumor risk in a Japanese population. <i>International Journal of Cancer</i> , 2016 , 139, 2714-2721	7.5	19
435	Coffee intake and the risk of colorectal adenoma: The colorectal adenoma study in Tokyo. <i>International Journal of Cancer</i> , 2015 , 137, 463-70	7.5	19
434	Genetic markers of immunoglobulin G and susceptibility to breast cancer. <i>Human Immunology</i> , 2012 , 73, 1155-8	2.3	19
433	Calcium intake and the 10-year incidence of self-reported vertebral fractures in women and men: the Japan Public Health Centre-based Prospective Study. <i>British Journal of Nutrition</i> , 2009 , 101, 285-94	3.6	19

432	Different effects of hypertension and hypercholesterolemia on the natural history of aortic atherosclerosis by the stage of intimal lesions. <i>Atherosclerosis</i> , 1997 , 128, 85-95	3.1	19
431	Different etiological role of <i>Helicobacter pylori</i> (Hp) infection in carcinogenesis between differentiated and undifferentiated gastric cancers: a nested case-control study using IgG titer against Hp surface antigen. <i>Acta Oncologica</i> , 2008 , 47, 360-5	3.2	19
430	Four-year follow-up of effects of toluene diisocyanate exposure on the respiratory system in polyurethane foam manufacturing workers. II. Four-year changes in the effects on the respiratory system. <i>International Archives of Occupational and Environmental Health</i> , 1992 , 63, 565-9	3.2	19
429	Mortality and cause of death among first-generation Japanese in Sã Paulo, Brazil. <i>International Journal of Epidemiology</i> , 1989 , 18, 647-51	7.8	19
428	Cigarette smoking and the risk of head and neck cancer in the Japanese population: a systematic review and meta-analysis. <i>Japanese Journal of Clinical Oncology</i> , 2016 , 46, 580-95	2.8	19
427	Circulating sex hormone levels and colorectal cancer risk in Japanese postmenopausal women: The JPHC nested case-control study. <i>International Journal of Cancer</i> , 2019 , 145, 1238-1244	7.5	18
426	Evidence-based cancer prevention recommendations for Japanese. <i>Japanese Journal of Clinical Oncology</i> , 2018 , 48, 576-586	2.8	18
425	IGKC and FcR genotypes and humoral immunity to HER2 in breast cancer. <i>Immunobiology</i> , 2014 , 219, 113-7	3.4	18
424	Fish consumption and colorectal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2013 , 43, 935-41	2.8	18
423	Zinc and heme iron intakes and risk of colorectal cancer: a population-based prospective cohort study in Japan. <i>American Journal of Clinical Nutrition</i> , 2012 , 96, 864-73	7	18
422	Pathway analyses identify TGFBR2 as potential breast cancer susceptibility gene: results from a consortium study among Asians. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2012 , 21, 1176-84	4	18
421	Determination of sub-ppb cadmium in urine by solid-phase extraction and inductively coupled plasma-mass spectrometry. <i>Analytical Sciences</i> , 2008 , 24, 1049-52	1.7	18
420	Development of an automated 3D segmentation program for volume quantification of body fat distribution using CT. <i>Japanese Journal of Radiological Technology</i> , 2008 , 64, 1177-81		18
419	Long-term vitamin C supplementation has no markedly favourable effect on serum lipids in middle-aged Japanese subjects. <i>British Journal of Nutrition</i> , 2004 , 91, 81-90	3.6	18
418	<i>Helicobacter pylori</i> infection and atrophic gastritis in middle-aged Japanese residents of Sã Paulo and Lima. <i>International Journal of Epidemiology</i> , 1999 , 28, 577-82	7.8	18
417	Sustained Weight Loss and Risk of Breast Cancer in Women 50 Years and Older: A Pooled Analysis of Prospective Data. <i>Journal of the National Cancer Institute</i> , 2020 , 112, 929-937	9.7	18
416	Quantitative Assessment of the Retina Using OCT and Associations with Cognitive Function. <i>Ophthalmology</i> , 2020 , 127, 107-118	7.3	18
415	Dietary acrylamide intake and the risk of endometrial or ovarian cancers in Japanese women. <i>Cancer Science</i> , 2018 , 109, 3316-3325	6.9	18

414	Working Hours and Risk of Acute Myocardial Infarction and Stroke Among Middle-Aged Japanese Men - The Japan Public Health Center-Based Prospective Study Cohort II. <i>Circulation Journal</i> , 2019 , 83, 1072-1079	2.9	17
413	The association between adult attained height and sitting height with mortality in the European Prospective Investigation into Cancer and Nutrition (EPIC). <i>PLoS ONE</i> , 2017 , 12, e0173117	3.7	17
412	Dietary consumption of antioxidant vitamins and subsequent lung cancer risk: The Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2018 , 142, 2441-2460	7.5	17
411	Coping behaviors and suicide in the middle-aged and older Japanese general population: the Japan Public Health Center-based Prospective Study. <i>Annals of Epidemiology</i> , 2014 , 24, 199-205	6.4	17
410	Inclusion of a Genetic Risk Score into a Validated Risk Prediction Model for Colorectal Cancer in Japanese Men Improves Performance. <i>Cancer Prevention Research</i> , 2017 , 10, 535-541	3.2	17
409	Dietary intake of antioxidant vitamins and risk of stroke: the Japan Public Health Center-based Prospective Study. <i>European Journal of Clinical Nutrition</i> , 2017 , 71, 1179-1185	5.2	17
408	Socioeconomic status inconsistency and risk of stroke among Japanese middle-aged women. <i>Stroke</i> , 2014 , 45, 2592-8	6.7	17
407	Dietary isoflavone intake, polymorphisms in the CYP17, CYP19, 17beta-HSD1, and SHBG genes, and risk of breast cancer in case-control studies in Japanese, Japanese Brazilians, and non-Japanese Brazilians. <i>Nutrition and Cancer</i> , 2010 , 62, 466-75	2.8	17
406	Validity of a self-administered food frequency questionnaire used in the 5-year follow-up survey of the JPHC Study to assess selenium intake: comparison with dietary records and blood levels. <i>Journal of Epidemiology</i> , 2003 , 13, S92-7	3.4	17
405	Validity of a self-administered food frequency questionnaire in the 5-year follow-up survey of the JPHC Study Cohort I to assess sodium and potassium intake: comparison with dietary records and 24-hour urinary excretion level. <i>Journal of Epidemiology</i> , 2003 , 13, S102-5	3.4	17
404	Risk factors for lung cancer in Rio de Janeiro, Brazil: a case-control study. <i>Lung Cancer</i> , 1994 , 11, 179-90	5.9	17
403	Occupational sitting time and risk of all-cause mortality among Japanese workers. <i>Scandinavian Journal of Work, Environment and Health</i> , 2015 , 41, 519-28	4.3	17
402	Measures of body fatness and height in early and mid-to-late adulthood and prostate cancer: risk and mortality in The Pooling Project of Prospective Studies of Diet and Cancer. <i>Annals of Oncology</i> , 2020 , 31, 103-114	10.3	17
401	Adjustment of Cell-Type Composition Minimizes Systematic Bias in Blood DNA Methylation Profiles Derived by DNA Collection Protocols. <i>PLoS ONE</i> , 2016 , 11, e0147519	3.7	17
400	Vitamin D Receptor Gene Polymorphism and the Risk of Colorectal Cancer: A Nested Case-Control Study. <i>PLoS ONE</i> , 2016 , 11, e0164648	3.7	17
399	A Collaborative Analysis of Individual Participant Data from 19 Prospective Studies Assesses Circulating Vitamin D and Prostate Cancer Risk. <i>Cancer Research</i> , 2019 , 79, 274-285	10.1	17
398	Smoking, Alcohol, and Biliary Tract Cancer Risk: A Pooling Project of 26 Prospective Studies. <i>Journal of the National Cancer Institute</i> , 2019 , 111, 1263-1278	9.7	16
397	Dietary Acrylamide Intake and Risk of Esophageal, Gastric, and Colorectal Cancer: The Japan Public Health Center-Based Prospective Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2019 , 28, 1461-1468	4.4	16

396	Novel Risk Loci Identified in a Genome-Wide Association Study of Urolithiasis in a Japanese Population. <i>Journal of the American Society of Nephrology: JASN</i> , 2019 , 30, 855-864	12.7	16
395	Identification of novel breast cancer susceptibility loci in meta-analyses conducted among Asian and European descendants. <i>Nature Communications</i> , 2020 , 11, 1217	17.4	16
394	Validity of self-reported tooth counts and masticatory status study of a Japanese adult population. <i>Journal of Oral Rehabilitation</i> , 2018 , 45, 393-398	3.4	16
393	Cancer in Japan: Prevalence, prevention and the role of heterocyclic amines in human carcinogenesis. <i>Genes and Environment</i> , 2016 , 38, 16	2.8	16
392	Plasma tea catechins and risk of cardiovascular disease in middle-aged Japanese subjects: The JPHC study. <i>Atherosclerosis</i> , 2018 , 277, 90-97	3.1	16
391	Plasma isoflavones and risk of primary liver cancer in Japanese women and men with hepatitis virus infection: a nested case-control study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2015 , 24, 532-7	4	16
390	Modification of the excess risk of coronary heart disease due to smoking by seafood/fish intake. <i>American Journal of Epidemiology</i> , 2014 , 179, 1173-81	3.8	16
389	Association of anthropometric characteristics with the risk of malignant lymphoma and plasma cell myeloma in a Japanese population: a population-based cohort study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2010 , 19, 1623-31	4	16
388	Development of a quantitative food frequency questionnaire for assessing food, nutrient, and heterocyclic aromatic amines intake in Japanese Brazilians for a colorectal adenoma case-control study. <i>International Journal of Food Sciences and Nutrition</i> , 2009 , 60 Suppl 7, 128-39	3.7	16
387	Breastfeeding and breast cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 124-30	2.8	16
386	The lung cancer database project at the National Cancer Center, Japan: study design, corresponding rate and profiles of cohort. <i>Japanese Journal of Clinical Oncology</i> , 2006 , 36, 280-4	2.8	16
385	Prognostic significance of histopathological subtypes in stage I pure yolk sac tumour of the ovary. <i>British Journal of Cancer</i> , 1994 , 69, 529-36	8.7	16
384	Smoking, Serology, and Gastric Cancer Risk in Prospective Studies from China, Japan, and Korea. <i>Cancer Prevention Research</i> , 2019 , 12, 667-674	3.2	16
383	Education and gastric cancer risk-An individual participant data meta-analysis in the StoP project consortium. <i>International Journal of Cancer</i> , 2020 , 146, 671-681	7.5	16
382	Validity of a Self-administered Food Frequency Questionnaire for the Estimation of Acrylamide Intake in the Japanese Population: The JPHC FFQ Validation Study. <i>Journal of Epidemiology</i> , 2018 , 28, 482-487	3.4	16
381	Circulating inflammatory markers and colorectal cancer risk: A prospective case-cohort study in Japan. <i>International Journal of Cancer</i> , 2018 , 143, 2767-2776	7.5	15
380	The Murakami Cohort Study of vitamin D for the prevention of musculoskeletal and other age-related diseases: a study protocol. <i>Environmental Health and Preventive Medicine</i> , 2018 , 23, 28	4.2	15
379	Differences in suicide risk according to living arrangements in Japanese men and women--the Japan Public Health Center-based (JPHC) prospective study. <i>Journal of Affective Disorders</i> , 2011 , 131, 113-9	6.6	15

378	Alcohol drinking and lung cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2007 , 37, 168-74	2.8	15
377	Ethnic difference in serology of <i>Helicobacter pylori</i> CagA between Japanese and non-Japanese Brazilians for non-cardia gastric cancer. <i>Cancer Science</i> , 2003 , 94, 64-9	6.9	15
376	Effect of smoking on serum pepsinogen I level depends on serological status of <i>Helicobacter pylori</i> . <i>Japanese Journal of Cancer Research</i> , 2001 , 92, 243-8		15
375	Effects of three-month oral supplementation of beta-carotene and vitamin C on serum concentrations of carotenoids and vitamins in middle-aged subjects: a pilot study for a randomized controlled trial to prevent gastric cancer in high-risk Japanese population. <i>Japanese Journal of Cancer Research</i> , 2000 , 91, 464-70		15
374	Association between smoking and tumor progression in Japanese women with adenocarcinoma of the lung. <i>Japanese Journal of Cancer Research</i> , 1999 , 90, 129-35		15
373	Prognostic value of p53 protein accumulation in cancer cell nuclei in adenocarcinoma of the uterine cervix. <i>Japanese Journal of Cancer Research</i> , 1995 , 86, 1049-53		15
372	Adenocarcinoma of the lung with selective metastasis to the lung: clinical, histologic and DNA-cytofluorometric analyses. <i>Japanese Journal of Cancer Research</i> , 1992 , 83, 93-100		15
371	Work-family conflict and self-rated health among Japanese workers: How household income modifies associations. <i>PLoS ONE</i> , 2017 , 12, e0169903	3.7	15
370	CYP1A1, GSTM1 and GSTT1 genetic polymorphisms and gastric cancer risk among Japanese: A nested case-control study within a large-scale population-based prospective study. <i>International Journal of Cancer</i> , 2016 , 139, 759-68	7.5	15
369	Receiver operating characteristic analysis of prediction for gastric cancer development using serum pepsinogen and <i>Helicobacter pylori</i> antibody tests. <i>BMC Cancer</i> , 2017 , 17, 183	4.8	14
368	High serum total cholesterol is associated with suicide mortality in Japanese women. <i>Acta Psychiatrica Scandinavica</i> , 2017 , 136, 259-268	6.5	14
367	<i>Helicobacter pylori</i> infection, atrophic gastritis, and risk of pancreatic cancer: A population-based cohort study in a large Japanese population: the JPHC Study. <i>Scientific Reports</i> , 2019 , 9, 6099	4.9	14
366	Genome-wide association meta-analysis identifies GP2 gene risk variants for pancreatic cancer. <i>Nature Communications</i> , 2020 , 11, 3175	17.4	14
365	Alcohol, smoking, and obesity epidemiology in Japan. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2012 , 27 Suppl 2, 121-6	4	14
364	Interaction between Red Meat Intake and NAT2 Genotype in Increasing the Risk of Colorectal Cancer in Japanese and African Americans. <i>PLoS ONE</i> , 2015 , 10, e0144955	3.7	14
363	Endometrial cancer in relation to coffee, tea, and caffeine consumption: a prospective cohort study among middle-aged women in Sweden. <i>Nutrition and Cancer</i> , 2014 , 66, 1132-43	2.8	14
362	Alcohol and risk of lung cancer among Japanese men: data from a large-scale population-based cohort study, the JPHC study. <i>Cancer Causes and Control</i> , 2008 , 19, 1095-102	2.8	14
361	Bowel movement, state of stool, and subsequent risk for colorectal cancer: the Japan public health center-based prospective study. <i>Annals of Epidemiology</i> , 2006 , 16, 888-94	6.4	14

360	Serological Immunoglobulin G antibody titers to <i>Helicobacter pylori</i> in Japanese Brazilian and Non-Japanese Brazilian gastric cancer patients and controls in S̃ Paulo. <i>Japanese Journal of Cancer Research</i> , 2001 , 92, 829-35		14
359	Sensitivity of self-reports of cancer in a population-based prospective study: JPHC Study Cohort I. <i>Journal of Clinical Epidemiology</i> , 2001 , 54, 741-6	5.7	14
358	Prognostic significance of accumulation of gene and chromosome alterations and histological grade in node-negative breast carcinoma. <i>Japanese Journal of Clinical Oncology</i> , 1998 , 28, 5-11	2.8	14
357	Follow up of women with simple ovarian cysts detected by transvaginal sonography in the Tokyo metropolitan area. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 1999 , 106, 415-20	3.7	14
356	The Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT): Study Design and Participants. <i>Journal of Epidemiology</i> , 2020 , 30, 46-54	3.4	14
355	Diabetes and cancer risk: A Mendelian randomization study. <i>International Journal of Cancer</i> , 2020 , 146, 712-719	7.5	14
354	Meat intake and risk of gastric cancer in the Stomach cancer Pooling (StoP) project. <i>International Journal of Cancer</i> , 2020 , 147, 45-55	7.5	14
353	Chocolate consumption and risk of stroke among men and women: A large population-based, prospective cohort study. <i>Atherosclerosis</i> , 2017 , 260, 8-12	3.1	13
352	Association of plasma concentrations of branched-chain amino acids with risk of colorectal adenoma in a large Japanese population. <i>Annals of Oncology</i> , 2017 , 28, 818-823	10.3	13
351	Dietary fiber intake and total and cause-specific mortality: the Japan Public Health Center-based prospective study. <i>American Journal of Clinical Nutrition</i> , 2020 , 111, 1027-1035	7	13
350	Dietary magnesium intake and risk of incident coronary heart disease in men: A prospective cohort study. <i>Clinical Nutrition</i> , 2018 , 37, 1602-1608	5.9	13
349	Plasma levels of n-3 fatty acids and risk of coronary heart disease among Japanese: The Japan Public Health Center-based (JPHC) study. <i>Atherosclerosis</i> , 2018 , 272, 226-232	3.1	13
348	Reproductive history and risk of cognitive impairment in Japanese women. <i>Maturitas</i> , 2019 , 128, 22-28	5	13
347	Validity of a self-administered food frequency questionnaire in the estimation of heterocyclic aromatic amines. <i>Cancer Causes and Control</i> , 2014 , 25, 1015-28	2.8	13
346	Plasma 25-hydroxy vitamin D and subsequent prostate cancer risk in a nested Case-Control study in Japan: The JPHC study. <i>European Journal of Clinical Nutrition</i> , 2017 , 71, 132-136	5.2	13
345	Estimation of lifetime cumulative incidence and mortality risk of gastric cancer. <i>Japanese Journal of Clinical Oncology</i> , 2017 , 47, 1097-1102	2.8	13
344	Plasma isoflavone concentrations are not associated with gastric cancer risk among Japanese men and women. <i>Journal of Nutrition</i> , 2013 , 143, 1293-8	4.1	13
343	Validity of a self-administered food-frequency questionnaire in the estimation of amino acid intake. <i>British Journal of Nutrition</i> , 2009 , 101, 1393-9	3.6	13

342	A case-control study of the association between urinary cadmium concentration and endometriosis in infertile Japanese women. <i>Science of the Total Environment</i> , 2008 , 402, 171-5	10.2	13
341	Maintenance of a low-sodium, high-carotene and -vitamin C diet after a 1-year dietary intervention: the Hiraka dietary intervention follow-up study. <i>Preventive Medicine</i> , 2006 , 43, 14-9	4.3	13
340	Quantitative analysis of 8-hydroxyguanine in peripheral blood cells: an application for asbestosis patients. <i>International Archives of Occupational and Environmental Health</i> , 1993 , 65, S215-7	3.2	13
339	Three novel NEIL1 promoter polymorphisms in gastric cancer patients. <i>World Journal of Gastrointestinal Oncology</i> , 2010 , 2, 117-20	3.4	13
338	Citrus fruit intake and gastric cancer: The stomach cancer pooling (StoP) project consortium. <i>International Journal of Cancer</i> , 2019 , 144, 2936-2944	7.5	13
337	Circulating isoflavone and lignan concentrations and prostate cancer risk: a meta-analysis of individual participant data from seven prospective studies including 2,828 cases and 5,593 controls. <i>International Journal of Cancer</i> , 2018 , 143, 2677-2686	7.5	13
336	Dietary fiber intake and risk of breast cancer defined by estrogen and progesterone receptor status: the Japan Public Health Center-based Prospective Study. <i>Cancer Causes and Control</i> , 2017 , 28, 569-578	2.8	12
335	Anthropometric Risk Factors for Cancers of the Biliary Tract in the Biliary Tract Cancers Pooling Project. <i>Cancer Research</i> , 2019 , 79, 3973-3982	10.1	12
334	Fruits and vegetables intake and gastric cancer risk: A pooled analysis within the Stomach cancer Pooling Project. <i>International Journal of Cancer</i> , 2020 , 147, 3090-3101	7.5	12
333	Alcohol consumption, genetic variants in the alcohol- and folate metabolic pathways and colorectal cancer risk: the JPHC Study. <i>Scientific Reports</i> , 2016 , 6, 36607	4.9	12
332	Marital Transition and Risk of Stroke: How Living Arrangement and Employment Status Modify Associations. <i>Stroke</i> , 2016 , 47, 991-8	6.7	12
331	Smoking cessation and subsequent risk of cancer: A pooled analysis of eight population-based cohort studies in Japan. <i>Cancer Epidemiology</i> , 2017 , 51, 98-108	2.8	12
330	The relationship between vegetable/fruit consumption and gallbladder/bile duct cancer: A population-based cohort study in Japan. <i>International Journal of Cancer</i> , 2017 , 140, 1009-1019	7.5	12
329	Neighborhood deprivation and risk of cancer incidence, mortality and survival: results from a population-based cohort study in Japan. <i>PLoS ONE</i> , 2014 , 9, e106729	3.7	12
328	Higher levels of antibodies to the tumour-associated antigen cyclin B1 in cancer-free individuals than in patients with breast cancer. <i>Clinical and Experimental Immunology</i> , 2014 , 178, 75-8	6.2	12
327	Plasma levels of adiponectin and primary liver cancer risk in middle-aged Japanese adults with hepatitis virus infection: a nested case-control study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2013 , 22, 2250-7	4	12
326	Education level and physical functional limitations among Japanese community residents-gender difference in prognosis from stroke. <i>BMC Public Health</i> , 2009 , 9, 131	4.1	12
325	Validity of a self-administered food frequency questionnaire (FFQ) and its generalizability to the estimation of dietary folate intake in Japan. <i>Nutrition Journal</i> , 2005 , 4, 26	4.3	12

324	Lifestyle and Health Related Factors Among Randomly Selected Japanese Residents in the City of Sao Paulo, Brazil, and their Comparisons with Japanese in Japan. <i>Journal of Epidemiology</i> , 1994 , 4, 37-46	3.4	12
323	Coping strategies and cancer incidence and mortality: The Japan Public Health Center-based prospective study. <i>Cancer Epidemiology</i> , 2016 , 40, 126-33	2.8	12
322	Body Mass Index and Risks of Incident Ischemic Stroke Subtypes: The Japan Public Health Center-Based Prospective (JPHC) Study. <i>Journal of Epidemiology</i> , 2019 , 29, 325-333	3.4	12
321	Association between adherence to the Japanese diet and all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. <i>European Journal of Nutrition</i> , 2021 , 60, 1327-1336	5.2	12
320	Plasma adiponectin levels, ADIPOQ variants, and incidence of type 2 diabetes: A nested case-control study. <i>Diabetes Research and Clinical Practice</i> , 2017 , 127, 254-264	7.4	11
319	Body mass index change during adulthood and risk of oesophageal squamous-cell carcinoma in a Japanese population: the Japan Public Health (JPHC)-based prospective study. <i>British Journal of Cancer</i> , 2017 , 117, 1715-1722	8.7	11
318	Coffee Drinking and Reduced Risk of Liver Cancer: Update on Epidemiological Findings and Potential Mechanisms. <i>Current Nutrition Reports</i> , 2019 , 8, 182-186	6	11
317	Dietary patterns and prostate cancer risk in Japanese: the Japan Public Health Center-based Prospective Study (JPHC Study). <i>Cancer Causes and Control</i> , 2018 , 29, 589-600	2.8	11
316	Dietary patterns and colorectal cancer risk in middle-aged adults: A large population-based prospective cohort study. <i>Clinical Nutrition</i> , 2018 , 37, 1019-1026	5.9	11
315	Green tea consumption and mortality in Japanese men and women: a pooled analysis of eight population-based cohort studies in Japan. <i>European Journal of Epidemiology</i> , 2019 , 34, 917-926	12.1	11
314	Association between parity and dentition status among Japanese women: Japan public health center-based oral health study. <i>BMC Public Health</i> , 2013 , 13, 993	4.1	11
313	Smoking and alcohol and subsequent risk of myelodysplastic syndromes in Japan: the Japan Public Health Centre-based Prospective Study. <i>British Journal of Haematology</i> , 2017 , 178, 747-755	4.5	11
312	Saturated fat intake and cardiovascular disease in Japanese population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015 , 22, 435-9	4	11
311	Association of postmenopausal endogenous sex hormones with global methylation level of leukocyte DNA among Japanese women. <i>BMC Cancer</i> , 2012 , 12, 323	4.8	11
310	The Impact of C-Reactive Protein on Risk of Stroke, Stroke Subtypes, and Ischemic Heart Disease in Middle-Aged Japanese: the Japan Public Health Center-Based Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012 , 19,	4	11
309	Racially restricted contribution of immunoglobulin Fc γ and Fc γ receptor genotypes to humoral immunity to human epidermal growth factor receptor 2 in breast cancer. <i>Clinical and Experimental Immunology</i> , 2013 , 171, 273-7	6.2	11
308	Interpopulation and intrapopulation variability of nutrient intake in five regions of Japan. <i>European Journal of Clinical Nutrition</i> , 1998 , 52, 176-9	5.2	11
307	Long-term study of urinary bisphenol A in elementary school children. <i>Environmental Health and Preventive Medicine</i> , 2008 , 13, 332-7	4.2	11

306	Alcohol drinking and total cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2007 , 37, 692-700	2.8	11
305	Breast cancer in first-degree relatives and risk of lung cancer: assessment of the existence of gene sex interactions. <i>Japanese Journal of Clinical Oncology</i> , 2007 , 37, 419-23	2.8	11
304	Comparison of selenium status between Japanese living in Tokyo and Japanese brazilians in Sã Paulo, Brazil. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2001 , 10, 197-9	1	11
303	Association of BMI and height with the risk of endometrial cancer, overall and by histological subtype: a population-based prospective cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2019 , 28, 196-202	2	11
302	Cruciferous vegetable intake and mortality in middle-aged adults: A prospective cohort study. <i>Clinical Nutrition</i> , 2019 , 38, 631-643	5.9	11
301	Non-High-Density Lipoprotein Cholesterol and Risk of Stroke Subtypes and Coronary Heart Disease: The Japan Public Health Center-Based Prospective (JPHC) Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 363-374	4	11
300	Epidemiological profiles of chronic low back and knee pain in middle-aged and elderly Japanese from the Murakami cohort. <i>Journal of Pain Research</i> , 2018 , 11, 3161-3169	2.9	11
299	Daily Total Physical Activity and Incident Cardiovascular Disease in Japanese Men and Women: Japan Public Health Center-Based Prospective Study. <i>Circulation</i> , 2017 , 135, 1471-1473	16.7	10
298	Smoking and subsequent risk of leukemia in Japan: The Japan Public Health Center-based Prospective Study. <i>Journal of Epidemiology</i> , 2017 , 27, 305-310	3.4	10
297	Neighborhood contextual factors for smoking among middle-aged Japanese: a multilevel analysis. <i>Health and Place</i> , 2015 , 31, 17-23	4.6	10
296	Development of a risk prediction model for lung cancer: The Japan Public Health Center-based Prospective Study. <i>Cancer Science</i> , 2018 , 109, 854-862	6.9	10
295	Coffee and green tea consumption and subsequent risk of acute myeloid leukemia and myelodysplastic syndromes in Japan. <i>International Journal of Cancer</i> , 2018 , 142, 1130-1138	7.5	10
294	Educational Levels and Risk of Suicide in Japan: The Japan Public Health Center Study (JPHC) Cohort I. <i>Journal of Epidemiology</i> , 2016 , 26, 315-21	3.4	10
293	Glycemic index and glycemic load and risk of colorectal cancer: a population-based cohort study (JPHC Study). <i>Cancer Causes and Control</i> , 2016 , 27, 583-93	2.8	10
292	Alcohol consumption and bladder cancer risk with or without the flushing response: The Japan Public Health Center-based Prospective Study. <i>International Journal of Cancer</i> , 2017 , 141, 2480-2488	7.5	10
291	Risk of lung cancer and consumption of vegetables and fruit in Japanese: A pooled analysis of cohort studies in Japan. <i>Cancer Science</i> , 2015 , 106, 1057-65	6.9	10
290	Association of CYP19A1 polymorphisms with risks for atypical adenomatous hyperplasia and bronchioloalveolar carcinoma in the lungs. <i>Carcinogenesis</i> , 2010 , 31, 1794-9	4.6	10
289	Dietary differences with green tea intake among middle-aged Japanese men and women. <i>Preventive Medicine</i> , 1997 , 26, 704-10	4.3	10

288	Adult height and all-cause and cause-specific mortality in the Japan Public Health Center-based Prospective Study (JPHC). <i>PLoS ONE</i> , 2018 , 13, e0197164	3.7	10
287	Validation of a Blood Biomarker for Identification of Individuals at High Risk for Gastric Cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2018 , 27, 1472-1479	4	10
286	Diet quality and depression risk in a Japanese population: the Japan Public Health Center (JPHC)-based Prospective Study. <i>Scientific Reports</i> , 2019 , 9, 7150	4.9	9
285	Expectations of and recommendations for a cancer survivorship guideline in Japan: a literature review of guidelines for cancer survivorship. <i>Japanese Journal of Clinical Oncology</i> , 2019 , 49, 812-822	2.8	9
284	Transethnic Meta-Analysis of Genome-Wide Association Studies Identifies Three New Loci and Characterizes Population-Specific Differences for Coronary Artery Disease. <i>Circulation Genomic and Precision Medicine</i> , 2020 , 13, e002670	5.2	9
283	Habitual tub bathing and risks of incident coronary heart disease and stroke. <i>Heart</i> , 2020 , 106, 732-737	5.1	9
282	Correlation between global methylation level of peripheral blood leukocytes and serum C reactive protein level modified by MTHFR polymorphism: a cross-sectional study. <i>BMC Cancer</i> , 2018 , 18, 184	4.8	9
281	Female reproductive factors and risk of all-cause and cause-specific mortality among women: The Japan Public Health Center-based Prospective Study (JPHC study). <i>Annals of Epidemiology</i> , 2018 , 28, 597-604.e6	6.4	9
280	Menstrual and reproductive factors and type 2 diabetes risk: The Japan Public Health Center-based Prospective Study. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 147-153	3.9	9
279	Descriptive epidemiology of diabetes prevalence and HbA1c distributions based on a self-reported questionnaire and a health checkup in the JPHC diabetes study. <i>Journal of Epidemiology</i> , 2014 , 24, 460-8 ³⁻⁴	3.4	9
278	The association between complete and partial non-response to psychosocial questions and suicide: the JPHC Study. <i>European Journal of Public Health</i> , 2015 , 25, 424-30	2.1	9
277	Use of vitamin supplements and risk of total cancer and cardiovascular disease among the Japanese general population: a population-based survey. <i>BMC Public Health</i> , 2011 , 11, 540	4.1	9
276	Validity of self-reported cancer among a Japanese population: recent results from a population-based prospective study in Japan (JPHC Study). <i>Cancer Epidemiology</i> , 2011 , 35, 250-3	2.8	9
275	Within- and between-person variations in portion sizes of foods consumed by the Japanese population. <i>Nutrition and Cancer</i> , 1997 , 29, 140-5	2.8	9
274	Modelling of mortality data from a multi-centre study in Japan by means of Poisson regression with error in variables. <i>International Journal of Epidemiology</i> , 1997 , 26, 501-7	7.8	9
273	A pilot study for a randomized controlled trial to prevent gastric cancer in high-risk Japanese population: study design and feasibility evaluation. <i>Japanese Journal of Cancer Research</i> , 1996 , 87, 676-9		9
272	Cigarette smoking, alcohol drinking, and oral cavity and pharyngeal cancer in the Japanese: a population-based cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2018 , 27, 171-179	2	9
271	Prediagnostic circulating inflammation biomarkers and esophageal squamous cell carcinoma: A case-cohort study in Japan. <i>International Journal of Cancer</i> , 2020 , 147, 686-691	7.5	9

270	Exploring the interactions between <i>Helicobacter pylori</i> (Hp) infection and other risk factors of gastric cancer: A pooled analysis in the Stomach cancer Pooling (StoP) Project. <i>International Journal of Cancer</i> , 2021 , 149, 1228-1238	7.5	9
269	Sex differences in the prevalence of <i>Helicobacter pylori</i> infection: an individual participant data pooled analysis (StoP Project). <i>European Journal of Gastroenterology and Hepatology</i> , 2019 , 31, 593-598	2.2	9
268	Association of dietary diversity with total mortality and major causes of mortality in the Japanese population: JPHC study. <i>European Journal of Clinical Nutrition</i> , 2020 , 74, 54-66	5.2	9
267	Reproductive factors and gallbladder/bile duct cancer: a population-based cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2017 , 26, 292-300	2	8
266	Smoking and Pancreatic Cancer Incidence: A Pooled Analysis of 10 Population-Based Cohort Studies in Japan. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2019 , 28, 1370-1378	4	8
265	Genome-wide association meta-analysis and Mendelian randomization analysis confirm the influence of ALDH2 on sleep duration in the Japanese population. <i>Sleep</i> , 2019 , 42,	1.1	8
264	Coffee consumption and mortality in Japanese men and women: A pooled analysis of eight population-based cohort studies in Japan (Japan Cohort Consortium). <i>Preventive Medicine</i> , 2019 , 123, 270-277	4.3	8
263	Association between plasma concentrations of branched-chain amino acids and adipokines in Japanese adults without diabetes. <i>Scientific Reports</i> , 2018 , 8, 1043	4.9	8
262	Coffee, green tea and liver cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. <i>Japanese Journal of Clinical Oncology</i> , 2019 , 49, 972-984	2.8	8
261	Positive predictive values for self-reported fractures in an adult Japanese population. <i>Environmental Health and Preventive Medicine</i> , 2011 , 16, 129-32	4.2	8
260	Effect of five-year supplementation of vitamin C on serum vitamin C concentration and consumption of vegetables and fruits in middle-aged Japanese: a randomized controlled trial. <i>Journal of the American College of Nutrition</i> , 2003 , 22, 208-16	3.5	8
259	The mercury content of hair of Japanese immigrants in various locations in South America. <i>Science of the Total Environment</i> , 1987 , 63, 69-76	10.2	8
258	Chromosome fragility of lymphocytes from breast cancer patients in relation to epidemiologic data. <i>Japanese Journal of Cancer Research</i> , 1988 , 79, 1024-30		8
257	High-Negative Anti- IgG Antibody Titers and Long-Term Risk of Gastric Cancer: Results from a Large-Scale Population-Based Cohort Study in Japan. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020 , 29, 420-426	4	8
256	Trends in the proportions of stroke subtypes and coronary heart disease in the Japanese men and women from 1995 to 2009. <i>Atherosclerosis</i> , 2016 , 248, 219-23	3.1	8
255	Tuberculosis infection and lung adenocarcinoma: Mendelian randomization and pathway analysis of genome-wide association study data from never-smoking Asian women. <i>Genomics</i> , 2020 , 112, 1223-1232	4.3	8
254	Ethnic differences in serum pepsinogen levels among Japanese and non-Japanese Brazilian gastric cancer patients and controls. <i>Cancer Detection and Prevention</i> , 2000 , 24, 564-71		8
253	Food frequency questionnaire is a valid tool in the nutritional assessment of Brazilian women of diverse ethnicity. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2009 , 18, 76-80	1	8

252	Prediagnostic Calcium Intake and Lung Cancer Survival: A Pooled Analysis of 12 Cohort Studies. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2017 , 26, 1060-1070	4	7
251	Effect of body-mass index on the risk of gastric cancer: A population-based cohort study in A Japanese population. <i>Cancer Epidemiology</i> , 2019 , 63, 101622	2.8	7
250	Circulating Inflammation Markers and Risk of Gastric and Esophageal Cancers: A Case-Cohort Study Within the Japan Public Health Center-Based Prospective Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2019 , 28, 829-832	4	7
249	Occupational sitting time and subsequent risk of cancer: The Japan Public Health Center-based Prospective Study. <i>Cancer Science</i> , 2020 , 111, 974-984	6.9	7
248	Variations in the estimated intake of acrylamide from food in the Japanese population. <i>Nutrition Journal</i> , 2020 , 19, 17	4.3	7
247	Risk of thyroid cancer in relation to height, weight, and body mass index in Japanese individuals: a population-based cohort study. <i>Cancer Medicine</i> , 2018 , 7, 2200-2210	4.8	7
246	Predictive performance of a genetic risk score using 11 susceptibility alleles for the incidence of Type 2 diabetes in a general Japanese population: a nested case-control study. <i>Diabetic Medicine</i> , 2018 , 35, 602-611	3.5	7
245	Metabolome analysis for pancreatic cancer risk in nested case-control study: Japan Public Health Center-based prospective Study. <i>Cancer Science</i> , 2018 , 109, 1672-1681	6.9	7
244	Smoking and subsequent risk of acute myeloid leukaemia: A pooled analysis of 9 cohort studies in Japan. <i>Hematological Oncology</i> , 2018 , 36, 262-268	1.3	7
243	Report of the Japan Diabetes Society (JDS)/Japanese Cancer Association (JCA) Joint Committee on Diabetes and Cancer, Second Report. <i>Diabetology International</i> , 2016 , 7, 12-15	2.3	7
242	History of diabetes and risk of suicide and accidental death in Japan: The Japan Public Health Centre-based Prospective Study, 1990-2012. <i>Diabetes and Metabolism</i> , 2016 , 42, 184-91	5.4	7
241	Epstein-Barr Virus Antibody Titers Are Not Associated with Gastric Cancer Risk in East Asia. <i>Digestive Diseases and Sciences</i> , 2018 , 63, 2765-2772	4	7
240	Smoking, Alcohol Consumption, and Risks for Biliary Tract Cancer and Intrahepatic Bile Duct Cancer. <i>Journal of Epidemiology</i> , 2019 , 29, 180-186	3.4	7
239	Trends in cancer prognosis in a population-based cohort survey: can recent advances in cancer therapy affect the prognosis?. <i>Cancer Epidemiology</i> , 2015 , 39, 97-103	2.8	7
238	Serum brain-derived neurotrophic factor and antidepressant-naive major depression after lung cancer diagnosis. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 1233-7	2.8	7
237	Projected cancer mortality among Japanese males under different smoking prevalence scenarios: evidence for tobacco control goal setting. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 483-9	2.8	7
236	Projecting the probability of survival free from cancer and cardiovascular incidence through lifestyle modification in Japan. <i>Preventive Medicine</i> , 2009 , 48, 128-33	4.3	7
235	The association between cancer risk and age at onset of smoking in Japanese. <i>Journal of Epidemiology</i> , 2010 , 20, 128-35	3.4	7

234	Differing effects of vasopressin on regional cerebral blood flow of dogs following intracisternal vs. intra-arterial administration. <i>Life Sciences</i> , 1994 , 54, PL241-6	6.8	7
233	Cervico-thoraco-abdominal (3-field) lymph node dissection for carcinoma in the thoracic esophagus. <i>Kurume Medical Journal</i> , 1992 , 39, 167-74	0.9	7
232	Fermented soy products intake and risk of cardiovascular disease and total cancer incidence: The Japan Public Health Center-based Prospective study. <i>European Journal of Clinical Nutrition</i> , 2021 , 75, 954-968	5.2	7
231	Smoking and colorectal cancer: A pooled analysis of 10 population-based cohort studies in Japan. <i>International Journal of Cancer</i> , 2021 , 148, 654-664	7.5	7
230	Coffee Consumption and Lung Cancer Risk: The Japan Public Health Center-Based Prospective Study. <i>Journal of Epidemiology</i> , 2018 , 28, 207-213	3.4	7
229	Coffee and tea consumption and mortality from all causes, cardiovascular disease and cancer: a pooled analysis of prospective studies from the Asia Cohort Consortium. <i>International Journal of Epidemiology</i> , 2021 ,	7.8	7
228	The association of active and secondhand smoking with oral health in adults: Japan public health center-based study. <i>Tobacco Induced Diseases</i> , 2015 , 13, 19	3.2	6
227	Identification of a novel uterine leiomyoma GWAS locus in a Japanese population. <i>Scientific Reports</i> , 2020 , 10, 1197	4.9	6
226	Weight loss from 20 years of age is associated with cognitive impairment in middle-aged and elderly individuals. <i>PLoS ONE</i> , 2017 , 12, e0185960	3.7	6
225	Menstrual and reproductive factors in the risk of thyroid cancer in Japanese women: the Japan Public Health Center-Based Prospective Study. <i>European Journal of Cancer Prevention</i> , 2018 , 27, 361-369 ²		6
224	Humoral Immunity to Cytomegalovirus Glycoprotein B in Patients With Breast Cancer and Matched Controls: Contribution of Immunoglobulin μ and Fc γ Receptor Genes. <i>Journal of Infectious Diseases</i> , 2016 , 213, 611-7	7	6
223	Revisit of an unanswered question by pooled analysis of eight cohort studies in Japan: Does cigarette smoking and alcohol drinking have interaction for the risk of esophageal cancer?. <i>Cancer Medicine</i> , 2019 , 8, 6414-6425	4.8	6
222	Impact of five modifiable lifestyle habits on the probability of cancer occurrence in a Japanese population-based cohort: results from the JPHC study. <i>Preventive Medicine</i> , 2013 , 57, 685-9	4.3	6
221	A novel method to quantify base substitution mutations at the 10 per bp level in DNA samples. <i>Cancer Letters</i> , 2017 , 403, 152-158	9.9	6
220	Rice, bread, noodle and cereal intake and colorectal cancer in Japanese men and women: the Japan Public Health Center-based prospective Study (JPHC Study). <i>British Journal of Cancer</i> , 2014 , 110, 1316-21	8.7	6
219	Validation of a quantitative FFQ for a study of diet and risk of colorectal adenoma among Japanese Brazilians. <i>Public Health Nutrition</i> , 2013 , 16, 1445-53	3.3	6
218	Effectiveness of the combined approach for assessing social gradients in stroke risk among married women in Japan. <i>Journal of Epidemiology</i> , 2012 , 22, 324-30	3.4	6
217	Four food-frequency categories of fruit intake as a predictor of plasma ascorbic acid level in middle-aged Japanese men. <i>Annals of Epidemiology</i> , 1998 , 8, 378-83	6.4	6

216	Development and validation of specific carotene food composition tables for use in nutritional epidemiologic studies for Japanese populations. <i>Journal of Epidemiology</i> , 2001 , 11, 266-75	3.4	6
215	Does high gastric cancer risk associated with low serum ferritin level reflect achlorhydria? An examination via cross-sectional study. <i>Japanese Journal of Cancer Research</i> , 1993 , 84, 844-51		6
214	Association between meat intake and mortality due to all-cause and major causes of death in a Japanese population. <i>PLoS ONE</i> , 2020 , 15, e0244007	3.7	6
213	The U-shaped association between body mass index and gastric cancer risk in the Helicobacter pylori Biomarker Cohort Consortium: A nested case-control study from eight East Asian cohort studies. <i>International Journal of Cancer</i> , 2020 , 147, 777-784	7.5	6
212	Doneness preferences, meat and meat-derived heterocyclic amines intake, and N-acetyltransferase 2 polymorphisms: association with colorectal adenoma in Japanese Brazilians. <i>European Journal of Cancer Prevention</i> , 2020 , 29, 7-14	2	6
211	Relationship between unhealthy sleep status and dry eye symptoms in a Japanese population: The JPHC-NEXT study. <i>Ocular Surface</i> , 2021 , 21, 306-312	6.5	6
210	Report of the Japan diabetes society/Japanese cancer association joint committee on diabetes and cancer, Second report. <i>Cancer Science</i> , 2016 , 107, 369-71	6.9	6
209	Association between serum liver enzymes and all-cause mortality: The Japan Public Health Center-based Prospective Study. <i>Liver International</i> , 2019 , 39, 1566-1576	7.9	6
208	Association of Sleep Duration With All- and Major-Cause Mortality Among Adults in Japan, China, Singapore, and Korea. <i>JAMA Network Open</i> , 2021 , 4, e2122837	10.4	6
207	Meat subtypes and colorectal cancer risk: A pooled analysis of 6 cohort studies in Japan. <i>Cancer Science</i> , 2019 , 110, 3603-3614	6.9	5
206	Relationships of diabetes and hyperglycaemia with intraocular pressure in a Japanese population: the JPHC-NEXT Eye Study. <i>Scientific Reports</i> , 2020 , 10, 5355	4.9	5
205	Time Spent Walking and Risk of Diabetes in Japanese Adults: The Japan Public Health Center-Based Prospective Diabetes Study. <i>Journal of Epidemiology</i> , 2016 , 26, 224-32	3.4	5
204	Smoking is a risk factor for development of adult T-cell leukemia/lymphoma in Japanese human T-cell leukemia virus type-1 carriers. <i>Cancer Causes and Control</i> , 2016 , 27, 1059-66	2.8	5
203	Changes in Smoking Status and Mortality From All Causes and Lung Cancer: A Longitudinal Analysis of a Population-based Study in Japan. <i>Journal of Epidemiology</i> , 2019 , 29, 11-17	3.4	5
202	Fish intake and risk of mortality due to aortic dissection and aneurysm: A pooled analysis of the Japan cohort consortium. <i>Clinical Nutrition</i> , 2019 , 38, 1678-1683	5.9	5
201	Online version of the self-administered food frequency questionnaire for the Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT) protocol: Relative validity, usability, and comparison with a printed questionnaire. <i>Journal of Epidemiology</i> , 2017 , 27, 435-446	3.4	5
200	Association of plasma C-reactive protein level with the prevalence of colorectal adenoma: the Colorectal Adenoma Study in Tokyo. <i>Scientific Reports</i> , 2017 , 7, 4456	4.9	5
199	New breast cancer risk variant discovered at 10q25 in East Asian women. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2013 , 22, 1297-303	4	5

198	Association between polymorphisms in glutathione S-transferase Mu3 and IgG titer levels in serum against Helicobacter pylori. <i>Journal of Human Genetics</i> , 2009 , 54, 557-63	4.3	5
197	Alcohol consumption, alcohol-induced flushing and incidence of acute myocardial infarction among middle-aged men in Japan--Japan Public Health Center-based prospective study. <i>Atherosclerosis</i> , 2007 , 194, 512-6	3.1	5
196	RESPONSE: Re: Soy, Isoflavones, and Breast Cancer Risk in Japan. <i>Journal of the National Cancer Institute</i> , 2003 , 95, 1881-1882	9.7	5
195	Alternating sequential chemotherapy with high-dose ifosfamide and doxorubicin/cyclophosphamide for adult non-small round cell soft tissue sarcomas. <i>Journal of Orthopaedic Science</i> , 2005 , 10, 258-63	1.6	5
194	Effects of human diets of two different Japanese populations on cancer incidence in rat hepatic drug-metabolizing and antioxidant enzyme systems. <i>Nutrition and Cancer</i> , 1996 , 26, 63-71	2.8	5
193	Anomalous Magnetoresistance in Amorphous Si?Au System. <i>Physica Status Solidi (B): Basic Research</i> , 1981 , 105, K5-K8	1.3	5
192	Fermented and nonfermented soy foods and the risk of breast cancer in a Japanese population-based cohort study. <i>Cancer Medicine</i> , 2021 , 10, 757-771	4.8	5
191	Dietary Acrylamide Intake and the Risk of Pancreatic Cancer: The Japan Public Health Center-Based Prospective Study. <i>Nutrients</i> , 2020 , 12,	6.7	5
190	Dietary Inflammatory Index Is Associated With Inflammation in Japanese Men. <i>Frontiers in Nutrition</i> , 2021 , 8, 604296	6.2	5
189	Identification of two novel breast cancer loci through large-scale genome-wide association study in the Japanese population. <i>Scientific Reports</i> , 2019 , 9, 17332	4.9	5
188	Study Design and Baseline Profiles of Participants in the Uonuma CKD Cohort Study in Niigata, Japan. <i>Journal of Epidemiology</i> , 2020 , 30, 170-176	3.4	5
187	Low carbohydrate diet and all cause and cause-specific mortality. <i>Clinical Nutrition</i> , 2021 , 40, 2016-2024	5.9	5
186	Associations of coffee and tea consumption with lung cancer risk. <i>International Journal of Cancer</i> , 2020 , 148, 2457	7.5	5
185	Body size and weight change over adulthood and risk of breast cancer by menopausal and hormone receptor status: a pooled analysis of 20 prospective cohort studies. <i>European Journal of Epidemiology</i> , 2021 , 36, 37-55	12.1	5
184	Body mass index and colorectal cancer risk: A Mendelian randomization study. <i>Cancer Science</i> , 2021 , 112, 1579-1588	6.9	5
183	Genome-wide association study (GWAS) of ovarian cancer in Japanese predicted regulatory variants in 22q13.1. <i>PLoS ONE</i> , 2018 , 13, e0209096	3.7	5
182	Comparison of weighed food record procedures for the reference methods in two validation studies of food frequency questionnaires. <i>Journal of Epidemiology</i> , 2017 , 27, 331-337	3.4	4
181	Association of BMI, Smoking, and Alcohol with Multiple Myeloma Mortality in Asians: A Pooled Analysis of More than 800,000 Participants in the Asia Cohort Consortium. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2019 , 28, 1861-1867	4	4

180	Validity and Reproducibility of a Self-Administered Food Frequency Questionnaire for the Assessment of Sugar Intake in Middle-Aged Japanese Adults. <i>Nutrients</i> , 2019 , 11,	6.7	4
179	Relationship between dietary non-enzymatic antioxidant capacity and type 2 diabetes risk in the Japan Public Health Center-based Prospective Study. <i>Nutrition</i> , 2019 , 66, 62-69	4.8	4
178	Female reproductive factors and risk of lymphoid neoplasm: The Japan Public Health Center-based Prospective Study. <i>Cancer Science</i> , 2019 , 110, 1442-1452	6.9	4
177	Fat mass and obesity-associated gene polymorphisms, pre-diagnostic plasma adipokine levels and the risk of colorectal cancer: The Japan Public Health Center-based Prospective Study. <i>PLoS ONE</i> , 2020 , 15, e0229005	3.7	4
176	Total physical activity and risk of chronic low back and knee pain in middle-aged and elderly Japanese people: The Murakami cohort study. <i>European Journal of Pain</i> , 2020 , 24, 863-872	3.7	4
175	The Validity and Reproducibility of Dietary Non-enzymatic Antioxidant Capacity Estimated by Self-administered Food Frequency Questionnaires. <i>Journal of Epidemiology</i> , 2018 , 28, 428-436	3.4	4
174	Menstrual and reproductive factors and risk of vertebral fractures in Japanese women: the Japan Public Health Center-based prospective (JPHC) study. <i>Osteoporosis International</i> , 2018 , 29, 2791-2801	5.3	4
173	Physical activity and 10-year incidence of self-reported vertebral fractures in Japanese women: the Japan Public Health Center-based Prospective Study. <i>Osteoporosis International</i> , 2014 , 25, 2565-71	5.3	4
172	Subcutaneous hematoma associated with manual cervical massage during carotid artery stenting. A case report. <i>Interventional Neuroradiology</i> , 2011 , 17, 386-90	1.9	4
171	Comparison of postmenopausal endogenous sex hormones among Japanese, Japanese Brazilians, and non-Japanese Brazilians. <i>BMC Medicine</i> , 2011 , 9, 16	11.4	4
170	Perceptions of change in diet have limited utility for improving estimates of past food frequency of individuals. <i>Nutrition and Cancer</i> , 1995 , 23, 299-307	2.8	4
169	Dietary Acrylamide Intake and Risk of Lung Cancer: The Japan Public Health Center Based Prospective Study. <i>Nutrients</i> , 2020 , 12,	6.7	4
168	Dietary Acrylamide Intake and the Risk of Liver Cancer: The Japan Public Health Center-Based Prospective Study. <i>Nutrients</i> , 2020 , 12,	6.7	4
167	Weight change during middle age and risk of stroke and coronary heart disease: The Japan Public Health Center-based Prospective Study. <i>Atherosclerosis</i> , 2021 , 322, 67-73	3.1	4
166	Dairy foods, calcium, and risk of breast cancer overall and for subtypes defined by estrogen receptor status: a pooled analysis of 21 cohort studies. <i>American Journal of Clinical Nutrition</i> , 2021 , 114, 450-461	7	4
165	Association of Choroidal Thickness with Intermediate Age-Related Macular Degeneration in a Japanese Population. <i>Ophthalmology Retina</i> , 2021 , 5, 528-535	3.8	4
164	Association between educational level and total and cause-specific mortality: a pooled analysis of over 694 000 individuals in the Asia Cohort Consortium. <i>BMJ Open</i> , 2019 , 9, e026225	3	4
163	Fruit and vegetable intake and pancreatic cancer risk in a population-based cohort study in Japan. <i>International Journal of Cancer</i> , 2019 , 144, 1858-1866	7.5	4

162	Association of Vegetable, Fruit, and Okinawan Vegetable Consumption With Incident Stroke and Coronary Heart Disease. <i>Journal of Epidemiology</i> , 2020 , 30, 37-45	3.4	4
161	Soy food and isoflavones are not associated with changes in serum lipids and glycohemoglobin concentrations among Japanese adults: a cohort study. <i>European Journal of Nutrition</i> , 2020 , 59, 2075-2087	5.2	4
160	Comparison between the impact of fermented and unfermented soy intake on the risk of liver cancer: the JPHC Study. <i>European Journal of Nutrition</i> , 2021 , 60, 1389-1401	5.2	4
159	Dietary Acrylamide Intake and the Risk of Hematological Malignancies: The Japan Public Health Center-Based Prospective Study. <i>Nutrients</i> , 2021 , 13,	6.7	4
158	Association between meat and saturated fatty acid intake and lung cancer risk: The Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2020 , 147, 3019-3028	7.5	4
157	Fruit and vegetable intake and the risk of overall cancer in Japanese: A pooled analysis of population-based cohort studies. <i>Journal of Epidemiology</i> , 2017 , 27, 152-162	3.4	3
156	Higher Dietary Non-enzymatic Antioxidant Capacity Is Associated with Decreased Risk of All-Cause and Cardiovascular Disease Mortality in Japanese Adults. <i>Journal of Nutrition</i> , 2019 ,	4.1	3
155	Diet Quality Affects the Association between Census-Based Neighborhood Deprivation and All-Cause Mortality in Japanese Men and Women: The Japan Public Health Center-Based Prospective Study. <i>Nutrients</i> , 2019 , 11,	6.7	3
154	Midlife cancer/diabetes and risk of dementia and mild cognitive impairment: A population-based prospective cohort study in Japan. <i>Psychiatry and Clinical Neurosciences</i> , 2019 , 73, 597-599	6.2	3
153	Association of estimated dietary acid load with albuminuria in Japanese adults: a cross-sectional study. <i>BMC Nephrology</i> , 2019 , 20, 194	2.7	3
152	The functional ALDH2 polymorphism is associated with breast cancer risk: A pooled analysis from the Breast Cancer Association Consortium. <i>Molecular Genetics & Genomic Medicine</i> , 2019 , 7, e707	2.3	3
151	Metabolic Syndrome, Physical Activity, and Inflammation: A Cross-Sectional Analysis of 110 Circulating Biomarkers in Japanese Adults. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020 , 29, 1639-1646	4	3
150	Predictors of decline in vitamin D status in middle-aged and elderly individuals: a 5-year follow-up study. <i>British Journal of Nutrition</i> , 2020 , 124, 729-735	3.6	3
149	Consumption of flavonoid-rich fruits and risk of CHD: a prospective cohort study. <i>British Journal of Nutrition</i> , 2020 , 124, 952-959	3.6	3
148	Soy Food Intake and Pancreatic Cancer Risk: The Japan Public Health Center-based Prospective Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020 , 29, 1214-1221	4	3
147	Estimation of lifetime cumulative mortality risk of lung cancer by smoking status in Japan. <i>Japanese Journal of Clinical Oncology</i> , 2020 , 50, 1218-1224	2.8	3
146	Modifiable factors associated with symptomatic knee osteoarthritis: The Murakami cohort study. <i>Maturitas</i> , 2019 , 128, 53-59	5	3
145	Coffee and Green Tea Consumption and Subsequent Risk of Malignant Lymphoma and Multiple Myeloma in Japan: The Japan Public Health Center-based Prospective Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2017 , 26, 1352-1356	4	3

144	Female reproductive factors, exogenous hormone use, and pancreatic cancer risk: the Japan Public Health Center-based prospective study. <i>European Journal of Cancer Prevention</i> , 2017 , 26, 378-384	2	3
143	Further study of translational research for preventive medicine. <i>Preventive Medicine</i> , 2012 , 55, 573-4	4.3	3
142	Vitamin C supplementation in relation to inflammation in individuals with atrophic gastritis: a randomised controlled trial in Japan. <i>British Journal of Nutrition</i> , 2013 , 109, 1089-95	3.6	3
141	Diabetes mellitus defined by hemoglobin A1c value: Risk characterization for incidence among Japanese subjects in the JPHC Diabetes Study. <i>Journal of Diabetes Investigation</i> , 2011 , 2, 359-65	3.9	3
140	Effect of cooking loss in the assessment of vitamin intake for epidemiological data in Japan. <i>European Journal of Clinical Nutrition</i> , 2011 , 65, 546-52	5.2	3
139	Reply to F Imamura. <i>American Journal of Clinical Nutrition</i> , 2010 , 91, 1805-1806	7	3
138	Nutritional composition of dishes commonly consumed by Japanese Brazilians in S ^o Paulo, Brazil. <i>International Journal of Food Sciences and Nutrition</i> , 2010 , 61, 549-72	3.7	3
137	Drinking habit as a base for blood pressure elevation--difference in epidemiological significance by beverage type. <i>Applied Human Science: Journal of Physiological Anthropology</i> , 1997 , 16, 47-53		3
136	Soy Isoflavone Consumption Is Not Associated with Increased Risk of Advanced Prostate Cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2007 , 16, 2169.2-2169	4	3
135	Importance of standardization of hemoglobin A1c in the analysis of factors that predict hemoglobin A1c levels in non-diabetic residents of three distinct areas of Japan. <i>Diabetes Research and Clinical Practice</i> , 2001 , 53, 91-7	7.4	3
134	Proportional cancer incidence according to selected sites--comparison between residents in the city of S. Paulo, Brazil: Japanese and Brazilian/Portuguese descent. <i>Revista De Saude Publica</i> , 1991 , 25, 188-92	2.4	3
133	Low Carbohydrate Diet and Risk of Cancer Incidence: the Japan Public Health Center-based Prospective Study. <i>Cancer Science</i> , 2021 ,	6.9	3
132	Usefulness of the Recall-Based Self-Report as a Means of Case Definition in Epidemiological Studies on Senile Cataracts.. [<i>Minzoku Eisei</i>] <i>Race Hygiene</i> , 2002 , 68, 43-53		3
131	Food frequency questionnaire reproducibility for middle-aged and elderly Japanese. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2019 , 28, 362-370	1	3
130	Suction decompression methods for giant internal carotid ophthalmic aneurysms by using revised double lumen balloon catheters. <i>Acta Neurochirurgica Supplementum</i> , 2008 , 103, 9-10	1.7	3
129	Soy and isoflavone consumption and subsequent risk of prostate cancer mortality: the Japan Public Health Center-based Prospective Study. <i>International Journal of Epidemiology</i> , 2020 , 49, 1553-1561	7.8	3
128	Intensity-specific validity and reliability of the Japan Public Health Center-based prospective study-physical activity questionnaire. <i>Preventive Medicine Reports</i> , 2020 , 20, 101169	2.6	3
127	Association between dietary sugar intake and colorectal adenoma among cancer screening examinees in Japan. <i>Cancer Science</i> , 2020 , 111, 3862-3872	6.9	3

126	Validity of Self-Reported Periodontitis in Japanese Adults: The Japan Public Health Center-Based Prospective Study for the Next-Generation Oral Health Study. <i>Asia-Pacific Journal of Public Health</i> , 2020 , 32, 346-353	2	3
125	Inclusion of a gene-environment interaction between alcohol consumption and the aldehyde dehydrogenase 2 genotype in a risk prediction model for upper aerodigestive tract cancer in Japanese men. <i>Cancer Science</i> , 2020 , 111, 3835-3844	6.9	3
124	High Myopia and Its Associated Factors in JPHC-NEXT Eye Study: A Cross-Sectional Observational Study. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	3
123	Smoking and Helicobacter pylori infection: an individual participant pooled analysis (Stomach Cancer Pooling- StoP Project). <i>European Journal of Cancer Prevention</i> , 2019 , 28, 390-396	2	3
122	Associations between changes in fruit and vegetable consumption and weight change in Japanese adults. <i>European Journal of Nutrition</i> , 2021 , 60, 217-227	5.2	3
121	The Association Between Habitual Sleep Duration and Mortality According to Sex and Age: The Japan Public Health Center-based Prospective Study. <i>Journal of Epidemiology</i> , 2021 , 31, 109-118	3.4	3
120	Consumption of flavonoid-rich fruits, flavonoids from fruits and stroke risk: a prospective cohort study. <i>British Journal of Nutrition</i> , 2021 , 126, 1717-1724	3.6	3
119	Sugary Drink Consumption and Subsequent Colorectal Cancer Risk: The Japan Public Health Center-Based Prospective Cohort Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021 , 30, 782-788	4.1	3
118	Dietary Acrylamide Intake and the Risks of Renal Cell, Prostate, and Bladder Cancers: A Japan Public Health Center-Based Prospective Study. <i>Nutrients</i> , 2021 , 13,	6.7	3
117	Alcohol consumption, tobacco smoking, and subsequent risk of renal cell carcinoma: The JPHC study. <i>Cancer Science</i> , 2021 , 112, 5068-5077	6.9	3
116	Effect of smoking on the association between alcohol consumption and cancer mortality among middle-aged Japanese men: JPHC Study Cohort I. <i>Iarc (international Agency for Research on Cancer) Scientific Publications</i> , 2002 , 156, 165-8		3
115	Validity and reliability of a self-administered food frequency questionnaire for the JPHC study: The assessment of amino acid intake. <i>Journal of Epidemiology</i> , 2017 , 27, 242-247	3.4	2
114	Two truncating variants in FANCC and breast cancer risk. <i>Scientific Reports</i> , 2019 , 9, 12524	4.9	2
113	Reactive oxygen species and gastric cancer risk: a large nested case-control study in Japan. <i>European Journal of Epidemiology</i> , 2015 , 30, 589-94	12.1	2
112	Comparison of plasma levels of nutrient-related biomarkers among Japanese populations in Tokyo, Japan, S Paulo, Brazil, and Hawaii, USA. <i>European Journal of Cancer Prevention</i> , 2015 , 24, 155-61	2	2
111	Passive smoking and type2 diabetes among never-smoking women: The Japan Public Health Center-based Prospective Study. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1352-1358	3.9	2
110	Changes in the living arrangement and risk of stroke in Japan; does it matter who lives in the household? Who among the family matters?. <i>PLoS ONE</i> , 2017 , 12, e0173860	3.7	2
109	High serum total cholesterol is associated with suicide mortality in Japanese women independently of menopause. <i>Acta Psychiatrica Scandinavica</i> , 2018 , 137, 80-81	6.5	2

108	History of having a macrosomic infant and the risk of diabetes: the Japan public health center-based prospective diabetes study. <i>PLoS ONE</i> , 2013 , 8, e84542	3.7	2
107	Association of body mass index with risk of age-related cataracts in a middle-aged Japanese population: the JPHC Study. <i>Environmental Health and Preventive Medicine</i> , 2010 , 15, 367-73	4.2	2
106	Ethnic differences in blood pressure level between Japanese school children in Bolivia and native Bolivians. <i>International Journal of Epidemiology</i> , 1989 , 18, 100-4	7.8	2
105	Sugary drink consumption and risk of kidney and bladder cancer in Japanese adults. <i>Scientific Reports</i> , 2021 , 11, 21701	4.9	2
104	Identifying the Profile of -Negative Gastric Cancers: A Case-Only Analysis within the Stomach Cancer Pooling (StoP) Project. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021 ,	4	2
103	Changes and differences in eating habits of immigrants from Okinawa and those from the Japanese mainland in the republic of Bolivia.. [<i>Minzoku Eisei</i>] <i>Race Hygiene</i> , 1989 , 55, 124-132		2
102	Epidemiology of nonmelanoma skin cancer in Japan: Occupational type, lifestyle, and family history of cancer. <i>Cancer Science</i> , 2020 , 111, 4257-4265	6.9	2
101	Myopia, corneal endothelial cell density and morphology in a Japanese population-based cross-sectional study: the JPHC-NEXT Eye Study. <i>Scientific Reports</i> , 2021 , 11, 6366	4.9	2
100	Apolipoprotein A2 Isoforms in Relation to the Risk of Myocardial Infarction: A Nested Case-Control Analysis in the JPHC Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 483-490	4	2
99	Serum anti-DIDO1, anti-CPSF2, and anti-FOXJ2 antibodies as predictive risk markers for acute ischemic stroke. <i>BMC Medicine</i> , 2021 , 19, 131	11.4	2
98	Smoking cessation, weight gain and risk of cardiovascular disease. <i>Heart</i> , 2021 ,	5.1	2
97	Family History and Gastric Cancer Risk: A Pooled Investigation in the Stomach Cancer Pooling (STOP) Project Consortium. <i>Cancers</i> , 2021 , 13,	6.6	2
96	Comparison of plasma levels of obesity-related biomarkers among Japanese populations in Tokyo, Japan, Sã Paulo, Brazil, and Hawaii, USA. <i>European Journal of Cancer Prevention</i> , 2016 , 25, 41-9	2	2
95	Plasma C-peptide and glycated albumin and subsequent risk of cancer: From a large prospective case-cohort study in Japan. <i>International Journal of Cancer</i> , 2019 , 144, 718-729	7.5	2
94	Family history of cancer and subsequent risk of cancer: A large-scale population-based prospective study in Japan. <i>International Journal of Cancer</i> , 2020 , 147, 331-337	7.5	2
93	Cross-Sectional Association Between Employment Status and Self-Rated Health Among Middle-Aged Japanese Women: The Influence of Socioeconomic Conditions and Work-Life Conflict. <i>Journal of Epidemiology</i> , 2020 , 30, 396-403	3.4	2
92	Alcohol Drinking and Bladder Cancer Risk From a Pooled Analysis of Ten Cohort Studies in Japan. <i>Journal of Epidemiology</i> , 2020 , 30, 309-313	3.4	2
91	Dietary fiber intake and risk of gastric cancer: The Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2021 , 148, 2664-2673	7.5	2

90	Intake of Vegetables and Fruits and the Risk of Cataract Incidence in a Japanese Population: The Japan Public Health Center-Based Prospective Study. <i>Journal of Epidemiology</i> , 2021 , 31, 21-29	3.4	2
89	Association Between Dietary Fish and PUFA Intake in Midlife and Dementia in Later Life: The JPHC Saku Mental Health Study. <i>Journal of Alzheimer's Disease</i> , 2021 , 79, 1091-1104	4.3	2
88	Association of serum levels of antibodies against ALDOA and FH4 with transient ischemic attack and cerebral infarction. <i>BMC Neurology</i> , 2021 , 21, 274	3.1	2
87	Association of sugary drink consumption with all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. <i>Preventive Medicine</i> , 2021 , 148, 106561	4.3	2
86	Dietary heterocyclic aromatic amine intake and cancer risk: epidemiological evidence from Japanese studies. <i>Genes and Environment</i> , 2021 , 43, 33	2.8	2
85	Reliability of self-reported questionnaire for epidemiological investigation of Helicobacter pylori eradication in a population-based cohort study. <i>Scientific Reports</i> , 2021 , 11, 15605	4.9	2
84	Female reproductive factors and risk of external causes of death among women: The Japan Public Health Center-based Prospective Study (JPHC Study). <i>Scientific Reports</i> , 2019 , 9, 14329	4.9	1
83	Lack of social support and social trust as potential risk factors for dry eye disease: JPHC-NEXT study. <i>Ocular Surface</i> , 2019 , 17, 278-284	6.5	1
82	Physical activity and subsequent risk of kidney, bladder and upper urinary tract cancer in the Japanese population: the Japan Public Health Centre-based Prospective Study. <i>British Journal of Cancer</i> , 2019 , 120, 571-574	8.7	1
81	Estimation of the performance of a risk prediction model for gastric cancer occurrence in Japan: Evidence from a small external population. <i>Cancer Epidemiology</i> , 2020 , 67, 101766	2.8	1
80	Impact of alcohol drinking on cancer risk with consideration of flushing response: The Japan Public Health Center-based Prospective Study Cohort (JPHC study). <i>Preventive Medicine</i> , 2020 , 133, 106026	4.3	1
79	Relationship between nerve fiber layer defect and the presence of epiretinal membrane in a Japanese population: The JPHC-NEXT Eye Study. <i>Scientific Reports</i> , 2020 , 10, 779	4.9	1
78	Occupational exposures and odds of gastric cancer: a StoP project consortium pooled analysis. <i>International Journal of Epidemiology</i> , 2020 , 49, 422-434	7.8	1
77	IGHG, IGKC, and FCGR genes and endogenous antibody responses to GARP in patients with breast cancer and matched controls. <i>Human Immunology</i> , 2018 , 79, 632-637	2.3	1
76	Relationship between the variety of food consumed and sufficient nutrition in middle-aged Japanese. <i>Nutrition and Food Science</i> , 2013 , 43, 344-355	1.5	1
75	Dietary factors in gastrointestinal tract cancers – An Asian perspective. <i>Acta Oncologica</i> , 2007 , 46, 405-406	3.2	1
74	Chip ligating human genomic DNA serves as storage material and template for polymerase chain reaction. <i>Biotechnology Letters</i> , 2003 , 25, 509-12	3	1
73	Cancer Patterns and Lifestyle Among Japanese Brazilian in Sao Paulo. <i>Journal of Epidemiology</i> , 1996 , 6, 169-173	3.4	1

72	Association between C-reactive protein and risk of overall and 18 site-specific cancers in a Japanese case-cohort.. <i>British Journal of Cancer</i> , 2022 ,	8.7	1
71	Excess body fatness during early to mid-adulthood and survival from colorectal and breast cancer: a pooled analysis of five international cohort studies. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021 ,	4	1
70	Meat consumption and gastric cancer risk: The Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2021 ,	7	1
69	Alcohol intake and stomach cancer risk in Japan: A pooled analysis of six cohort studies. <i>Cancer Science</i> , 2021 ,	6.9	1
68	Association of coffee, green tea, and caffeine with the risk of dementia in older Japanese people. <i>Journal of the American Geriatrics Society</i> , 2021 , 69, 3529-3544	5.6	1
67	GWAS identifies nine nephrolithiasis susceptibility loci related with metabolic metabolic and crystallization pathways		1
66	Epidemiology of Esophageal Squamous Cell Carcinoma 2020 , 1-13		1
65	Associations between reproductive factors and biliary tract cancers in women from the Biliary Tract Cancers Pooling Project. <i>Journal of Hepatology</i> , 2020 , 73, 863-872	13.4	1
64	????????? ?????? Nihon Nyugan Kenshin Gakkaishi (Journal of Japan Association of Breast Cancer Screening), 2010 , 19, 4-15	0.1	1
63	Relationship between Meat/Fish Consumption and Biliary Tract Cancer: The Japan Public Health Center-Based Prospective Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020 , 29, 95-102	4	1
62	Soy Intake and Colorectal Cancer Risk: Results from a Pooled Analysis of Prospective Cohort Studies Conducted in China and Japan. <i>Journal of Nutrition</i> , 2020 , 150, 2442-2450	4.1	1
61	Long-term antihypertensive drug use and risk of cancer: The Japan Public Health Center-based prospective study. <i>Cancer Science</i> , 2021 , 112, 1997-2005	6.9	1
60	Reproductive Factors and Lung Cancer Risk among Never-Smoking Japanese Women with 21 Years of Follow-Up: A Cohort Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021 , 30, 1185-1192	4	1
59	Body Mass Index, Height, Weight Change, and Subsequent Lung Cancer Risk: The Japan Public Health Center-Based Prospective Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021 , 30, 1708-1716	4.716	1
58	Serum anti-AP3D1 antibodies are risk factors for acute ischemic stroke related with atherosclerosis. <i>Scientific Reports</i> , 2021 , 11, 13450	4.9	1
57	Genotype-stratified GWAS meta-analysis reveals novel loci associated with alcohol consumption		1
56	Dietary glycemic index, glycemic load, and endometrial cancer risk: The Japan Public Health Center-based Prospective Study. <i>Cancer Science</i> , 2021 , 112, 3682-3690	6.9	1
55	A Personal Breast Cancer Risk Stratification Model Using Common Variants and Environmental Risk Factors in Japanese Females. <i>Cancers</i> , 2021 , 13,	6.6	1

54	Cruciferous vegetable intake and colorectal cancer risk: Japan public health center-based prospective study. <i>European Journal of Cancer Prevention</i> , 2019 , 28, 420-427	2	1
53	Association Between Okinawan Vegetables Consumption and Risk of Type 2 Diabetes in Japanese Communities: The JPHC Study. <i>Journal of Epidemiology</i> , 2020 , 30, 227-235	3.4	1
52	Dietary calcium and vitamin K are associated with osteoporotic fracture risk in middle-aged and elderly Japanese women, but not men: the Murakami Cohort Study. <i>British Journal of Nutrition</i> , 2021 , 125, 319-328	3.6	1
51	Effectiveness of screening using fecal occult blood test and colonoscopy on the risk of colorectal cancer: The Japan Public Health Center-based Prospective Study. <i>Journal of Epidemiology</i> , 2021 ,	3.4	1
50	Working cancer survivors' physical and mental characteristics compared to cancer-free workers in Japan: a nationwide general population-based study. <i>Journal of Cancer Survivorship</i> , 2021 , 15, 912-921	5.1	1
49	Endogenous antibody responses to mucin 1 in a large multiethnic cohort of patients with breast cancer and healthy controls: Role of immunoglobulin and Fcγ receptor genes. <i>Immunobiology</i> , 2018 , 223, 178-182	3.4	1
48	The association between plasma C-peptide concentration and the risk of prostate cancer: a nested case-control study within a Japanese population-based prospective study. <i>European Journal of Cancer Prevention</i> , 2018 , 27, 461-467	2	1
47	Prediagnostic circulating inflammation-related biomarkers and gastric cancer: A case-cohort study in Japan. <i>Cytokine</i> , 2021 , 144, 155558	4	1
46	Association of dietary intakes of vitamin B12, vitamin B6, folate, and methionine with the risk of esophageal cancer: the Japan Public Health Center-based (JPHC) prospective study. <i>BMC Cancer</i> , 2021 , 21, 982	4.8	1
45	Coffee consumption and gastric cancer: a pooled analysis from the Stomach cancer Pooling Project consortium. <i>European Journal of Cancer Prevention</i> , 2021 ,	2	1
44	Peanut Consumption and Risk of Stroke and Ischemic Heart Disease in Japanese Men and Women: The JPHC Study. <i>Stroke</i> , 2021 , 52, 3543-3550	6.7	1
43	"True" <i>Helicobacter pylori</i> infection and non-cardia gastric cancer: A pooled analysis within the Stomach Cancer Pooling (StoP) Project.. <i>Helicobacter</i> , 2022 , e12883	4.9	1
42	Association Between Physical Activity and Risk of Disabling Dementia in Japan.. <i>JAMA Network Open</i> , 2022 , 5, e224590	10.4	1
41	Salt intake and gastric cancer: a pooled analysis within the Stomach cancer Pooling (StoP) Project.. <i>Cancer Causes and Control</i> , 2022 , 33, 779	2.8	1
40	Association of Marital Status With Total and Cause-Specific Mortality in Asia. <i>JAMA Network Open</i> , 2022 , 5, e2214181	10.4	1
39	Minimum Cost Estimation of a Baseline Survey for a Molecular Epidemiology Cohort Study: Collecting Participants in a Model Region in Japan. <i>Journal of Epidemiology</i> , 2016 , 26, 522-529	3.4	0
38	Urinary Stones and Risk of Coronary Heart Disease and Stroke: the Japan Public Health Center-Based Prospective Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 1208-1215	4	0
37	Education, household income, and depressive symptoms in middle-aged and older Japanese adults. <i>BMC Public Health</i> , 2021 , 21, 2120	4.1	0

36	Effectiveness of a Cancer Risk Prediction Tool on Lifestyle Habits: A Randomized Controlled Trial. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021 , 30, 1063-1071	4	○
35	Body mass index and risk of recurrent falls in community-dwelling Japanese aged 40-74 years: The Murakami cohort study. <i>Geriatrics and Gerontology International</i> , 2021 , 21, 498-505	2.9	○
34	Relation Between Body Mass Index and Dry Eye Disease: The Japan Public Health Center-Based Prospective Study for the Next Generation. <i>Eye and Contact Lens</i> , 2021 , 47, 449-455	3.2	○
33	Quantifying the association of low-intensity and late initiation of tobacco smoking with total and cause-specific mortality in Asia. <i>Tobacco Control</i> , 2021 , 30, 328-335	5.3	○
32	Validity of a food frequency questionnaire for the estimation of total polyphenol intake estimates and its major food sources in the Japanese population: the JPHC FFQ Validation Study. <i>Journal of Nutritional Science</i> , 2021 , 10, e35	2.7	○
31	Plasma phospholipid n-3 polyunsaturated fatty acids and major depressive disorder in Japanese elderly: the Japan Public Health Center-based Prospective Study. <i>Scientific Reports</i> , 2021 , 11, 4003	4.9	○
30	Risk of stroke in cancer survivors using a propensity score-matched cohort analysis. <i>Scientific Reports</i> , 2021 , 11, 5599	4.9	○
29	Impact of reproductive factors on breast cancer incidence: Pooled analysis of nine cohort studies in Japan. <i>Cancer Medicine</i> , 2021 , 10, 2153-2163	4.8	○
28	Menstrual and reproductive factors and limitations in activities of daily living: A case-control study within the Japan Public Health Center-based Prospective Study. <i>Journal of Obstetrics and Gynaecology Research</i> , 2021 , 47, 3903-3912	1.9	○
27	Having hobbies and the risk of cardiovascular disease incidence: A Japan public health center-based study. <i>Atherosclerosis</i> , 2021 , 335, 1-7	3.1	○
26	Serum perfluoroalkyl substances and breast cancer risk in Japanese women: A case-control study. <i>Science of the Total Environment</i> , 2021 , 800, 149316	10.2	○
25	Long-term exposure to fine particle matter and all-cause mortality and cause-specific mortality in Japan: the JPHC Study.. <i>BMC Public Health</i> , 2022 , 22, 466	4.1	○
24	The association between midlife living arrangement and psychiatrist-diagnosed depression in later life: who among your family members reduces the risk of depression?. <i>Translational Psychiatry</i> , 2022 , 12, 156	8.6	○
23	Body mass index and height in relation to brain tumor risk in a Japanese population. <i>Annals of Epidemiology</i> , 2020 , 51, 1-6	6.4	
22	Body mass index and mortality among middle-aged Japanese individuals with diagnosed diabetes: The Japan Public Health Center-based prospective study (JPHC study). <i>Diabetes Research and Clinical Practice</i> , 2020 , 164, 108198	7.4	
21	P1-348 Leisure-time physical activity and breast cancer risk defined by oestrogen and progesterone receptor status: the Japan public health center-based prospective study. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, A163-A163	5.1	
20	P2-338 Association of body mass index with risk of age-related cataracts in a middle-aged Japanese population. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, A315-A316	5.1	
19	P2-214 Rice intake and type 2 diabetes in Japanese men and women: the Japan Public Health Center-Based Prospective Study. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, A280-A281	5.1	

- 18 P2-277 Alcohol drinking and primary liver cancer in Japanese: a pooled analysis of four cohort studies. *Journal of Epidemiology and Community Health*, **2011**, 65, A298-A298 5.1
- 17 P1-446 Comparison of postmenopausal endogenous sex hormones among Japanese, Japanese Brazilians, and non-Japanese Brazilians. *Journal of Epidemiology and Community Health*, **2011**, 65, A190-A191 5.1
- 16 Reply to AA Ogunleye. *American Journal of Clinical Nutrition*, **2010**, 92, 668-669 7
- 15 Reply to Brockton's letter. *Cancer Causes and Control*, **2008**, 19, 1003-1003 2.8
- 14 Endocrine disrupting chemicals in Hungarian canned foods. *Environmental Health and Preventive Medicine*, **2002**, 6, 273-5 4.2
- 13 Occupational Allergic Dermatitis Induced by an Epoxy Hardener Alkylamine. *Journal of Occupational Health*, **2002**, 44, 264-266 2.3
- 12 ?????????? ?????????? *Environmental Mutagen Research*, **2004**, 26, 233-239
- 11 Soy and Breast Cancer Prevention. *Nutrition and Disease Prevention*, **2005**, 43-72
- 10 What does a healthy diet mean to the Japanese?. *Journal of Food System Research*, **2020**, 27, 80-87 0.2
- 9 Epidemiology of ESCC **2015**, 1-12
- 8 Interaction between visfatin and resistin and the risk of colorectal adenoma.. *Journal of Clinical Oncology*, **2014**, 32, 438-438 2.2
- 7 Dietary glycemic index, glycemic load and mortality: Japan Public Health Center-based prospective study. *European Journal of Nutrition*, **2021**, 60, 4607-4620 5.2
- 6 Non-alcoholic beverages intake and risk of CVD among Japanese men and women: the Japan Public Health Center study. *British Journal of Nutrition*, **2021**, 1-8 3.6
- 5 Cross-sectional associations between the types/amounts of beverages consumed and the glycemia status: The Japan Public Health Center-based Prospective Diabetes study.. *Metabolism Open*, **2022**, 14, 100185 2.8
- 4 Association between meat intake and mortality due to all-cause and major causes of death in a Japanese population **2020**, 15, e0244007
- 3 Association between meat intake and mortality due to all-cause and major causes of death in a Japanese population **2020**, 15, e0244007
- 2 Association between meat intake and mortality due to all-cause and major causes of death in a Japanese population **2020**, 15, e0244007
- 1 Association between meat intake and mortality due to all-cause and major causes of death in a Japanese population **2020**, 15, e0244007

