

# Paola Mantellini

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9151133/publications.pdf>

Version: 2024-02-01

48  
papers

1,640  
citations

279701

23  
h-index

289141

40  
g-index

52  
all docs

52  
docs citations

52  
times ranked

1877  
citing authors

#	ARTICLE	IF	CITATIONS
1	Adenoma detection by Endocuff-assisted versus standard colonoscopy in an organized screening program: the <i>deltaVision</i> randomized controlled trial. <i>Endoscopy</i> , 2022, 54, 138-147.	1.0	20
2	Cancer screening programmes in Italy during the COVID-19 pandemic: an update of a nationwide survey on activity volumes and delayed diagnoses. <i>Annali Dell'Istituto Superiore Di Sanita</i> , 2022, 58, 16-24.	0.2	13
3	A dynamic web-based decision aid to improve informed choice in organised breast cancer screening. A pragmatic randomised trial in Italy. <i>British Journal of Cancer</i> , 2020, 123, 714-721.	2.9	22
4	Patients' experience of screening CT colonography with reduced and full bowel preparation in a randomised trial. <i>European Radiology</i> , 2019, 29, 2457-2464.	2.3	15
5	Does low-dose aspirin use for cardiovascular disease prevention reduce colorectal cancer deaths? A comparison of two cohorts in the Florence district, Italy. <i>European Journal of Cancer Prevention</i> , 2018, 27, 134-139.	0.6	8
6	Cost analysis of colorectal cancer screening with CT colonography in Italy. <i>European Journal of Health Economics</i> , 2018, 19, 735-746.	1.4	5
7	Faecal immunochemical test in subjects not attending screening computed tomography colonography and colonoscopy in a randomized trial. <i>European Journal of Cancer Prevention</i> , 2018, 27, 105-109.	0.6	4
8	Reducing harms from treatment. Sixteen years of surgery of the axilla for screen-detected breast cancers in Italy. <i>Breast</i> , 2018, 42, 15-22.	0.9	1
9	Detection rate and predictive factors of sessile serrated polyps in an organised colorectal cancer screening programme with immunochemical faecal occult blood test: the EQUiPE study (Evaluating Quality Indicators of the) <i>Tj ETQq1 1 0.784314 rgBT /Overl</i>	1.0	14
10	Impact of a new sampling buffer on faecal haemoglobin stability in a colorectal cancer screening programme by the faecal immunochemical test. <i>European Journal of Cancer Prevention</i> , 2017, 26, 285-291.	0.6	11
11	Reduced and Full-Preparation CT Colonography, Fecal Immunochemical Test, and Colonoscopy for Population Screening of Colorectal Cancer: A Randomized Trial. <i>Journal of the National Cancer Institute</i> , 2016, 108, djv319.	3.0	70
12	Appropriateness of endoscopic surveillance recommendations in organised colorectal cancer screening programmes based on the faecal immunochemical test. <i>Gut</i> , 2016, 65, 1822-1828.	6.1	25
13	Quality of colonoscopy in an organised colorectal cancer screening programme with immunochemical faecal occult blood test: the EQUiPE study (Evaluating Quality Indicators of the) <i>Tj ETQq1 1 0.784314 rgBT /Overl</i>	1.0	14
14	Colorectal cancer screening of immigrants to Italy. Figures from the 2013 National Survey. <i>Preventive Medicine</i> , 2015, 81, 132-137.	1.6	16
15	Audit system on Quality of breast cancer diagnosis and Treatment (QT): results of quality indicators on screen-detected lesions in Italy, 2011-2012. <i>Epidemiologia E Prevenzione</i> , 2015, 39, 40-7.	1.1	7
16	Breast cancer mortality trends in Italy by region and screening programme, 1980-2008. <i>Journal of Medical Screening</i> , 2014, 21, 189-193.	1.1	3
17	The impact of immunochemical faecal occult blood testing on colorectal cancer incidence. <i>Digestive and Liver Disease</i> , 2014, 46, 82-86.	0.4	105
18	Screening for colorectal cancer with FOBT, virtual colonoscopy and optical colonoscopy: study protocol for a randomized controlled trial in the Florence district (SAVE study). <i>Trials</i> , 2013, 14, 74.	0.7	27

#	ARTICLE	IF	CITATIONS
19	hr-HPV testing in the follow-up of women with cytological abnormalities and negative colposcopy. <i>British Journal of Cancer</i> , 2013, 109, 1766-1774.	2.9	14
20	Feasibility of evaluating quality cancer care using registry data and electronic health records: a population-based study. <i>International Journal for Quality in Health Care</i> , 2012, 24, 411-418.	0.9	28
21	Familial risk of colorectal cancer in subjects attending an organised screening programme. <i>Digestive and Liver Disease</i> , 2012, 44, 80-83.	0.4	11
22	Direct mailing of faecal occult blood tests for colorectal cancer screening: a randomized population study from Central Italy. <i>Journal of Medical Screening</i> , 2011, 18, 121-127.	1.1	30
23	Quality indicators in breast cancer care. <i>European Journal of Cancer</i> , 2010, 46, 2344-2356.	1.3	180
24	Immunochemical faecal occult blood test: number of samples and positivity cutoff. What is the best strategy for colorectal cancer screening?. <i>British Journal of Cancer</i> , 2009, 100, 259-265.	2.9	82
25	Cost evaluation in a colorectal cancer screening programme by faecal occult blood test in the District of Florence. <i>Journal of Medical Screening</i> , 2008, 15, 175-181.	1.1	26
26	CT colonography after incomplete colonoscopy in subjects with positive faecal occult blood test. <i>World Journal of Gastroenterology</i> , 2008, 14, 4499.	1.4	31
27	Sensitivity of latex agglutination faecal occult blood test in the Florence District population-based colorectal cancer screening programme. <i>British Journal of Cancer</i> , 2007, 96, 1750-1754.	2.9	29
28	Full-Field Digital Versus Screen-Film Mammography: Comparative Accuracy in Concurrent Screening Cohorts. <i>American Journal of Roentgenology</i> , 2007, 189, 860-866.	1.0	143
29	Breast Cancer Diagnostic Methods: Screen-Detected and Clinical Cases. An Italian Survey of Women's Experiences. <i>Tumori</i> , 2007, 93, 452-460.	0.6	3
30	Association of FOBT-assessed faecal Hb content with colonic lesions detected in the Florence screening programme. <i>British Journal of Cancer</i> , 2007, 96, 218-221.	2.9	81
31	Audit system on Quality of breast cancer diagnosis and Treatment (QT): results from the survey on screen-detected lesions in Italy, 2003-2004. <i>Epidemiologia E Prevenzione</i> , 2006, 30, 59-63.	1.1	9
32	Comparison of early performance indicators for screening projects within the European Breast Cancer Network: 1989-2000. <i>European Journal of Cancer Prevention</i> , 2005, 14, 107-116.	0.6	42
33	Second reading of screening mammograms increases cancer detection and recall rates. Results in the Florence screening programme. <i>Journal of Medical Screening</i> , 2005, 12, 103-106.	1.1	62
34	The role of arbitration of discordant reports at double reading of screening mammograms. <i>Journal of Medical Screening</i> , 2005, 12, 125-127.	1.1	26
35	Conventional versus digital mammography in the analysis of screen-detected lesions with low positive predictive value. <i>European Journal of Radiology</i> , 2005, 55, 258-263.	1.2	12
36	Colorectal cancer screening programme by faecal occult blood test in Tuscany: first round results. <i>European Journal of Cancer Prevention</i> , 2004, 13, 19-26.	0.6	68

#	ARTICLE	IF	CITATIONS
37	Interobserver agreement in the histologic diagnosis of colorectal polyps the experience of the multicenter adenoma colorectal study (SMAC). <i>Journal of Clinical Epidemiology</i> , 2003, 56, 209-214.	2.4	83
38	Colorectal Cancer Screening by Fecal Occult Blood Testing: Results of a Population-Based Experience. <i>Tumori</i> , 2000, 86, 384-388.	0.6	22
39	Do patients with rectosigmoid adenomas 5 mm or less in diameter need total colonoscopy?. <i>Gastrointestinal Endoscopy</i> , 1999, 50, 314-321.	0.5	20
40	Cost Analysis in a Population Based Screening Programme for Colorectal Cancer: Comparison of Immunochemical and Guaiac Faecal Occult Blood Testing. <i>Journal of Medical Screening</i> , 1997, 4, 142-146.	1.1	35
41	Distal hyperplastic polyps do not predict proximal adenomas: results from a multicentric study of colorectal adenomas. <i>Gastrointestinal Endoscopy</i> , 1997, 46, 124-130.	0.5	42
42	Experimentally-Induced Duodenal Ulcers in Rats are Associated with a Reduction of Gastric and Duodenal Calcitonin Gene-Related Peptide-Like Immunoreactivity. , 1995, , 53-59.		1
43	Ruthenium red selectively antagonizes capsaicin-induced release of vasoactive intestinal polypeptide (VIP) from the human colon. <i>Neuroscience Letters</i> , 1991, 126, 134-136.	1.0	5
44	Capsaicin-induced release of neurokinin A from muscle and mucosa of gastric corpus: Correlation with capsaicin-evoked release of calcitonin gene-related peptide. <i>Neuropeptides</i> , 1991, 19, 137-145.	0.9	25
45	Enzyme Immunoassay for Tachykinin-Like Immunoreactivity in the Guinea Pig Spinal Cord. <i>Journal of Neurochemistry</i> , 1991, 56, 281-286.	2.1	5
46	Decrease of Duodenal Calcitonin Gene-Related Peptide- and Substance P-Like Immunoreactivity in Rat Duodenal Ulcers. <i>Advances in Experimental Medicine and Biology</i> , 1991, 298, 129-135.	0.8	8
47	Duodenal SP-like immunoreactivity is decreased in experimentally-induced duodenal ulcers. <i>Neuroscience Letters</i> , 1990, 112, 352-355.	1.0	7
48	Duodenal ulcers are associated with a depletion of duodenal calcitonin gene-related peptide-like immunoreactivity in rats. <i>European Journal of Pharmacology</i> , 1989, 164, 389-391.	1.7	15