

Fioretta Palombi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9156869/publications.pdf>

Version: 2024-02-01

15
papers

535
citations

840776

11
h-index

996975

15
g-index

15
all docs

15
docs citations

15
times ranked

769
citing authors

#	ARTICLE	IF	CITATIONS
1	Relevance of lysosomal Ca ²⁺ signalling machinery in cancer. <i>Cell Calcium</i> , 2022, 102, 102539.	2.4	8
2	Naringenin is a powerful inhibitor of SARS-CoV-2 infection in vitro. <i>Pharmacological Research</i> , 2021, 163, 105255.	7.1	88
3	The Discovery of Naringenin as Endolysosomal Two-Pore Channel Inhibitor and Its Emerging Role in SARS-CoV-2 Infection. <i>Cells</i> , 2021, 10, 1130.	4.1	20
4	Loss of Two-Pore Channel 2 (TPC2) Expression Increases the Metastatic Traits of Melanoma Cells by a Mechanism Involving the Hippo Signalling Pathway and Store-Operated Calcium Entry. <i>Cancers</i> , 2020, 12, 2391.	3.7	22
5	Could the Inhibition of Endo-Lysosomal Two-Pore Channels (TPCs) by the Natural Flavonoid Naringenin Represent an Option to Fight SARS-CoV-2 Infection?. <i>Frontiers in Microbiology</i> , 2020, 11, 970.	3.5	33
6	Naringenin Impairs Two-Pore Channel 2 Activity And Inhibits VEGF-Induced Angiogenesis. <i>Scientific Reports</i> , 2017, 7, 5121.	3.3	77
7	Ned-19 inhibition of parasite growth and multiplication suggests a role for NAADP mediated signalling in the asexual development of <i>Plasmodium falciparum</i> . <i>Malaria Journal</i> , 2017, 16, 366.	2.3	5
8	NAADP-Dependent Ca ²⁺ Signaling Controls Melanoma Progression, Metastatic Dissemination and Neoangiogenesis. <i>Scientific Reports</i> , 2016, 6, 18925.	3.3	35
9	The Adherent/Invasive <i>Escherichia coli</i> Strain LF82 Invades and Persists in Human Prostate Cell Line RWPE-1, Activating a Strong Inflammatory Response. <i>Infection and Immunity</i> , 2016, 84, 3105-3113.	2.2	24
10	Visual evoked potentials of Niemann-Pick type C1 mice reveal an impairment of the visual pathway that is rescued by 2-hydroxypropyl- β -cyclodextrin. <i>Orphanet Journal of Rare Diseases</i> , 2015, 10, 133.	2.7	21
11	Regulation of Angiogenic Functions by Angiopoietins through Calcium-Dependent Signaling Pathways. <i>BioMed Research International</i> , 2015, 2015, 1-14.	1.9	16
12	VEGF-induced neoangiogenesis is mediated by NAADP and two-pore channel-2 ⁺ dependent Ca ²⁺ signaling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, E4706-15.	7.1	138
13	Cell-cell interactions in the local control of seminiferous tubule contractility. <i>Contraception</i> , 2002, 65, 289-291.	1.5	8
14	Dual control of seminiferous tubule contractility mediated by ET A and ET B endothelin receptor subtypes. <i>FASEB Journal</i> , 1997, 11, 276-286.	0.5	31
15	A neurofibrillar body in the perikaryon of nervus terminalis ganglion cells in teleosts. <i>Cell and Tissue Research</i> , 1969, 93, 395-403.	2.9	9