

# Kaori Ide

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/916768/kaori-ide-publications-by-year.pdf>

**Version:** 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

33  
papers

289  
citations

10  
h-index

16  
g-index

36  
ext. papers

331  
ext. citations

1.7  
avg, IF

2.48  
L-index

| #  | Paper  | IF  | Citations |
|----|--|-----|-----------|
| 33 | Narrow-band ultraviolet B therapy attenuates cutaneous T-cell responses in hapten-induced, experimental contact dermatitis in beagles. <i>Veterinary Dermatology</i> , <b>2021</b> , 32, 605-e161  | 1.8 |           |
| 32 | Effects of age, sex, and breed on the composition of free extractable ceramides in the stratum corneum of healthy dogs. <i>Veterinary Research Communications</i> , <b>2021</b> , 1  | 2.9 | 0         |
| 31 | Kestose supplementation exerts bifidogenic effect within fecal microbiota and increases fecal butyrate concentration in dogs. <i>Journal of Veterinary Medical Science</i> , <b>2020</b> , 82, 1-8   | 1.1 | 11        |
| 30 | Therapeutic Potential of an Endolysin Derived from Kayvirus S25-3 for Staphylococcal Impetigo. <i>Viruses</i> , <b>2019</b> , 11,  | 6.2 | 19        |
| 29 | Progenitor cells expressing nestin, a neural crest stem cell marker, differentiate into outer root sheath keratinocytes. <i>Veterinary Dermatology</i> , <b>2019</b> , 30, 365-e107  | 1.8 | 2         |
| 28 | First identification of a single amino acid change in the spike protein region of feline coronavirus detected from a coronavirus-associated cutaneous nodule in a cat. <i>Journal of Feline Medicine and Surgery Open Reports</i> , <b>2018</b> , 4, 2055116918801385        | 0.5 | 1         |
| 27 | Staphylococcus aureus penetrate the interkeratinocyte spaces created by skin-infiltrating neutrophils in a mouse model of impetigo <b>2017</b> , 141-148   |     |           |
| 26 | Staphylococcus aureus penetrate the interkeratinocyte spaces created by skin-infiltrating neutrophils in a mouse model of impetigo. <i>Veterinary Dermatology</i> , <b>2017</b> , 28, 126-e27  | 1.8 | 2         |
| 25 | Re-evaluation of the Cefovecin Disk Diffusion Test for Predicting Oxacillin-resistance in Staphylococcus pseudintermedius Isolated from Dogs. <i>The Japanese Journal of Veterinary Dermatology</i> , <b>2017</b> , 23, 73-76  | 0   |           |
| 24 | Comparison of the expression, activity, and fecal concentration of intestinal alkaline phosphatase between healthy dogs and dogs with chronic enteropathy. <i>American Journal of Veterinary Research</i> , <b>2016</b> , 77, 721-9  | 1.1 | 9         |
| 23 | Canine Dermatomyositis-like Skin Lesions in a Shiba Inu. <i>The Japanese Journal of Veterinary Dermatology</i> , <b>2015</b> , 21, 89  | 0   |           |
| 22 | Epidermal structure created by canine hair follicle keratinocytes enriched with bulge cells in a three-dimensional skin equivalent model in vitro: implications for regenerative therapy of canine epidermis. <i>Veterinary Dermatology</i> , <b>2013</b> , 24, 77-83.e19-20 | 1.8 | 10        |
| 21 | Usefulness of cefovecin disk-diffusion test for predicting mecA gene-containing strains of Staphylococcus pseudintermedius and clinical efficacy of cefovecin in dogs with superficial pyoderma. <i>Veterinary Dermatology</i> , <b>2013</b> , 24, 162-7.e35-6               | 1.8 | 3         |
| 20 | Skin lipid profiling in normal and seborrhoeic shih tzu dogs. <i>Veterinary Dermatology</i> , <b>2013</b> , 24, 84-9.e21-21.8  |     | 7         |
| 19 | Transcription profile of chemokine receptors, cytokines and cytotoxic markers in peripheral blood of dogs with epitheliotropic cutaneous lymphoma. <i>Veterinary Dermatology</i> , <b>2013</b> , 24, 628-31, e155  | 1.8 | 0         |
| 18 | Gene transcription analysis in lesional skin of canine epitheliotropic cutaneous lymphoma using quantitative real-time RT-PCR. <i>Veterinary Immunology and Immunopathology</i> , <b>2011</b> , 144, 329-36  | 2   | 6         |
| 17 | Staphylococcus pseudintermedius exfoliative toxin EX1 selectively digests canine desmoglein 1 and causes subcorneal clefts in canine epidermis. <i>Veterinary Dermatology</i> , <b>2011</b> , 22, 319-26   | 1.8 | 30        |

|    |  |     |    |
|----|--|-----|----|
| 16 | Alteration of stratum corneum ceramide profiles in spontaneous canine model of atopic dermatitis. <i>Experimental Dermatology</i> , <b>2011</b> , 20, 732-6  | 4   | 45 |
| 15 | Identification of a novel <i>Staphylococcus pseudintermedius</i> exfoliative toxin gene and its prevalence in isolates from canines with pyoderma and healthy dogs. <i>FEMS Microbiology Letters</i> , <b>2010</b> , 312, 169-75                             | 2.9 | 35 |
| 14 | Two dogs with juvenile-onset skin diseases with involvement of extremities. <i>Journal of Veterinary Medical Science</i> , <b>2010</b> , 72, 1513-6  | 1.1 | 2  |
| 13 | Expression analysis of desmosomal components of the novel canine epidermal keratinocyte cell line (MSCEK). <i>Journal of Veterinary Medical Science</i> , <b>2010</b> , 72, 1479-82  | 1.1 | 3  |
| 12 | Fibrodysplasia ossificans progressiva in a Maine Coon cat with prominent ossification in dorsal muscle. <i>Journal of Veterinary Medical Science</i> , <b>2009</b> , 71, 1649-52   | 1.1 | 15 |
| 11 | Disseminated histiocytic sarcoma with excessive hemophagocytosis in a cat. <i>Journal of Veterinary Medical Science</i> , <b>2009</b> , 71, 817-20   | 1.1 | 26 |
| 10 | Quantitative analysis of mRNA transcripts of Hox, SHH, PTCH, Wnt, and Fzd genes in canine hematopoietic progenitor cells and various in vitro colonies differentiated from the cells. <i>Journal of Veterinary Medical Science</i> , <b>2009</b> , 71, 69-77 | 1.1 | 4  |
| 9  | Investigation of various methods for the cryopreservation of canine bone marrow-derived CD34(+) cells. <i>Journal of Veterinary Medical Science</i> , <b>2008</b> , 70, 1211-7   | 1.1 | 6  |
| 8  | Induction of chemoresistance in a cultured canine cell line by retroviral transduction of the canine multidrug resistance 1 gene. <i>American Journal of Veterinary Research</i> , <b>2007</b> , 68, 95-100  | 1.1 | 7  |
| 7  | A retrospective study and gene analysis of canine sterile panniculitis. <i>Journal of Veterinary Medical Science</i> , <b>2007</b> , 69, 915-24  | 1.1 | 26 |
| 6  | Canine PHA-stimulated adherent cell enhance interferon-gamma production and proliferation of autologous peripheral blood mononuclear cells. <i>Veterinary and Comparative Oncology</i> , <b>2005</b> , 3, 25-31  | 2.5 |    |
| 5  | Generation of canine dendritic cells from peripheral blood mononuclear cells. <i>Journal of Veterinary Medical Science</i> , <b>2003</b> , 65, 663-9   | 1.1 | 17 |
| 4  | Enhancement of reactive oxygen species production from canine blood leukocytes by human recombinant interleukin-12. <i>Veterinary Immunology and Immunopathology</i> , <b>2003</b> , 93, 1-8   | 2   | 2  |
| 3  | Usefulness of Cefovecin Disk-Diffusion Test for Predicting MecaGene-Containing Strains of <i>Staphylococcus Pseudintermedius</i> and Clinical Efficacy of Cefovecin in Dogs with Superficial Pyoderma 176-181  |     |    |
| 2  | Skin Lipid Profiling in Normal and Seborrhoeic Shih Tzu Dogs 92-97   |     |    |
| 1  | Epidermal Structure Created by Canine Hair Follicle Keratinocytes Enriched with Bulge Cells in a Three-Dimensional Skin Equivalent Model in Vitro: Implications for Regenerative Therapy of Canine Epidermis 85-91   |     |    |