Jianliang Xu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9170362/publications.pdf Version: 2024-02-01



Ιμανιμανίς Χιι

#	Article	IF	CITATIONS
1	mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants. Nature, 2021, 592, 616-622.	27.8	1,232
2	Myc Regulates Chromatin Decompaction and Nuclear Architecture during B Cell Activation. Molecular Cell, 2017, 67, 566-578.e10.	9.7	174
3	Nanobodies from camelid mice and llamas neutralize SARS-CoV-2 variants. Nature, 2021, 595, 278-282.	27.8	154
4	An Anticancer Effect of Curcumin Mediated by Down-Regulating Phosphatase of Regenerating Liver-3 Expression on Highly Metastatic Melanoma Cells. Molecular Pharmacology, 2009, 76, 1238-1245.	2.3	76
5	APE1 is dispensable for S-region cleavage but required for its repair in class switch recombination. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, 17242-17247.	7.1	33
6	C-terminal region of activation-induced cytidine deaminase (AID) is required for efficient class switch recombination and gene conversion. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, 2253-2258.	7.1	25
7	<scp>SAMHD</scp> 1â€mediated <scp>dNTP</scp> degradation is required for efficient <scp>DNA</scp> repair during antibody class switch recombination. EMBO Journal, 2020, 39, e102931.	7.8	23
8	VEGF Promotes the Transcription of the Human PRL-3 Gene in HUVEC through Transcription Factor MEF2C. PLoS ONE, 2011, 6, e27165.	2.5	22
9	<i>Dregea volubilis</i> Ameliorates Concanavalin A-Induced Liver Injury by Facilitating Apoptosis of Activated T Cells. Experimental Biology and Medicine, 2008, 233, 1124-1132.	2.4	8
10	Activation-induced cytidine deaminase is dispensable for virus-mediated liver and skin tumor development in mouse models. International Immunology, 2014, 26, 397-406.	4.0	3