

Tomasz StokÅ,osa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/918138/publications.pdf>

Version: 2024-02-01

105
papers

2,865
citations

218677

26
h-index

175258

52
g-index

110
all docs

110
docs citations

110
times ranked

4305
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | The Clinical Tumor Lysis Syndrome in a Patient with Mixed Phenotype Acute Leukemia Undergoing Induction with Venetoclax and Azacitidine – a Case Report. <i>Chemotherapy</i> , 2022, , . | 1.6 | 2 |
| 2 | Tracking Clonal Evolution of Multiple Myeloma Using Targeted Next-Generation DNA Sequencing. <i>Biomedicines</i> , 2022, 10, 1674. | 3.2 | 3 |
| 3 | Whole-exome sequencing in patients with protein aggregate myopathies reveals causative mutations associated with novel atypical phenotypes. <i>Neurological Sciences</i> , 2021, 42, 2819-2827. | 1.9 | 2 |
| 4 | Predictive significance of selected gene mutations in relapsed and refractory chronic lymphocytic leukemia patients treated with ibrutinib. <i>European Journal of Haematology</i> , 2021, 106, 320-326. | 2.2 | 2 |
| 5 | Co-occurrence of unclassified myeloproliferative neoplasm and giant cell arteritis in a patient treated with allogeneic hematopoietic stem cell transplantation: a case report and literature review. <i>Central-European Journal of Immunology</i> , 2021, 46, 121-126. | 1.2 | 1 |
| 6 | In vivo, ex vivo and in vitro dasatinib activity in chronic lymphocytic leukemia. <i>Oncology Letters</i> , 2021, 21, 285. | 1.8 | 4 |
| 7 | Epithelial Cells of Deep Infiltrating Endometriosis Harbor Mutations in Cancer Driver Genes. <i>Cells</i> , 2021, 10, 749. | 4.1 | 8 |
| 8 | Gene Expression Profiling Predicts Sensitivity of Chronic Lymphocytic Leukemia Cells to Dasatinib. <i>HemaSphere</i> , 2021, 5, e514. | 2.7 | 0 |
| 9 | Results of Polish Adult Leukemia Study Group (PALG) project assessing TP53 mutations with next-generation sequencing technology in relapsed and refractory chronic lymphocytic leukemia patients – an 18-month update. <i>Acta Haematologica Polonica</i> , 2021, 52, 94-102. | 0.3 | 0 |
| 10 | Differential Regulation of Telomeric Complex by BCR-ABL1 Kinase in Human Cellular Models of Chronic Myeloid Leukemia – From Single Cell Analysis to Next-Generation Sequencing. <i>Genes</i> , 2020, 11, 1145. | 2.4 | 10 |
| 11 | First familial cases of type 2 congenital erythrocytosis (ECYT2) with a Chuvash pathogenic variant in VHL gene in Poland: example of the clinical utility of next-generation sequencing in diagnostics of orphan diseases. <i>Acta Haematologica Polonica</i> , 2020, 51, 220-225. | 0.3 | 0 |
| 12 | Wenetoklaks w monoterapii przewlekłej, ciężkiej limfocytowej przed powtórny przeszczepieniem allogenicznym krwiotwórczych komórek macierzystych. <i>Hematologia</i> , 2020, 11, 95-100. | 0.0 | 0 |
| 13 | Clonal Hematopoiesis with Somatic Mutations in "AYA" Generation of Patients with Chronic Myeloid Leukemia. <i>Blood</i> , 2020, 136, 23-24. | 1.4 | 0 |
| 14 | Germline missense NF1 mutation in an elderly patient with a blastic plasmacytoid dendritic cell neoplasm. <i>International Journal of Hematology</i> , 2019, 110, 102-106. | 1.6 | 5 |
| 15 | Predictive Significance of Selected Gene Mutations Identified Using Next Generation Sequencing in Relapsed and Refractory Chronic Lymphocytic Leukemia Patients Treated with Ibrutinib. <i>Blood</i> , 2019, 134, 5456-5456. | 1.4 | 0 |
| 16 | ASXL1 Mutations Detectable at Diagnosis May Predict Response to Imatinib in Patients with Chronic Myeloid Leukemia. <i>Blood</i> , 2019, 134, 4148-4148. | 1.4 | 4 |
| 17 | DNA methylation signature in blood does not predict calendar age in patients with chronic lymphocytic leukemia but may alert to the presence of disease. <i>Forensic Science International: Genetics</i> , 2018, 34, e15-e17. | 3.1 | 13 |
| 18 | Tyrosine kinase inhibitor-induced defects in DNA repair sensitize FLT3(ITD)-positive leukemia cells to PARP1 inhibitors. <i>Blood</i> , 2018, 132, 67-77. | 1.4 | 54 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | The Role of Shelterin Complex and Post-Translational Non-Enzymatic Modification in Telomere Maintenance in Chronic Myeloid Leukemia. <i>Blood</i> , 2018, 132, 5426-5426. | 1.4 | 0 |
| 20 | Genomic landscape of human erythroleukemia K562 cell line, as determined by next-generation sequencing and cytogenetics. <i>Acta Haematologica Polonica</i> , 2017, 48, 343-349. | 0.3 | 3 |
| 21 | Mutational Analysis of Recurrent Meningioma Progressing From Atypical to Rhabdoid Subtype. <i>World Neurosurgery</i> , 2017, 97, 754.e1-754.e6. | 1.3 | 9 |
| 22 | Titin Truncating Variants in Dilated Cardiomyopathy â€“ Prevalence and Genotype-Phenotype Correlations. <i>PLoS ONE</i> , 2017, 12, e0169007. | 2.5 | 63 |
| 23 | Gene expression and mutation-guided synthetic lethality eradicates proliferating and quiescent leukemia cells. <i>Journal of Clinical Investigation</i> , 2017, 127, 2392-2406. | 8.2 | 64 |
| 24 | Normal ABL1 is a tumor suppressor and therapeutic target in human and mouse leukemias expressing oncogenic ABL1 kinases. <i>Blood</i> , 2016, 127, 2131-2143. | 1.4 | 32 |
| 25 | A rare mutation in a rare tumorâ€™s SMARCB1-deficient malignant glomus tumor. <i>Genes Chromosomes and Cancer</i> , 2016, 55, 107-109. | 2.8 | 8 |
| 26 | PPAR γ ligands increase antileukemic activity of second- and third-generation tyrosine kinase inhibitors in chronic myeloid leukemia cells. <i>Blood Cancer Journal</i> , 2016, 6, e377-e377. | 6.2 | 26 |
| 27 | Next Generation Sequencing in Oncology. , 2016, , 63-74. | | 0 |
| 28 | Role of Shelterin Complex and Alternative Telomere Lengthening in Genomic Instability and Disease Progression in Chronic Myeloid Leukemia. <i>Blood</i> , 2016, 128, 1880-1880. | 1.4 | 0 |
| 29 | Detailed Clinical, Immunological and Molecular Analysis of NOTCH1, SF3B1 and MYD88 mutations in Chronic Lymphocytic Leukemia Patients Reveals Accumulation of Negative Prognostic Features in NOTCH1 and SF3B1 mutated Individuals. <i>Blood</i> , 2016, 128, 5570-5570. | 1.4 | 0 |
| 30 | Imatinib in the treatment of chronic myeloid leukemia: current perspectives on optimal dose. <i>Blood and Lymphatic Cancer: Targets and Therapy</i> , 2015, , 101. | 2.7 | 0 |
| 31 | A study in Polish patients with cardiomyopathy emphasizes pathogenicity of phospholamban (PLN) mutations at amino acid position 9 and low penetrance of heterozygous null PLN mutations. <i>BMC Medical Genetics</i> , 2015, 16, 21. | 2.1 | 28 |
| 32 | Somatic Mutations in Commonly Mutated Genes in Myeloid Malignancies May Preexist or Arise in the Course of Chronic Myeloid Leukemia - Different Scenarios of Progression Revealed By Targeted Next-Generation Sequencing. <i>Blood</i> , 2015, 126, 2771-2771. | 1.4 | 1 |
| 33 | HIF1-Alpha and MYC Transcription Factor Signatures in B-Cell Acute Lymphoblastic Leukemia Are Associated with Positive Minimal Residual Disease Status: Therapeutic Implications. <i>Blood</i> , 2015, 126, 1436-1436. | 1.4 | 0 |
| 34 | Differential expression of BIRC family genes in chronic myeloid leukaemia â€“ BIRC3 and BIRC8 as potential new candidates to identify disease progression. <i>British Journal of Haematology</i> , 2014, 164, 740-742. | 2.5 | 9 |
| 35 | Statins inhibit ABCB1 and ABCG2 drug transporter activity in chronic myeloid leukemia cells and potentiate antileukemic effects of imatinib. <i>Experimental Hematology</i> , 2014, 42, 439-447. | 0.4 | 21 |
| 36 | BRAF â€“ A new player in hematological neoplasms. <i>Blood Cells, Molecules, and Diseases</i> , 2014, 53, 77-83. | 1.4 | 26 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Effects of First and Next-Generation Tyrosine Kinase Inhibitors on Telomere-Mediated Chromosomal Instability in Chronic Myeloid Leukemia Cells. <i>Blood</i> , 2014, 124, 5510-5510. | 1.4 | 2 |
| 38 | HDAC Inhibitors As Potential New Agents Improving the Efficacy of Monoclonal Antibodies. <i>Blood</i> , 2014, 124, 3641-3641. | 1.4 | 0 |
| 39 | Gene Expression and Mutation Analysis (GEMA) â€“Guided Precision Medicine Targeting PARP1 to Induce Synthetic Lethality in DNA-PK â€“Deficient Quiescent and BRCA-Deficient Proliferating Leukemia Stem and Progenitor Cells. <i>Blood</i> , 2014, 124, 480-480. | 1.4 | 0 |
| 40 | Genomic instability may originate from imatinib-refractory chronic myeloid leukemia stem cells. <i>Blood</i> , 2013, 121, 4175-4183. | 1.4 | 105 |
| 41 | BCR-ABL1 kinase inhibits uracil DNA glycosylase UNG2 to enhance oxidative DNA damage and stimulate genomic instability. <i>Leukemia</i> , 2013, 27, 629-634. | 7.2 | 36 |
| 42 | Diverse mechanisms of mTOR activation in chronic and blastic phase of chronic myelogenous leukemia. <i>Experimental Hematology</i> , 2013, 41, 462-469. | 0.4 | 6 |
| 43 | Synthesis and Optical Properties of Tetraarylâ€“1,4â€“dihydropyrrolo[3,2â€“b</i>]pyrroles. <i>Asian Journal of Organic Chemistry</i> , 2013, 2, 411-415. | 2.7 | 80 |
| 44 | Anti-oxidant vitamin E prevents accumulation of imatinib-resistant BCR-ABL1 kinase mutations in CML-CP xenografts in NSG mice. <i>Leukemia</i> , 2013, 27, 2253-2254. | 7.2 | 9 |
| 45 | ARMS-PCR for detection of BRAF V600E hotspot mutation in comparison with Real-Time PCR-based techniques.. <i>Acta Biochimica Polonica</i> , 2013, 60, . | 0.5 | 16 |
| 46 | Normal ABL1 Is a Tumor Suppressor and Therapeutic Target In BCR-ABL1â€“positive Leukemias. <i>Blood</i> , 2013, 122, 1466-1466. | 1.4 | 0 |
| 47 | Differential Expression of BIRC Family Genes In The Course Of Chronic Myeloid Leukemia â€“ BIRC3 and BIRC8 As Potential New Candidates To Identify Disease Progression. <i>Blood</i> , 2013, 122, 2718-2718. | 1.4 | 0 |
| 48 | The PERK-eIF2Î± phosphorylation arm is a pro-survival pathway of BCR-ABL signaling and confers resistance to imatinib treatment in chronic myeloid leukemia cells. <i>Cell Cycle</i> , 2012, 11, 4069-4078. | 2.6 | 58 |
| 49 | Rac2-MRC-cllâ€“generated ROS cause genomic instability in chronic myeloid leukemia stem cells and primitive progenitors. <i>Blood</i> , 2012, 119, 4253-4263. | 1.4 | 147 |
| 50 | Terapia przewlekłej, białaczkowej szpikowej â€“ terapia i wyzwania na przyszłość. <i>Acta Haematologica Polonica</i> , 2012, 43, 249-257. | 0.3 | 0 |
| 51 | Bright, Color-Tunable Fluorescent Dyes Based on Î€-Expanded Diketopyrrolopyrroles. <i>Organic Letters</i> , 2012, 14, 2670-2673. | 4.6 | 79 |
| 52 | BCR-ABL1 Kinase Inhibits DNA Glycosylases to Enhance Oxidative DNA Damage and Stimulate Genomic Instability. <i>Blood</i> , 2012, 120, 520-520. | 1.4 | 1 |
| 53 | Monoubiquitinated Fanconi anemia D2 (FANCD2-Ub) is required for BCR-ABL1 kinase-induced leukemogenesis. <i>Leukemia</i> , 2011, 25, 1259-1267. | 7.2 | 5 |
| 54 | Imatinib Sensitivity in BCR-ABL1â€“Positive Chronic Myeloid Leukemia Cells Is Regulated by the Remaining Normal ABL1 Allele. <i>Cancer Research</i> , 2011, 71, 5381-5386. | 0.9 | 10 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Sorafenib Affects Membrane Complement Inhibitors and Improves Antitumor Activity of Rituximab., Blood, 2011, 118, 3723-3723. | 1.4 | 0 |
| 56 | Targeting Rac2 - Mitochondrial Respiratory Chain Complex III Signaling to Prevent Genomic Instability in Leukemia Stem and Progenitor Cells. Blood, 2011, 118, 2736-2736. | 1.4 | 0 |
| 57 | Statins Increase Antileukemic Potency of Imatinib Through the Inhibition of MDR/ABCB1 and BCRP/ABCG2 Drug Transporters Activity. Blood, 2011, 118, 2742-2742. | 1.4 | 0 |
| 58 | Preyl Transferases Are Involved in the Regulation of CD20 Levels and Influence Anti-CD20 Monoclonal Antibody-Mediated Activation of Complement-Dependent Cytotoxicity., Blood, 2011, 118, 3722-3722. | 1.4 | 0 |
| 59 | Src Family Tyrosine Kinases Are Involved in the Transcriptional Regulation of CD20 Levels. Blood, 2011, 118, 1661-1661. | 1.4 | 0 |
| 60 | Mitochondrial Respiratory Chain Complex III Causes Genomic Instability In CML-CP.. Blood, 2010, 116, 1211-1211. | 1.4 | 24 |
| 61 | Potiation of the Antileukemic Effects of Imatinib through the Modulation of BCRP/ABCG2 Activity.. Blood, 2010, 116, 3400-3400. | 1.4 | 0 |
| 62 | Proteasome Inhibition Potentiates Antitumor Effects of Photodynamic Therapy in Mice through Induction of Endoplasmic Reticulum Stress and Unfolded Protein Response. Cancer Research, 2009, 69, 4235-4243. | 0.9 | 96 |
| 63 | ROS-Induced DNA Damage Causing Genomic Instability in CML Stem and/or Progenitor Cells and in Quiescent and/or Proliferating Cells: Role of Mitochondrial Respiratory Chain Complex III.. Blood, 2009, 114, 3268-3268. | 1.4 | 1 |
| 64 | BCR/ABL Requires Fanconi Anemia D2 (FANCD2) Protein to Transform Hematopoetic Stem Cells.. Blood, 2009, 114, 3249-3249. | 1.4 | 0 |
| 65 | Zinc protoporphyrin IX, a heme oxygenase-1 inhibitor, demonstrates potent antitumor effects but is unable to potentiate antitumor effects of chemotherapeutics in mice. BMC Cancer, 2008, 8, 197. | 2.6 | 59 |
| 66 | Statins Impair Antitumor Effects of Rituximab by Inducing Conformational Changes of CD20. PLoS Medicine, 2008, 5, e64. | 8.4 | 115 |
| 67 | BCR/ABL Inhibits Mismatch Repair to Protect from Apoptosis and Induce Point Mutations. Cancer Research, 2008, 68, 2576-2580. | 0.9 | 92 |
| 68 | Ciglitazone, an agonist of peroxisome proliferator-activated receptor β , exerts potentiated cytostatic/cytotoxic effects against tumor cells when combined with lovastatin. International Journal of Oncology, 2008, , . | 3.3 | 2 |
| 69 | Monoubiquitination of the Fanconi Anemia D2 (FANCD2) Protein Regulates the Transforming Potential of BCR/ABL. Blood, 2008, 112, 3189-3189. | 1.4 | 0 |
| 70 | Erythropoietin reduces cisplatin-induced neurotoxicity without impairment of cytotoxic effects against tumor cells. International Journal of Oncology, 2007, 31, 1547-52. | 3.3 | 3 |
| 71 | Enhanced phosphorylation of Nbs1, a member of DNA repair/checkpoint complex Mre11-RAD50-Nbs1, can be targeted to increase the efficacy of imatinib mesylate against BCR/ABL-positive leukemia cells. Blood, 2007, 110, 651-660. | 1.4 | 28 |
| 72 | Potentiated antitumor effects of the combination treatment with statins and pamidronate in vitro and in vivo. International Journal of Oncology, 2007, , . | 3.3 | 6 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | BCR/ABL Kinase Inhibits Mismatch Repair To Reduce Apoptosis and Induce Point Mutations.. Blood, 2007, 110, 32-32. | 1.4 | 6 |
| 74 | BCR/ABL Kinase Elevates ROS-Mediated Oxidative DNA Damage in CML Stem/Progenitor Cells and Affects the Efficiency and Fidelity of DNA Repair To Induce Genetic Instability.. Blood, 2007, 110, 34-34. | 1.4 | 0 |
| 75 | Statins Impair Antitumor Effects of CD20 mAb by Inducing Conformational Changes of CD20.. Blood, 2007, 110, 2341-2341. | 1.4 | 0 |
| 76 | Potentiated antitumor effects of the combination treatment with statins and pamidronate in vitro and in vivo. International Journal of Oncology, 2007, 30, 1413-25. | 3.3 | 8 |
| 77 | BCR/ABL kinase induces self-mutagenesis via reactive oxygen species to encode imatinib resistance. Blood, 2006, 108, 319-327. | 1.4 | 271 |
| 78 | Heme oxygenase-1 protects tumor cells against photodynamic therapy-mediated cytotoxicity. Oncogene, 2006, 25, 3365-3374. | 5.9 | 163 |
| 79 | ATR-Chk1 Axis Protects BCR/ABL Leukemia Cells from the Lethal Effect of DNA Double-Strand Breaks. Cell Cycle, 2006, 5, 994-1000. | 2.6 | 53 |
| 80 | Enhanced Phosphorylation of Nbs1, a Member of DNA Repair/Checkpoint Complex RAD50-Mre11-Nbs1, Can Be Targeted Simultaneously with BCR/ABL Kinase To Eliminate Leukemia Cells.. Blood, 2006, 108, 2127-2127. | 1.4 | 1 |
| 81 | Impaired homologous recombination DNA repair and enhanced sensitivity to DNA damage in prostate cancer cells exposed to anchorage-independence. Oncogene, 2005, 24, 3748-3758. | 5.9 | 18 |
| 82 | The influence of photodynamic therapy on the immune response. Photodiagnosis and Photodynamic Therapy, 2005, 2, 283-298. | 2.6 | 83 |
| 83 | Direct tumor damage mechanisms of photodynamic therapy.. Acta Biochimica Polonica, 2005, 52, 339-352. | 0.5 | 222 |
| 84 | Enhanced Phosphorylation of Nbs1, a Member of the DNA Repair/Checkpoint Activation Complex Rad50/Mre11/Nbs1, Prolongs Cell Cycle S Phase and Contributes to Drug Resistance in BCR/ABL-Positive Leukemias.. Blood, 2005, 106, 2867-2867. | 1.4 | 1 |
| 85 | BCR/ABL Kinase Disrupts Formation of Mismatch Repair Complex To Induce Genomic Instability.. Blood, 2005, 106, 2864-2864. | 1.4 | 0 |
| 86 | ATR-Chk1 Axis Is Activated, but the Function of Chk1 Is Disrupted in BCR/ABL Leukemia Cells Responding to DNA Damage.. Blood, 2005, 106, 2883-2883. | 1.4 | 1 |
| 87 | Technology evaluation: SB-485232, GlaxoSmithKline. Current Opinion in Molecular Therapeutics, 2005, 7, 85-93. | 2.8 | 6 |
| 88 | Prospects for p53-based cancer therapy. Acta Biochimica Polonica, 2005, 52, 321-8. | 0.5 | 7 |
| 89 | Direct tumor damage mechanisms of photodynamic therapy. Acta Biochimica Polonica, 2005, 52, 339-52. | 0.5 | 63 |
| 90 | BCR/ABL Recruits p53 Tumor Suppressor Protein to Induce Drug Resistance. Cell Cycle, 2004, 3, 1463-1472. | 2.6 | 28 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Increased local vascular endothelial growth factor expression associated with antitumor activity of proteasome inhibitor. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2004, 9, 193-204. | 4.9 | 5 |
| 92 | BCR/ABL oncogenic kinase promotes unfaithful repair of the reactive oxygen species-dependent DNA double-strand breaks. <i>Blood</i> , 2004, 104, 3746-3753. | 1.4 | 252 |
| 93 | BCR/ABL Oncogenic Kinase Promotes Unfaithful Repair of the Reactive Oxygen Species - Dependent DNA Double-Strand Breaks.. <i>Blood</i> , 2004, 104, 712-712. | 1.4 | 2 |
| 94 | A single injection of immature dendritic cells is able to induce antitumour response against a murine colon adenocarcinoma with a low apoptotic index. <i>Oncology Reports</i> , 2002, 9, 991-4. | 2.6 | 9 |
| 95 | Lactacystin Inhibits Cathepsin A Activity in Melanoma Cell Lines. <i>Tumor Biology</i> , 2001, 22, 211-215. | 1.8 | 22 |
| 96 | Separation of cathepsin A-like enzyme and the proteasome: evidence that lactacystin/lactone is not a specific inhibitor of the proteasome. <i>International Journal of Biochemistry and Cell Biology</i> , 2000, 32, 747-757. | 2.8 | 45 |
| 97 | Lovastatin and simvastatin are modulators of the proteasome. <i>International Journal of Biochemistry and Cell Biology</i> , 2000, 32, 957-965. | 2.8 | 40 |
| 98 | Interleukin 12 and indomethacin exert a synergistic, angiogenesis-dependent antitumor activity in mice. <i>Life Sciences</i> , 2000, 66, 1223-1230. | 4.3 | 13 |
| 99 | A Combination of Retinoic Acid and Proteasome Inhibitors for the Treatment of Leukemias Is Potentially Dangerous. <i>Blood</i> , 1999, 94, 1827-1828. | 1.4 | 6 |
| 100 | The potentiated antileukemic effects of doxorubicin and interleukin-12 combination are not dependent on nitric oxide production. <i>Cancer Letters</i> , 1999, 147, 67-75. | 7.2 | 9 |
| 101 | Effective chemo-immunotherapy of L1210 leukemia in vivo using interleukin-12 combined with doxorubicin but not with cyclophosphamide, paclitaxel or cisplatin. <i>International Journal of Cancer</i> , 1998, 77, 720-727. | 5.1 | 39 |
| 102 | Erythropoietin Prevents the Development of Interleukin-12-Induced Anemia and Thrombocytopenia But Does Not Decrease Its Antitumor Activity in Mice. <i>Blood</i> , 1998, 91, 4387-4388. | 1.4 | 14 |
| 103 | Antitumor effects of the combination immunotherapy with interleukin-12 and tumor necrosis factor $\hat{\pm}$ in mice. <i>Cancer Immunology, Immunotherapy</i> , 1997, 45, 100-108. | 4.2 | 63 |
| 104 | Potential of the anti-tumor effect of actinomycin D by tumor necrosis factor $\hat{\pm}$ in mice: Correlation between in vitro and in vivo results. , 1996, 66, 374-379. | | 27 |
| 105 | Decreased natural killer cell activity in whole-blood donors does not seem to result in increased cancer incidence. <i>Transfusion</i> , 1994, 34, 359-360. | 1.6 | 7 |