## MarÃ-a Belén Muñoz-GarcÃ-a

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9184803/publications.pdf

Version: 2024-02-01



| #  | Article  | IF                | CITATIONS        |
|----|--|-------------------|------------------|
| 1  | WHAT CAN COVID-19 REMOTE LEARNING TEACH US FOR FUTURE REGULAR FACE-TO-FACE TEACHING OF GEOLOGICAL MAPPING. , 2021, , .   |                   | Ο                |
| 2  | Fluid-inclusion petrography in calcite stalagmites: Implications for entrapment processes. Journal of Sedimentary Research, 2021, 91, 1206-1226.   | 1.6               | 3                |
| 3  | Palaeoecological and palaeoenviromental reconstruction of the upper Miocene vertebrate karstic<br>site of Corral de Lobato, central-eastern Spain. Palaeogeography, Palaeoclimatology, Palaeoecology,<br>2020, 556, 109877.                            | 2.3               | 2                |
| 4  | Middle Jurassic–Early Cretaceous tectono-sedimentary evolution of the southwestern Iberian Basin<br>(central Spain): Major palaeogeographical changes in the geotectonic framework of the Western<br>Tethys. Earth-Science Reviews, 2019, 199, 102983. | 9.1               | 25               |
| 5  | Characterization of a locally deposited material on Arnela Beach (Galicia Coast, Spain). Journal of<br>Geochemical Exploration, 2017, 174, 164-171.  | 3.2               | 0                |
| 6  | Speleothem Architectural Analysis: Integrated approach for stalagmite-based paleoclimate research.<br>Sedimentary Geology, 2017, 353, 28-45.   | 2.1               | 28               |
| 7  | Comparison of speleothem fabrics and microstratigraphic stacking patterns in calcite stalagmites as indicators of paleoenvironmental change. Quaternary International, 2016, 407, 74-85.   | 1.5               | 23               |
| 8  | Long-term hydrological changes in northern Iberia (4.9–0.9 ky BP) from speleothem Mg/Ca ratios and<br>cave monitoring (Ojo GuareA±a Karst Complex, Spain). Environmental Earth Sciences, 2015, 74, 7741-7753.  | 2.7               | 15               |
| 9  | Palaeoenvironmental Interpretation of Palaeosols and Palustrine Carbonates of the Earliest<br>Cretaceous Terrestrial Ecosystems in the SerranÃa de Cuenca, Iberian Ranges, Spain. Springer Geology,<br>2014, , 1191-1195.                              | 0.3               | 1                |
| 10 | The Karstic Habitat of Spelaeogriphaceans from the Las Hoyas Fossil Site (Upper Barremian, SerranÃa de) Tj ETQc  | 0 0 0 rgB1 0.3    | /Qverlock 1      |
| 11 | The Stratigraphy and Rifting Evolution of the Oxfordian–Barremian (Upper Jurassic–Lower) Tj ETQq1 1 0.784<br>655-658.  | 314 rgBT /<br>0.3 | Overlock 10<br>1 |
| 12 | Reply to Comment by DomÃnguez-Villar on "Land surface temperature changes in Northern Iberia since<br>4000yr BP, based in l´13C of speleothems―(MartÃn-Chivelet et al., 2011). Global and Planetary Change,<br>2013, 101, 129-130.                     | 3.5               | 0                |
| 13 | The uppermost deposits of the stratigraphic succession of the Farafra Depression (Western Desert,) Tj ETQq1 1 (  | ).784314 t<br>2.0 | rg&T /Overloo    |
| 14 | The Blake geomagnetic excursion recorded in a radiometrically dated speleothem. Earth and Planetary Science Letters, 2012, 353-354, 173-181.   | 4.4               | 50               |
| 15 | Porosity and hydric behavior of typical calcite microfabrics in stalagmites. Sedimentary Geology, 2012, 265-266, 72-86.  | 2.1               | 7                |
| 16 | Land surface temperature changes in Northern Iberia since 4000yrBP, based on δ13C of speleothems.<br>Global and Planetary Change, 2011, 77, 1-12.  | 3.5               | 122              |
| 17 | Sedimentary facies and three-dimensional reconstructions of upper Oligocene meander belts from the Loranca Basin, Spain. AAPG Bulletin, 2010, 94, 241-257.   | 1.5               | 12               |