

# Francisco J Salazar

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9185283/publications.pdf>

Version: 2024-02-01

17  
papers

204  
citations

1040018

9  
h-index

1125717

13  
g-index

18  
all docs

18  
docs citations

18  
times ranked

219  
citing authors

#	ARTICLE	IF	CITATIONS
1	Sex Differences in the Renal Changes Elicited by Angiotensin II Blockade During the Nephrogenic Period. <i>Hypertension</i> , 2007, 49, 1429-1435.	2.7	50
2	Age- and Sodium-Sensitive Hypertension and Sex-Dependent Renal Changes in Rats With a Reduced Nephron Number. <i>Hypertension</i> , 2008, 51, 1184-1189.	2.7	38
3	Role of COX-2-derived metabolites in regulation of the renal hemodynamic response to norepinephrine. <i>American Journal of Physiology - Renal Physiology</i> , 2001, 281, F975-F982.	2.7	23
4	Role of angiotensin II in arterial pressure and renal hemodynamics in rats with altered renal development: age- and sex-dependent differences. <i>American Journal of Physiology - Renal Physiology</i> , 2013, 304, F33-F40.	2.7	17
5	Role of cyclooxygenase-2-derived metabolites and nitric oxide in regulating renal function. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2000, 279, R1641-R1646.	1.8	16
6	Sex-dependent hypertension and renal changes in aged rats with altered renal development. <i>American Journal of Physiology - Renal Physiology</i> , 2014, 307, F461-F470.	2.7	15
7	Renal effects induced by prolonged mPGES1 inhibition. <i>American Journal of Physiology - Renal Physiology</i> , 2014, 306, F68-F74.	2.7	13
8	Renal effects of prolonged high protein intake and COX2 inhibition on hypertensive rats with altered renal development. <i>American Journal of Physiology - Renal Physiology</i> , 2011, 301, F327-F333.	2.7	11
9	Nitric oxide, prostaglandins and angiotensin II in the regulation of renal medullary blood flow during volume expansion. <i>Journal of Physiology and Biochemistry</i> , 2016, 72, 1-8.	3.0	9
10	COX2 inhibition during nephrogenic period induces ANG II hypertension and sex-dependent changes in renal function during aging. <i>American Journal of Physiology - Renal Physiology</i> , 2014, 306, F534-F541.	2.7	6
11	Cardiac, renal and uterine hemodynamics changes throughout pregnancy in rats with a prolonged high fat diet from an early age. <i>PLoS ONE</i> , 2020, 15, e0234861.	2.5	3
12	SGLT2 inhibition potentiates the cardiovascular, renal, and metabolic effects of sGC stimulation in hypertensive rats with prolonged exposure to high-fat diet. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2022, 322, H523-H536.	3.2	2
13	Gender differences in the renal changes induced by a prolonged high-fat diet in rats with altered renal development. <i>Journal of Physiology and Biochemistry</i> , 2021, 77, 431-441.	3.0	1
14	Altered renal hemodynamic and excretory response to aminoacids when nephron number is diminished. <i>FASEB Journal</i> , 2007, 21, A894.	0.5	0
15	Gender differences in the altered renal excretory response to an acute volume expansion in rats with low glomerular number. <i>FASEB Journal</i> , 2007, 21, A1417.	0.5	0
16	Greater Renal Sensitivity to Angiotensin II in Rats with a Lower Nephron Number. <i>FASEB Journal</i> , 2008, 22, 735.2.	0.5	0
17	Altered renal hemodynamic and excretory function in rats treated with a COX2 inhibitor during the nephrogenic period. <i>FASEB Journal</i> , 2009, 23, 969.12.	0.5	0