

Gudrun F Debes

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9196385/publications.pdf>

Version: 2024-02-01

32
papers

2,907
citations

331538

21
h-index

477173

29
g-index

32
all docs

32
docs citations

32
times ranked

4190
citing authors

#	ARTICLE	IF	CITATIONS
1	Immunoregulation by antibody secreting cells in inflammation, infection, and cancer. <i>Immunological Reviews</i> , 2021, 303, 103-118.	2.8	12
2	Skin-Homing Regulatory B Cells Required for Suppression of Cutaneous Inflammation. <i>Journal of Investigative Dermatology</i> , 2021, 141, 1995-2005.e6.	0.3	10
3	Approaches to overcome flow cytometry limitations in the analysis of cells from veterinary relevant species. <i>BMC Veterinary Research</i> , 2020, 16, 83.	0.7	5
4	Lymphoplasmacellular mucositis ameliorated by $\alpha 4 \beta 7$ integrin inhibitor vedolizumab in a patient with ulcerative colitis. <i>JAAD Case Reports</i> , 2019, 5, 663-665.	0.4	1
5	B Cells and Melanoma Pathogenesis. <i>Journal of Investigative Dermatology</i> , 2019, 139, 1422-1424.	0.3	3
6	IgM Plasma Cells Reside in Healthy Skin and Accumulate with Chronic Inflammation. <i>Journal of Investigative Dermatology</i> , 2019, 139, 2477-2487.	0.3	29
7	Skin-Associated B Cells in Health and Inflammation. <i>Journal of Immunology</i> , 2019, 202, 1659-1666.	0.4	57
8	T Cell Migration from Inflamed Skin to Draining Lymph Nodes Requires Intralymphatic Crawling Supported by ICAM-1/LFA-1 Interactions. <i>Cell Reports</i> , 2017, 18, 857-865.	2.9	96
9	IL-10+ Innate-like B Cells Are Part of the Skin Immune System and Require $\alpha 4 \beta 1$ Integrin To Migrate between the Peritoneum and Inflamed Skin. <i>Journal of Immunology</i> , 2016, 196, 2514-2525.	0.4	56
10	Effector T Cell Egress via Afferent Lymph Modulates Local Tissue Inflammation. <i>Journal of Immunology</i> , 2015, 195, 3531-3536.	0.4	41
11	CRK proteins selectively regulate T cell migration into inflamed tissues. <i>Journal of Clinical Investigation</i> , 2015, 125, 1019-1032.	3.9	46
12	Timed Action of IL-27 Protects from Immunopathology while Preserving Defense in Influenza. <i>PLoS Pathogens</i> , 2014, 10, e1004110.	2.1	62
13	CXCR4 Is Dispensable for T Cell Egress from Chronically Inflamed Skin via the Afferent Lymph. <i>PLoS ONE</i> , 2014, 9, e95626.	1.1	24
14	Ovine skin-recirculating $\alpha \beta$ T cells express IFN- γ and IL-17 and exit tissue independently of CCR7. <i>Veterinary Immunology and Immunopathology</i> , 2013, 155, 87-97.	0.5	19
15	The Skin, a Novel Niche for Recirculating B Cells. <i>Journal of Immunology</i> , 2012, 188, 6027-6035.	0.4	86
16	Tissue Exit: a Novel Control Point in the Accumulation of Antigen-Specific CD8 T Cells in the Influenza A Virus-Infected Lung. <i>Journal of Virology</i> , 2012, 86, 3436-3445.	1.5	41
17	Tissue Exit: a Novel Control Point in the Accumulation of Antigen-Specific CD8 T Cells in the Influenza A Virus-Infected Lung. <i>Journal of Virology</i> , 2012, 86, 8918-8918.	1.5	0
18	CCL8 and skin T cells are an allergic attraction. <i>Nature Immunology</i> , 2011, 12, 111-112.	7.0	24

#	ARTICLE	IF	CITATIONS
19	Chemoattractant Receptors and Lymphocyte Egress from Extralymphoid Tissue: Changing Requirements during the Course of Inflammation. <i>Journal of Immunology</i> , 2010, 185, 4873-4882.	0.4	77
20	Helping and harming have something in common. <i>Nature Immunology</i> , 2009, 10, 138-140.	7.0	1
21	CD152 (CTLA-4) Determines CD4 T Cell Migration In Vitro and In Vivo. <i>PLoS ONE</i> , 2009, 4, e5702.	1.1	30
22	DCs metabolize sunlight-induced vitamin D3 to 'program' T cell attraction to the epidermal chemokine CCL27. <i>Nature Immunology</i> , 2007, 8, 285-293.	7.0	584
23	Chemotaxis Assay: Analysis of Migration of Lymphocyte Subsets. , 2006, , 418-423.		2
24	Trafficking of Lymphocyte Subpopulations. , 2006, , 154-172.		0
25	Chemotactic Responses of IL-4-, IL-10-, and IFN- γ -Producing CD4+ T Cells Depend on Tissue Origin and Microbial Stimulus. <i>Journal of Immunology</i> , 2006, 176, 557-566.	0.4	48
26	Chemokine receptor CCR7 required for T lymphocyte exit from peripheral tissues. <i>Nature Immunology</i> , 2005, 6, 889-894.	7.0	434
27	Developmental Stage, Phenotype, and Migration Distinguish Naive- and Effector/Memory-like CD4+ Regulatory T Cells. <i>Journal of Experimental Medicine</i> , 2004, 199, 303-313.	4.2	565
28	CC Chemokine Receptor 7 Expression by Effector/Memory CD4 + T Cells Depends on Antigen Specificity and Tissue Localization during Influenza A Virus Infection. <i>Journal of Virology</i> , 2004, 78, 7528-7535.	1.5	57
29	Expression of selectin ligands on murine effector and IL-10-producing CD4+T ϵ cells from non-infected and infected tissues. <i>European Journal of Immunology</i> , 2004, 34, 3070-3081.	1.6	28
30	Targeting T cell responses by selective chemokine receptor expression. <i>Seminars in Immunology</i> , 2003, 15, 277-286.	2.7	68
31	Chemotactic Responsiveness Toward Ligands for CXCR3 and CXCR4 Is Regulated on Plasma Blasts During the Time Course of a Memory Immune Response. <i>Journal of Immunology</i> , 2002, 169, 1277-1282.	0.4	323
32	In Vivo Differentiated Cytokine-Producing CD4+ T Cells Express Functional CCR7. <i>Journal of Immunology</i> , 2002, 168, 5441-5447.	0.4	78