

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/9196392/wei-xu-publications-by-citations.pdf>

**Version:** 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

28

papers

213

citations

9

h-index

13

g-index

30

ext. papers

306

ext. citations

4.1

avg, IF

2.93

L-index

#	Paper	IF	Citations
28	Low T3 syndrome is a strong prognostic predictor in diffuse large B cell lymphoma. <i>British Journal of Haematology</i> , <b>2017</b> , 177, 95-105	4.5	27
27	The impacts of zanubrutinib on immune cells in patients with chronic lymphocytic leukemia/small lymphocytic lymphoma. <i>Hematological Oncology</i> , <b>2019</b> , 37, 392-400	1.3	25
26	Circ-RPL15: a plasma circular RNA as novel oncogenic driver to promote progression of chronic lymphocytic leukemia. <i>Leukemia</i> , <b>2020</b> , 34, 919-923	10.7	24
25	Elevated absolute NK cell counts in peripheral blood predict good prognosis in chronic lymphocytic leukemia. <i>Journal of Cancer Research and Clinical Oncology</i> , <b>2018</b> , 144, 449-457	4.9	19
24	Low serum cholesterol levels predict inferior prognosis and improve NCCN-IPI scoring in diffuse large B cell lymphoma. <i>International Journal of Cancer</i> , <b>2018</b> , 143, 1884-1895	7.5	14
23	Low T3 syndrome as a predictor of poor prognosis in chronic lymphocytic leukemia. <i>International Journal of Cancer</i> , <b>2018</b> , 143, 466-477	7.5	13
22	NDRG2 mRNA levels and miR-28-5p and miR-650 activity in chronic lymphocytic leukemia. <i>BMC Cancer</i> , <b>2018</b> , 18, 1009	4.8	11
21	25-Hydroxy vitamin D deficiency predicts inferior prognosis in mantle cell lymphoma. <i>Journal of Cancer Research and Clinical Oncology</i> , <b>2020</b> , 146, 1003-1009	4.9	9
20	Circulating tumor DNA: clinical roles in diffuse large B cell lymphoma. <i>Annals of Hematology</i> , <b>2019</b> , 98, 255-269	3	9
19	Hyperfibrinogenemia is a poor prognostic factor in diffuse large B cell lymphoma. <i>Annals of Hematology</i> , <b>2018</b> , 97, 1841-1849	3	6
18	DA-EPOCH-R improves the outcome over that of R-CHOP regimen for DLBCL patients below 60 years, GCB phenotype, and those with high-risk IPI, but not for double expressor lymphoma. <i>Journal of Cancer Research and Clinical Oncology</i> , <b>2019</b> , 145, 117-127	4.9	6
17	NOTCH1 mutation and its prognostic significance in Chinese chronic lymphocytic leukemia: a retrospective study of 317 cases. <i>Cancer Medicine</i> , <b>2018</b> , 7, 1689-1696	4.8	5
16	The percentage of cells with 17p deletion and the size of 17p deletion subclones show prognostic significance in chronic lymphocytic leukemia. <i>Genes Chromosomes and Cancer</i> , <b>2019</b> , 58, 43-51	5	5
15	Diabetes mellitus predicts inferior survival in diffuse large B-cell lymphoma: a propensity score-matched analysis. <i>Cancer Management and Research</i> , <b>2019</b> , 11, 2849-2870	3.6	4
14	Prognostic significance of serum immunoglobulin paraprotein in patients with diffuse large B cell lymphoma. <i>British Journal of Haematology</i> , <b>2018</b> , 182, 131-134	4.5	4
13	Prognostic value of 25-hydroxy vitamin D in extranodal NK/T cell lymphoma. <i>Annals of Hematology</i> , <b>2021</b> , 100, 445-453	3	4
12	Prognostic nutritional index, a novel biomarker which predicts worse prognosis in diffuse large B cell lymphoma. <i>Leukemia Research</i> , <b>2021</b> , 110, 106664	2.7	4

11	Association of red blood cell distribution width and outcomes in patients with mantle cell lymphoma. <i>Cancer Medicine</i> , <b>2019</b> , 8, 2751-2758	4.8	3
10	The prognostic value of 25-hydroxy vitamin D deficiency and its interaction with c-Myc expression in diffuse large B cell lymphoma. <i>Annals of Hematology</i> , <b>2020</b> , 99, 2377-2384	3	3
9	Circulating low CD4/CD8 ratio is associated with poor prognosis in Waldenstrom macroglobulinemia patients. <i>Annals of Hematology</i> , <b>2021</b> , 100, 995-1002	3	3
8	A new SH2D1A mutation in a female adult XLP disease with hemophagocytic lymphohistiocytosis and NK-cell leukemia. <i>Annals of Hematology</i> , <b>2019</b> , 98, 2829-2831	3	3
7	The prognostic role of HBV infection in chronic lymphocytic leukemia. <i>Journal of Cancer Research and Clinical Oncology</i> , <b>2018</b> , 144, 1309-1315	4.9	3
6	98% IGHV gene identity is the optimal cutoff to dichotomize the prognosis of Chinese patients with chronic lymphocytic leukemia. <i>Cancer Medicine</i> , <b>2020</b> , 9, 999-1007	4.8	2
5	Low serum cholesterol levels predict inferior prognosis and improve prognostic index scoring for peripheral T-cell lymphoma, unspecified. <i>Leukemia Research</i> , <b>2021</b> , 103, 106534	2.7	2
4	25-Hydroxy vitamin D deficiency predicts inferior prognosis in Hodgkin lymphoma. <i>Leukemia Research</i> , <b>2021</b> , 105, 106580	2.7	2
3	Pretreatment whole blood Epstein-Barr virus DNA predicts prognosis in Hodgkin lymphoma. <i>Leukemia Research</i> , <b>2021</b> , 107, 106607	2.7	2
2	Definition of disease-progression risk stratification in untreated chronic lymphocytic leukemia using combined clinical, molecular and virological variables. <i>Hematological Oncology</i> , <b>2018</b> , 36, 656-662	1.3	1
1	Interim response in diffuse large B cell lymphoma on CT: what is the optimal size reduction (SPD) for predicting outcome?. <i>European Radiology</i> , <b>2020</b> , 30, 3094-3100	8	