

Tianxin Yang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9196627/publications.pdf>

Version: 2024-02-01

50
papers

1,767
citations

304743

22
h-index

276875

41
g-index

50
all docs

50
docs citations

50
times ranked

1567
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Activating P2Y1 receptors improves function in arteries with repressed autophagy. Cardiovascular Research, 2023, 119, 252-267. | 3.8 | 10 |
| 2 | Role of (pro)renin receptor in cyclosporin A-induced nephropathy. American Journal of Physiology - Renal Physiology, 2022, 322, F437-F448. | 2.7 | 5 |
| 3 | Na ⁺ -Retaining Action of COX-2 (Cyclooxygenase-2)/EP ₁ Pathway in the Collecting Duct via Activation of Intrarenal Renin-Angiotensin-Aldosterone System and Epithelial Sodium Channel. Hypertension, 2022, 79, 1190-1202. | 2.7 | 7 |
| 4 | Revisiting the relationship between (Pro)Renin receptor and the intrarenal RAS: focus on the soluble receptor. Current Opinion in Nephrology and Hypertension, 2022, 31, 351-357. | 2.0 | 2 |
| 5 | Site-1 Protease-Derived Soluble (Pro)Renin Receptor Contributes to Angiotensin II-Induced Hypertension in Mice. Hypertension, 2021, 77, 405-416. | 2.7 | 25 |
| 6 | Soluble (pro)renin receptor induces endothelial dysfunction and hypertension in mice with diet-induced obesity via activation of angiotensin II type 1 receptor. Clinical Science, 2021, 135, 793-810. | 4.3 | 24 |
| 7 | Serine Protease HTRA1 as a Novel Target Antigen in Primary Membranous Nephropathy. Journal of the American Society of Nephrology: JASN, 2021, 32, 1666-1681. | 6.1 | 61 |
| 8 | (Pro)renin receptor antagonist PRO20 attenuates nephrectomy-induced nephropathy in rats via inhibition of intrarenal RAS and Wnt/β-catenin signaling. Physiological Reports, 2021, 9, e14881. | 1.7 | 10 |
| 9 | Targeting AT ₂ receptors in renal disease. American Journal of Physiology - Renal Physiology, 2021, 320, F1025-F1027. | 2.7 | 2 |
| 10 | Mutagenesis of the Cleavage Site of Pro Renin Receptor Abrogates Angiotensin II-Induced Hypertension in Mice. Hypertension, 2021, 78, 115-127. | 2.7 | 13 |
| 11 | Soluble (Pro)Renin Receptor as a Negative Regulator of NCC (Na ⁺ -Cl ⁻) Cotransporter. JCI Insight, 2020, 5, . | 2.7 | 8 |
| 12 | Soluble (pro)renin receptor regulation of ENaC involved in aldosterone signaling in cultured collecting duct cells. American Journal of Physiology - Renal Physiology, 2020, 318, F817-F825. | 2.7 | 18 |
| 13 | (Pro)renin receptor decoy peptide PRO20 protects against adriamycin-induced nephropathy by targeting the intrarenal renin-angiotensin system. American Journal of Physiology - Renal Physiology, 2020, 319, F930-F940. | 2.7 | 15 |
| 14 | Soluble (pro)renin receptor promotes the fibrotic response in renal proximal tubule epithelial cells in vitro via the Akt/β-catenin/Snail signaling pathway. American Journal of Physiology - Renal Physiology, 2020, 319, F941-F953. | 2.7 | 15 |
| 15 | Soluble (pro)renin receptor treats metabolic syndrome in mice with diet-induced obesity via interaction with PPAR ^γ . JCI Insight, 2020, 5, . | 5.0 | 20 |
| 16 | Hydrogen sulfide upregulates renal AQP2 protein expression and promotes urine concentration. FASEB Journal, 2019, 33, 469-483. | 0.5 | 32 |
| 17 | Site-1 protease-derived soluble (pro)renin receptor targets vasopressin receptor 2 to enhance urine concentrating capability. JCI Insight, 2019, 4, . | 5.0 | 24 |
| 18 | (Pro)renin receptor contributes to pregnancy-induced sodium-water retention in rats via activation of intrarenal RAAS and ENaC. American Journal of Physiology - Renal Physiology, 2019, 316, F530-F538. | 2.7 | 14 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 19 | FGF21 Prevents Angiotensin II-Induced Hypertension and Vascular Dysfunction by Activation of ACE2/Angiotensin-(1 α -7) Axis in Mice. <i>Cell Metabolism</i> , 2018, 27, 1323-1337.e5. | 16.2 | 104 |
| 20 | Enzymatic sources and physio-pathological functions of soluble (pro)renin receptor. <i>Current Opinion in Nephrology and Hypertension</i> , 2018, 27, 77-82. | 2.0 | 16 |
| 21 | Soluble (pro)renin receptor as a potential therapy for diabetes insipidus. <i>American Journal of Physiology - Renal Physiology</i> , 2018, 315, F1416-F1421. | 2.7 | 2 |
| 22 | Role of (pro)renin receptor in albumin overload-induced nephropathy in rats. <i>American Journal of Physiology - Renal Physiology</i> , 2018, 315, F1759-F1768. | 2.7 | 27 |
| 23 | Sodium Butyrate Attenuates Angiotensin II-Induced Cardiac Hypertrophy by Inhibiting COX2/PGE2 Pathway via a HDAC5/HDAC6-Dependent Mechanism. <i>FASEB Journal</i> , 2018, 32, 580.6. | 0.5 | 0 |
| 24 | Overexpression of ELABELA in the Renal Medulla Attenuated DOCA/salt-Induced Hypertension in Sprague Dawley rats. <i>FASEB Journal</i> , 2018, 32, 716.6. | 0.5 | 0 |
| 25 | Deficiency of mPGES-1 exacerbates renal fibrosis and inflammation in mice with unilateral ureteral obstruction. <i>American Journal of Physiology - Renal Physiology</i> , 2017, 312, F121-F133. | 2.7 | 14 |
| 26 | Physiology and Pathophysiology of the Intrarenal Renin-Angiotensin System: An Update. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 1040-1049. | 6.1 | 176 |
| 27 | Collecting duct (pro)renin receptor targets ENaC to mediate angiotensin II-induced hypertension. <i>American Journal of Physiology - Renal Physiology</i> , 2017, 312, F245-F253. | 2.7 | 59 |
| 28 | (Pro)Renin receptor regulates potassium homeostasis through a local mechanism. <i>American Journal of Physiology - Renal Physiology</i> , 2017, 313, F641-F656. | 2.7 | 15 |
| 29 | Activation of Renal (Pro)Renin Receptor Contributes to High Fructose-Induced Salt Sensitivity. <i>Hypertension</i> , 2017, 69, 339-348. | 2.7 | 66 |
| 30 | NF- κ B-dependent upregulation of (pro)renin receptor mediates high-NaCl-induced apoptosis in mouse inner medullary collecting duct cells. <i>American Journal of Physiology - Cell Physiology</i> , 2017, 313, C612-C620. | 4.6 | 12 |
| 31 | (Pro)renin receptor mediates albumin-induced cellular responses: role of site-1 protease-derived soluble (pro)renin receptor in renal epithelial cells. <i>American Journal of Physiology - Cell Physiology</i> , 2017, 313, C632-C643. | 4.6 | 35 |
| 32 | The soluble (Pro) renin receptor does not influence lithium-induced diabetes insipidus but does provoke beiging of white adipose tissue in mice. <i>Physiological Reports</i> , 2017, 5, e13410. | 1.7 | 14 |
| 33 | A H ₂ S Donor GYY4137 Exacerbates Cisplatin-Induced Nephrotoxicity in Mice. <i>Mediators of Inflammation</i> , 2016, 2016, 1-10. | 3.0 | 29 |
| 34 | Activation of ENaC in collecting duct cells by prorenin and its receptor PRR: involvement of Nox4-derived hydrogen peroxide. <i>American Journal of Physiology - Renal Physiology</i> , 2016, 310, F1243-F1250. | 2.7 | 67 |
| 35 | High potassium promotes mutual interaction between (pro)renin receptor and the local renin-angiotensin-aldosterone system in rat inner medullary collecting duct cells. <i>American Journal of Physiology - Cell Physiology</i> , 2016, 311, C686-C695. | 4.6 | 12 |
| 36 | Protection of nitro-fatty acid against kidney diseases. <i>American Journal of Physiology - Renal Physiology</i> , 2016, 310, F697-F704. | 2.7 | 20 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Soluble (pro)renin receptor via β -catenin enhances urine concentration capability as a target of liver X receptor. Proceedings of the National Academy of Sciences of the United States of America, 2016, 113, E1898-906. | 7.1 | 83 |
| 38 | Antidiuretic Action of Collecting Duct (Pro)Renin Receptor Downstream of Vasopressin and PGE2 Receptor EP4. Journal of the American Society of Nephrology: JASN, 2016, 27, 3022-3034. | 6.1 | 67 |
| 39 | Intrarenal renin-angiotensin system mediates fatty acid-induced ER stress in the kidney. American Journal of Physiology - Renal Physiology, 2016, 310, F351-F363. | 2.7 | 54 |
| 40 | Renal medullary (pro)renin receptor contributes to angiotensin II-induced hypertension in rats via activation of the local renin-angiotensin system. BMC Medicine, 2015, 13, 278. | 5.5 | 63 |
| 41 | Aliskiren restores renal AQP2 expression during unilateral ureteral obstruction by inhibiting the inflammasome. American Journal of Physiology - Renal Physiology, 2015, 308, F910-F922. | 2.7 | 42 |
| 42 | Crosstalk between (Pro)renin receptor and COX-2 in the renal medulla during angiotensin II-induced hypertension. Current Opinion in Pharmacology, 2015, 21, 89-94. | 3.5 | 26 |
| 43 | COX-2 mediates angiotensin II-induced (pro)renin receptor expression in the rat renal medulla. American Journal of Physiology - Renal Physiology, 2014, 307, F25-F32. | 2.7 | 51 |
| 44 | Prostaglandin E-Prostanoid EP_4 Receptor Mediates Angiotensin II-Induced (Pro)Renin Receptor Expression in the Rat Renal Medulla. Hypertension, 2014, 64, 369-377. | 2.7 | 64 |
| 45 | The tempo of cardiovascular and metabolic responses to fasting is different between lean and obese mice. FASEB Journal, 2010, 24, 978.15. | 0.5 | 0 |
| 46 | Hydrogen peroxide stimulates chloride secretion in primary inner medullary collecting duct cells via mPGES-1-derived PGE_2 . American Journal of Physiology - Renal Physiology, 2007, 293, F1571-F1576. | 2.7 | 26 |
| 47 | Kidney-specific gene targeting: Insight into thiazolidinedione-induced fluid retention (Review Article). Nephrology, 2006, 11, 201-206. | 1.6 | 8 |
| 48 | Expression and function of COX isoforms in renal medulla: evidence for regulation of salt sensitivity and blood pressure. American Journal of Physiology - Renal Physiology, 2006, 290, F542-F549. | 2.7 | 69 |
| 49 | Renin expression in COX-2-knockout mice on normal or low-salt diets. American Journal of Physiology - Renal Physiology, 2000, 279, F819-F825. | 2.7 | 98 |
| 50 | Low Chloride Stimulation of Prostaglandin E_2 Release and Cyclooxygenase-2 Expression in a Mouse Macula Densa Cell Line. Journal of Biological Chemistry, 2000, 275, 37922-37929. | 3.4 | 145 |