

Gracia LiÃ©banas

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9198465/publications.pdf>

Version: 2024-02-01

39

papers

655

citations

623734

14

h-index

580821

25

g-index

39

all docs

39

docs citations

39

times ranked

464

citing authors

#	ARTICLE	IF	CITATIONS
1	Pratylenchus Ávovlasi sp. Nov. (Nematoda: Pratylenchidae) on Raspberries in North Italy with a Morphometrical and Molecular Characterization. <i>Plants</i> , 2021, 10, 1068.	3.5	3
2	The morphology of <i>Colpocephalum pectinatum</i> (Phthiraptera: Amblycera: Menoponidae) under scanning electron microscopy. <i>Arthropod Structure and Development</i> , 2021, 64, 101085.	1.4	2
3	Diversity of root-knot nematodes of the genus <i>Meloidogyne</i> Gómez, 1892 (Nematoda: Meloidogynidae) associated with olive plants and environmental cues regarding their distribution in southern Spain. <i>PLoS ONE</i> , 2018, 13, e0198236.	2.5	33
4	Root-lesion nematodes of the genus <i>Pratylenchus</i> (Nematoda: Pratylenchidae) from Costa Rica with molecular identification of <i>P. gutierrezi</i> and <i>P. panamaensis</i> topotypes. <i>European Journal of Plant Pathology</i> , 2016, 145, 973-998.	1.7	17
5	A new stem nematode, <i>Ditylenchus oncogenus</i> n. sp. (Nematoda: Tylenchida), parasitizing sowthistle from Adriatic coast dunes in southern Italy. <i>Journal of Helminthology</i> , 2016, 90, 152-165.	1.0	27
6	<i>Rotylenchus cretensis</i> n. sp. and <i>R. cypriensis</i> Antoniou 1980 (Nematoda: Hoplolaimidae) recovered from the rhizosphere of olive at Crete (Greece) with a molecular phylogeny of the genus. <i>European Journal of Plant Pathology</i> , 2016, 144, 167-184.	1.7	5
7	< i> <i>Discopersicus</i> </i> n. gen., a New Member of the Family Tylenchidae Orley, 1880 with Detailed SEM Study on Two Known Species of the Genus < i> <i>Discotylenchus</i> </i> Siddiqi, 1980 (Nematoda; Tylenchidae) from Iran. <i>Journal of Nematology</i> , 2016, 48, 214-221.	0.9	14
8	Parasitism effects on white clover by root-knot and cyst nematodes and molecular separation of <i>Heterodera daverti</i> from <i>H. trifolii</i> . <i>European Journal of Plant Pathology</i> , 2015, 143, 833-845.	1.7	19
9	<i>Rotylenchus castilloi</i> n. sp. (Nematoda: Hoplolaimidae), a new species with long stylet from northern Iran. <i>Zootaxa</i> , 2015, 3931, 88-100.	0.5	10
10	A new root-knot nematode <i>Meloidogyne spartelensis</i> n. sp. (Nematoda: Meloidogynidae) in Northern Morocco. <i>European Journal of Plant Pathology</i> , 2015, 143, 25-42.	1.7	16
11	Description of <i>Rotylenchus urmiaensis</i> n. sp. (Nematoda: Hoplolaimidae) from North-western Iran with molecular phylogeny of the genus. <i>Nematology</i> , 2015, 17, 607-619.	0.6	7
12	Characterisation of an Iberian population of <i>Rhyssocolpus iuventutis</i> Andrassy, 1971 (Dorylaimida: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 147 Td (Pr)	0.6	
13	Description of <i>Rotylenchus arasbaranensis</i> n. sp. from Iran with discussion on the taxonomic status of <i>Plesiotylenchus</i> Vovlas, Castillo & Lamberti, 1993 (Nematoda: Hoplolaimidae). <i>Nematology</i> , 2014, 16, 1019-1045.	0.6	10
14	Integrative taxonomy of the stunt nematodes of the genera < i> <i>Bitylenchus</i> </i> and < i> <i>Tylenchorhynchus</i> </i> (Nematoda, Telotylenchidae) with description of two new species and a molecular phylogeny. <i>Zoological Journal of the Linnean Society</i> , 2014, 172, 231-264.	2.3	36
15	Taxonomy of the genus <i>Oriverutus</i> Siddiqi, 1971 (Nematoda: Dorylaimida: Nordiidae). <i>Journal of Natural History</i> , 2014, 48, 1253-1278.	0.5	0
16	Morphological and molecular characterisation of <i>Pratylenchus oleae</i> n. sp. (Nematoda: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 147 Td (Pr) Pathology, 2014, 140, 53-67.	1.7	27
17	Comparative molecular and morphological characterisations in the nematode genus <i>Rotylenchus</i> : <i>Rotylenchus paravitis</i> n. sp., an example of cryptic speciation. <i>Zoologischer Anzeiger</i> , 2013, 252, 246-268.	0.9	43
18	Morphological and molecular characterisation of <i>Paralongidorus plesioepimikis</i> n. sp. (Nematoda: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50	0.6	15

#	ARTICLE	IF	CITATIONS
19	Molecular and morphological characterisation of <i>Paralongidorus iranicus</i> n. sp. and <i>P. bikanerensis</i> (Lal & Mathur, 1987) Siddiqi, Baujard & Mounport, 1993 (Nematoda: Longidoridae) from Iran. <i>Nematology</i> , 2012, 14, 427-443.	0.6	18
20	Molecular and morphological characterisation of <i>Rotylenchus vitis</i> n. sp. (Nematoda: Hoplolaimidae) infecting grapevine in southern Spain. <i>Nematology</i> , 2012, 14, 235-247.	0.6	13
21	< i> <i>Ditylenchus gigas</i> n. sp. parasitizing broad bean: a new stem nematode singled out from the < i> <i>Ditylenchus dipsaci</i> species complex using a polyphasic approach with molecular phylogeny. <i>Plant Pathology</i> , 2011, 60, 762-775.	2.4	77
22	Free-living dorylaimid nematodes from nature reserves in Costa Rica. The genus <i>Oriverutus</i> Siddiqi, 1971 “ species with specialised uterus. <i>Nematology</i> , 2011, 13, 307-318.	0.6	1
23	Free-living dorylaimid nematodes from nature reserves in Costa Rica. The genus <i>Oriverutus</i> Siddiqi, 1971 “ species with simple uterus. <i>Nematology</i> , 2011, 13, 463-475.	0.6	2
24	Molecular and morphological characterisation of <i>Sphaeronema alni</i> Turkina & Chizhov, 1986 (Nematoda: Sphaeronematidae) from Spain compared with a topotype population from Russia. <i>Nematology</i> , 2010, 12, 649-659.	0.6	7
25	Description of <i>Pratylenchus hispaniensis</i> n. sp. from Spain and considerations on the phylogenetic relationship among selected genera in the family Pratylenchidae. <i>Nematology</i> , 2010, 12, 429-451.	0.6	25
26	<i>Eutylenchus excretorius</i> Ebsary & Eveleigh, 1981 (Nematoda: Tylodoridae) from Spain with approaches to molecular phylogeny of related genera. <i>Nematology</i> , 2009, 11, 343-354.	0.6	13
27	A new root-knot nematode, < i> <i>Meloidogyne silvestris</i> n. sp. (Nematoda: Meloidogynidae), parasitizing European holly in northern Spain. <i>Plant Pathology</i> , 2009, 58, 606-619.	2.4	12
28	Soil enzymes, nematode community and selected physico-chemical properties as soil quality indicators in organic and conventional olive oil farming: Influence of seasonality and site features. <i>Applied Soil Ecology</i> , 2009, 41, 305-314.	4.3	111
29	Molecular phylogeny of the genus < i> <i>Rotylenchus</i> (Nematoda, Tylenchida) and description of a new species. <i>Zoologica Scripta</i> , 2008, 37, 521-537.	1.7	47
30	Nematodes of the order Dorylaimida from Andalucía Oriental, Spain. The genus <i>Enchodelus</i> Thorne, 1939. 2. Description of three known species with rounded tail and long odontostyle. <i>Nematology</i> , 2008, 10, 451-470.	0.6	2
31	Nematodes of the order Dorylaimida from Andalucía Oriental, Spain. The genus <i>Enchodelus</i> Thorne, 1939. 3. Description of two new and one known species with rounded tail and medium-sized odontostyle. <i>Nematology</i> , 2008, 10, 711-733.	0.6	1
32	Morphological and molecular characterisation of a new awl nematode, <i>Dolichodorus mediterraneus</i> sp. n. (Nematoda: Dolichodoridae), from Spain. <i>Nematology</i> , 2007, 9, 189-199.	0.6	3
33	Nematodes of the order Dorylaimida from Andalucía Oriental, Spain. The genus <i>Enchodelus</i> Thorne, 1939. 1. Description of two new and two known species with conical tail. <i>Nematology</i> , 2007, 9, 515-536.	0.6	4
34	A New Root-Knot Nematode Parasitizing Sea Rocket from Spanish Mediterranean Coastal Dunes: <i>Meloidogyne dunensis</i> n. sp. (Nematoda: Meloidogynidae). <i>Journal of Nematology</i> , 2007, 39, 190-202.	0.9	9
35	Re-establishment of <i>Hemicriconemoides promissus</i> (Nematoda: Criconematoidea) as a valid species, with additional data for <i>H. ortonwilliamsi</i> from Spain and <i>H. wessoni</i> from Florida. <i>Nematology</i> , 2006, 8, 511-519.	0.6	2
36	Host suitability of some crucifers for root-knot nematodes in southern Spain. <i>Nematology</i> , 2004, 6, 125-128.	0.6	12

#	ARTICLE	IF	CITATIONS
37	SEM studies on the Mediterranean olive root-knot nematode, <i>Meloidogyne baetica</i> , and histopathology on two additional natural hosts. <i>Nematology</i> , 2004, 6, 749-754.	0.6	7
38	Nematodes of the order Dorylaimida from Andalucía Oriental, Spain. New observations of <i>Chitwoodiellus parafuscus</i> , with comments on the genus <i>Chitwoodiellus</i> Jiménez-Guirado & Peña-Santiago, 1992. <i>Nematology</i> , 2002, 4, 371-380.	0.6	0
39	Nematodes of the order Dorylaimida from Andalucía Oriental, Spain: <i>Discolaimus Cobb</i> , 1913. 1. Description of two new species. <i>Nematology</i> , 2002, 4, 361-369.	0.6	0