
Junko Yamagishi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/920094/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Varietal difference in dynamics of non-structural carbohydrates in nodal segments of stem in two varieties of rice (<i>Oryza sativa</i> L.) at pre- and post-heading stages. Plant Production Science, 2022, 25, 30-42.	2.0	5
2	Detection and characterization of quantitative trait loci for coleoptile elongation under anaerobic conditions in rice. Plant Production Science, 2020, 23, 374-383.	2.0	10
3	Environmental Compensation Effect and Synergistic Mechanism of Optimized Nitrogen Management Increasing Nitrogen Use Efficiency in Indica Hybrid Rice. Frontiers in Plant Science, 2019, 10, 245.	3.6	13
4	Characterisation of a rice vacuolar invertase isoform, OsINV2, for growth and yield-related traits. Functional Plant Biology, 2019, 46, 777.	2.1	5
5	Evaluation of performance of sorghum varieties grown in Tokyo for sugar accumulation and its correlation with vacuolar invertase genes SbInv1 and SbInv2. Plant Production Science, 2018, 21, 328-338.	2.0	6
6	Genetic Evidence for the Role of a Rice Vacuolar Invertase as a Molecular Sink Strength Determinant. Rice, 2018, 11, 6.	4.0	46
7	Mild drying of sandy soil can physically limit the uptake of phosphorus by rainfed lowland rice in northeast Thailand. Soil Science and Plant Nutrition, 2018, 64, 677-685.	1.9	1
8	Residual effects of cultivation methods on weed seed banks and weeds in Cambodia. Weed Biology and Management, 2016, 16, 93-107.	1.4	6
9	Grain yield and phosphorus uptake of rainfed lowland rice under unsubmerged soil stress. Field Crops Research, 2016, 190, 54-59.	5.1	24
10	Morphological traits associated with vegetative growth of rice (Oryza sativa L.) during the recovery phase after early-season drought. European Journal of Agronomy, 2015, 64, 58-66.	4.1	25
11	Agronomic performance of an IR64 introgression line with large leaves derived from New Plant Type rice in aerobic culture. European Journal of Agronomy, 2014, 58, 11-17.	4.1	9
12	Root growth response of rainfed lowland rice to aerobic conditions in northeastern Thailand. Plant and Soil, 2013, 368, 557-567.	3.7	16
13	ã€Short Report】Grain Yield and Leaf Area Growth of Direct-Seeded Rice on Flooded and Aerobic Soils in Japan. Plant Production Science, 2013, 16, 276-279.	2.0	13
14	Allometric relationship between the size and number of shoots as a determinant of adaptations in rice to water-saving aerobic culture. Field Crops Research, 2012, 131, 17-25.	5.1	12
15	Long-term effects of organic manure application on the productivity of winter wheat grown in a crop rotation with maize in Japan. Field Crops Research, 2011, 120, 387-395.	5.1	24
16	Role of early vigor in adaptation of rice to water-saving aerobic culture: Effects of nitrogen utilization and leaf growth. Field Crops Research, 2011, 124, 124-131.	5.1	24
17	Ecophysiological study on weed seed banks and weeds in Cambodian paddy fields with contrasting water availability. Weed Biology and Management, 2010, 10, 261-272.	1.4	33
18	High-yielding Crop Management by Enhancing Growth in Reproductive Stage of Direct-Seeded Rainfed Lowland Rice (Oryza sativaL.) in Northeast Thailand. Plant Production Science, 2010, 13, 104-115.	2.0	5

Junko Yamagishi

#	Article	IF	CITATIONS
19	Response of Spikelet Number per Panicle in Rice Cultivars to Three Transplanting Densities. Plant Production Science, 2010, 13, 279-288.	2.0	10
20	Spatial variability in the growth of direct-seeded rainfed lowland rice (Oryza sativa L.) in northeast Thailand. Field Crops Research, 2009, 111, 251-261.	5.1	13
21	QTL analysis of panicle morphology response to irrigation regime in aerobic rice culture. Field Crops Research, 2009, 114, 295-303.	5.1	6
22	Identification of QTLs controlling rice drought tolerance at seedling stage in hydroponic culture. Euphytica, 2008, 160, 423-430.	1.2	73
23	Assessment of management of direct seeded rice production under different water conditions in Cambodia. Paddy and Water Environment, 2008, 6, 91-103.	1.8	27
24	Preflowering Abortion Reduces Spikelet Number in Upland Rice (<i>Oryza sativa</i> L.) under Water Stress. Crop Science, 2008, 48, 2389-2395.	1.8	87
25	Varietal Differences in Stem Diameter and Rooting Number of Phytomers in Conjunction with Root System Development of Field-Grown Rice (<i>Oryza sativa</i> L.). Plant Production Science, 2007, 10, 357-360.	2.0	6
26	Growth of Rice (Oryza SativaL.) Cultivars Under Upland Conditions With Different Levels of Water Supply3. Root System Development, Soil Moisture Changeand Plant Water Status. Plant Production Science, 2007, 10, 3-13.	2.0	46
27	Genotypic differences in grain yield of transplanted and direct-seeded rainfed lowland rice (Oryza) Tj ETQq1 1 C).784314 rg	gBT ₃ 90verlock
28	Evaluating the resistance of six rice cultivars to drought: restriction of deep rooting and the use of raised beds. Plant and Soil, 2007, 300, 149-161.	3.7	38
29	Visceral coverage with absorbable mesh followed by split-thickness skin graft in the treatment of ruptured giant omphalocele. Pediatric Surgery International, 2007, 23, 199-201.	1.4	19
30	Improvement of rice (Oryza sativa L.) growth in upland conditions with deep tillage and mulch. Soil and Tillage Research, 2007, 92, 30-44.	5.6	22
31	Genotypic Variation in Root Growth Angle in Rice (Oryza sativa L.) and its Association with Deep Root Development in Upland Fields with Different Water Regimes. Plant and Soil, 2006, 287, 117-129.	3.7	202
32	Effect of Planting Density on Grain Yield and Water Productivity of Rice (Oryza sativaL.) Grown in Flooded and Non-flooded Fields in Japan. Plant Production Science, 2006, 9, 298-311.	2.0	37
33	Growth of Three Rice (Oryza sativaL.) Cultivars under Upland Conditions with Different Levels of Water Supply. Plant Production Science, 2006, 9, 422-434.	2.0	38
34	Growth of Three Rice Cultivars (Oryza sativaL.) under Upland Conditions with Different Levels of Water Supply. Plant Production Science, 2006, 9, 435-445.	2.0	35
35	Size and Activity of Shoot Apical Meristems as Determinants of Floret Number in Rice Panicles. Plant Production Science, 2005, 8, 51-59.	2.0	19
36	QTLs for branching, floret formation, and pre-flowering floret abortion of rice panicle in a temperate japonica�×tropical japonica cross. Theoretical and Applied Genetics, 2004, 109, 1555-1561.	3.6	61

3

Junko Yamagishi

#	Article	IF	CITATIONS
37	Effects of Tillage on Along-Row Variability of Wheat and Maize Biomass. Plant Production Science, 2003, 6, 295-301.	2.0	1
38	Diversity of the Rachis-Branching System in a Panicle in Japónica Rice. Plant Production Science, 2003, 6, 59-64.	2.0	9
39	Spatial Variability Patterns of Wheat Growth and Soil Properties in a Small Field as Affected by Tillage Intensity. Plant Production Science, 2002, 5, 175-183.	2.0	12
40	Effects of Gibberellic Acid Application on Panicle Characteristics and Size of Shoot Apex in the First Bract Differentiation Stage in Rice. Plant Production Science, 2001, 4, 227-229.	2.0	6
41	Extent and Implications of Weed Spatial Variability in Arable Crop Fields. Plant Production Science, 2001, 4, 259-269.	2.0	10
42	The Spatial Variability Patterns of Maize Growth and Root Colonization by Arbuscular Mycorrhizal Fungi in aSmall Field. Plant Production Science, 2001, 4, 249-254.	2.0	6
43	The Maturing Processes of Field Reclaimed with Sub-surface Soil. The change of matter production during 17 years Japanese Journal of Crop Science, 1998, 67, 302-306.	0.2	1
44	Relation of Number of Spikelets per Panicle to the Characteristics of Shoot and the Size around Growing Point at Panicle Initiation Stage in Rice Varieties Japanese Journal of Crop Science, 1992, 61, 568-575.	0.2	11