

Hisao Naito

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9205988/hisao-naito-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

56
papers

1,075
citations

20
h-index

31
g-index

59
ext. papers

1,241
ext. citations

3.7
avg, IF

3.5
L-index

#	Paper	IF	Citations
56	Occupational exposure limits for acetaldehyde, 2-bromopropane, glyphosate, manganese and inorganic manganese compounds, and zinc oxide nanoparticle, and the biological exposure indices for cadmium and cadmium compounds and ethylbenzene, and carcinogenicity, occupational sensitizer, and reproductive toxicant classifications. <i>Journal of Occupational Health</i> , 2021 , 63, e12294	2.3	1
55	One-Pot Extraction and Quantification Method for Bile Acids in the Rat Liver by Capillary Liquid Chromatography Tandem Mass Spectrometry. <i>ACS Omega</i> , 2021 , 6, 8588-8597	3.9	0
54	The antihypertensive agent hydralazine reduced extracellular matrix synthesis and liver fibrosis in nonalcoholic steatohepatitis exacerbated by hypertension. <i>PLoS ONE</i> , 2020 , 15, e0243846	3.7	0
53	Increased risk of occupational trichloroethylene hypersensitivity syndrome at exposure levels higher than 15µg/L of urinary trichloroacetic acid, regardless of whether the patients had the HLA-B*13:01 allele. <i>Environmental Research</i> , 2020 , 191, 109972	7.9	2
52	Trichloroethylene and trichloroethanol induce skin sensitization with focal hepatic necrosis in guinea pigs. <i>Journal of Occupational Health</i> , 2020 , 62, e12142	2.3	4
51	High-fat and high-cholesterol diet decreases phosphorylated inositol-requiring kinase-1 and inhibits autophagy process in rat liver. <i>Scientific Reports</i> , 2019 , 9, 12514	4.9	5
50	In utero exposure to di(2-ethylhexyl)phthalate suppresses blood glucose and leptin levels in the offspring of wild-type mice. <i>Toxicology</i> , 2019 , 415, 49-55	4.4	8
49	Validation of the Japanese Version of the Yale Food Addiction Scale 2.0 (J-YFAS 2.0). <i>Nutrients</i> , 2019 , 11,	6.7	19
48	Occupational exposure limits for cumene, 2,4-dichlorophenoxy acetic acid, silicon carbide whisker, benzyl alcohol, and methylamine, and carcinogenicity, occupational sensitizer, and reproductive toxicant classifications. <i>Journal of Occupational Health</i> , 2019 , 61, 328-330	2.3	2
47	Smoking results in accumulation of ectopic fat in the liver. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2019 , 12, 1075-1080	3.4	6
46	Effect of laughter yoga on salivary cortisol and dehydroepiandrosterone among healthy university students: A randomized controlled trial. <i>Complementary Therapies in Clinical Practice</i> , 2018 , 32, 6-11	3.5	14
45	Occupational Exposure Limits for ethylidene norbornene, ethyleneimine, benomyl, and 2,3-epoxypropyl methacrylate, and classifications on carcinogenicity. <i>Journal of Occupational Health</i> , 2018 , 60, 333-335	2.3	1
44	Bile acid detoxifying enzymes limit susceptibility to liver fibrosis in female SHRSP5/Dmcr rats fed with a high-fat-cholesterol diet. <i>PLoS ONE</i> , 2018 , 13, e0192863	3.7	11
43	Occupational exposure limits for ethylene glycol monobutyl ether, isoprene, isopropyl acetate and propyleneimine, and classifications on carcinogenicity, occupational sensitizer and reproductive toxicant. <i>Journal of Occupational Health</i> , 2017 , 59, 364-366	2.3	1
42	Combination of Hypertension Along with a High Fat and Cholesterol Diet Induces Severe Hepatic Inflammation in Rats via a Signaling Network Comprising NF- κ B, MAPK, and Nrf2 Pathways. <i>Nutrients</i> , 2017 , 9,	6.7	14
41	Response to the letter to the editor: Metallothionein MT2A A-5G polymorphism and the risk for chronic kidney disease and diabetes. <i>Toxicological Sciences</i> , 2016 , 154, 198-199	4.4	
40	Efficacy of Dietary Lipid Control in Healing High-Fat and High-Cholesterol Diet-Induced Fibrotic Steatohepatitis in Rats. <i>PLoS ONE</i> , 2016 , 11, e0145939	3.7	8

39	Importance of detoxifying enzymes in differentiating fibrotic development between SHRSP5/Dmcr and SHRSP rats. <i>Environmental Health and Preventive Medicine</i> , 2016 , 21, 368-381	4.2	8
38	Metallothionein MT2A A-5G Polymorphism as a Risk Factor for Chronic Kidney Disease and Diabetes: Cross-Sectional and Cohort Studies. <i>Toxicological Sciences</i> , 2016 , 152, 181-93	4.4	18
37	Prenatal maternal blood triglyceride and fatty acid levels in relation to exposure to di(2-ethylhexyl)phthalate: a cross-sectional study. <i>Environmental Health and Preventive Medicine</i> , 2015 , 20, 168-78	4.2	26
36	Arsenite-mediated promotion of anchorage-independent growth of HaCaT cells through placental growth factor. <i>Journal of Investigative Dermatology</i> , 2015 , 135, 1147-1156	4.3	18
35	Association of maternal whole blood fatty acid status during the prenatal period with term birth dimensions: a cross-sectional study. <i>Journal of Perinatal Medicine</i> , 2015 , 43, 565-75	2.7	5
34	The Progression of Non-alcoholic Fatty Liver Disease and Lifestyle Intervention in Older Adults 2015 , 85-97		1
33	A trial to find appropriate animal models of dichloropropane-induced cholangiocarcinoma based on the hepatic distribution of glutathione S-transferases. <i>Journal of Occupational Health</i> , 2015 , 57, 548-54	2.3	4
32	Commentary to Krishna et al. (2014): brain deposition and neurotoxicity of manganese in adult mice exposed via the drinking water. <i>Archives of Toxicology</i> , 2014 , 88, 1185-6	5.8	6
31	Species and inter-individual differences in metabolic capacity of di(2-ethylhexyl)phthalate (DEHP) between human and mouse livers. <i>Environmental Health and Preventive Medicine</i> , 2014 , 19, 117-25	4.2	31
30	Association between maternal exposure to di(2-ethylhexyl) phthalate and reproductive hormone levels in fetal blood: the Hokkaido study on environment and children's health. <i>PLoS ONE</i> , 2014 , 9, e109039	3.7	87
29	A possible role of chenodeoxycholic acid and glycine-conjugated bile acids in fibrotic steatohepatitis in a dietary rat model. <i>Digestive Diseases and Sciences</i> , 2014 , 59, 1490-501	4	17
28	High-fat-cholesterol diet mainly induced necrosis in fibrotic steatohepatitis rat by suppressing caspase activity. <i>Life Sciences</i> , 2013 , 93, 673-80	6.8	16
27	Dysregulated bile acid synthesis, metabolism and excretion in a high fat-cholesterol diet-induced fibrotic steatohepatitis in rats. <i>Digestive Diseases and Sciences</i> , 2013 , 58, 2212-22	4	25
26	Exposure to DEHP decreased four fatty acid levels in plasma of prepartum mice. <i>Toxicology</i> , 2013 , 309, 52-60	4.4	21
25	Occupational trichloroethylene hypersensitivity syndrome: human herpesvirus 6 reactivation and rash phenotypes. <i>Journal of Dermatological Science</i> , 2013 , 72, 218-24	4.3	24
24	Effects of sub-acute and sub-chronic inhalation of 1-bromopropane on neurogenesis in adult rats. <i>Toxicology</i> , 2013 , 304, 76-82	4.4	8
23	Effects of exposure to 1-bromopropane on astrocytes and oligodendrocytes in rat brain. <i>Journal of Occupational Health</i> , 2013 , 55, 29-38	2.3	8
22	Comparison of Barium and Arsenic Concentrations in Well Drinking Water and in Human Body Samples and a Novel Remediation System for These Elements in Well Drinking Water. <i>PLoS ONE</i> , 2013 , 8, e66681	3.7	35

21	Sex differences in metabolism of trichloroethylene and trichloroethanol in guinea pigs. <i>Journal of Occupational Health</i> , 2013 , 55, 443-9	2.3	7
20	Simple and rapid quantitation of 21 bile acids in rat serum and liver by UPLC-MS-MS: effect of high fat diet on glycine conjugates of rat bile acids. <i>Nagoya Journal of Medical Science</i> , 2013 , 75, 57-71	0.7	30
19	Modulation of ammonium perfluorooctanoate-induced hepatic damage by genetically different PPAR α in mice. <i>Archives of Toxicology</i> , 2012 , 86, 63-74	5.8	21
18	Evidence for diazinon-mediated inhibition of cis-permethrin metabolism and its effects on reproductive toxicity in adult male mice. <i>Reproductive Toxicology</i> , 2012 , 34, 489-97	3.4	15
17	The modulation of hepatic adenosine triphosphate and inflammation by eicosapentaenoic acid during severe fibrotic progression in the SHRSP5/Dmcr rat model. <i>Life Sciences</i> , 2012 , 90, 934-43	6.8	18
16	Simultaneous changes in high-fat and high-cholesterol diet-induced steatohepatitis and severe fibrosis and those underlying molecular mechanisms in novel SHRSP5/Dmcr rat. <i>Environmental Health and Preventive Medicine</i> , 2012 , 17, 444-56	4.2	17
15	Development of novel rat model for high-fat and high-cholesterol diet-induced steatohepatitis and severe fibrosis progression in SHRSP5/Dmcr. <i>Environmental Health and Preventive Medicine</i> , 2012 , 17, 173-82	4.2	38
14	Differences in metabolite burden of di(2-ethylhexyl)phthalate in pregnant and postpartum dams and their offspring in relation to drug-metabolizing enzymes in mice. <i>Archives of Toxicology</i> , 2012 , 86, 563-9	5.8	12
13	Plasticizers May Activate Human Hepatic Peroxisome Proliferator-Activated Receptor α Less Than That of a Mouse but May Activate Constitutive Androstane Receptor in Liver. <i>PPAR Research</i> , 2012 , 2012, 201284	4.3	27
12	Incidence of young onset insulin-requiring diabetes mellitus among 18- to 30-year-olds in Dhaka, Bangladesh (1994-2003). <i>Nagoya Journal of Medical Science</i> , 2012 , 74, 149-56	0.7	2
11	Ammonium perfluorooctanoate may cause testosterone reduction by adversely affecting testis in relation to PPAR α . <i>Toxicology Letters</i> , 2011 , 205, 265-72	4.4	23
10	Hepatic peroxisome proliferator-activated receptor α may have an important role in the toxic effects of di(2-ethylhexyl)phthalate on offspring of mice. <i>Toxicology</i> , 2011 , 289, 1-10	4.4	33
9	Study profile on baseline survey of Daiko Study in the Japan Multi-Institutional Collaborative Cohort Study (J-MICC Study). <i>Nagoya Journal of Medical Science</i> , 2011 , 73, 187-95	0.7	10
8	Differential response to trichloroethylene-induced hepatosteatosis in wild-type and PPAR α -humanized mice. <i>Environmental Health Perspectives</i> , 2010 , 118, 1557-63	8.4	30
7	Bisphenol A may cause testosterone reduction by adversely affecting both testis and pituitary systems similar to estradiol. <i>Toxicology Letters</i> , 2010 , 194, 16-25	4.4	162
6	"Hypothesis of seven balances": molecular mechanisms behind alcoholic liver diseases and association with PPAR α . <i>Journal of Occupational Health</i> , 2009 , 51, 391-403	2.3	23
5	Microgram-order ammonium perfluorooctanoate may activate mouse peroxisome proliferator-activated receptor alpha, but not human PPAR α . <i>Toxicology</i> , 2009 , 265, 27-33	4.4	42
4	Biological monitoring of pyrethroid exposure of pest control workers in Japan. <i>Journal of Occupational Health</i> , 2007 , 49, 509-14	2.3	32

3	Differential effects of aging, drinking and exercise on serum cholesterol levels dependent on the PPARA-V227A polymorphism. <i>Journal of Occupational Health</i> , 2007 , 49, 353-62	2.3	12
2	8-Hydroxydeoxyguanosine levels in human leukocyte and urine according to exposure to organophosphorus pesticides and paraoxonase 1 genotype. <i>International Archives of Occupational and Environmental Health</i> , 2007 , 80, 217-27	3.2	38
1	Association of V227A PPARalpha polymorphism with altered serum biochemistry and alcohol drinking in Japanese men. <i>Pharmacogenetics and Genomics</i> , 2006 , 16, 569-77	1.9	24