Nagamani Bora

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9207410/publications.pdf

Version: 2024-02-01

840119 1199166 16 558 11 12 citations h-index g-index papers 19 19 19 767 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Characterization of Actinomycetes from Smear Ripened Cheeses – A Polyphasic Approach. , 2015, , 51-101.		1
2	Analyzing the Metagenome of Smear Cheese Flora Using Next Generation Sequencing Tools. , 2015, , 137-153.		1
3	Actinobacterial Diversity and Dynamics as Revealed by Denaturing Gradient Gel Electrophoresis. , 2015, , 103-136.		1
4	Biodiversity of the Surface Microbial Consortia from Limburger, Reblochon, Livarot, Tilsit, and Gubbeen Cheeses., 2014,, 219-250.		2
5	Transcriptomic analysis of Lactococcus chungangensis sp. nov. and its potential in cheese making. Journal of Dairy Science, 2014, 97, 7363-7372.	1.4	11
6	Biodiversity of the Surface Microbial Consortia from Limburger, Reblochon, Livarot, Tilsit, and Gubbeen Cheeses. Microbiology Spectrum, 2014, 2, CM-0010-2012.	1.2	45
7	The Implementation of a Design of Experiments Strategy to Increase Recombinant Protein Yields in Yeast (Review). Methods in Molecular Biology, 2012, 866, 115-127.	0.4	12
8	Large-Scale Production of Secreted Proteins in Pichia pastoris. Methods in Molecular Biology, 2012, 866, 217-235.	0.4	2
9	Antifoam addition to shake flask cultures of recombinant Pichia pastoris increases yield. Microbial Cell Factories, 2011, 10, 17.	1.9	37
10	Surface microbial consortia from Livarot, a French smear-ripened cheese. Canadian Journal of Microbiology, 2011, 57, 651-660.	0.8	76
11	Understanding the yeast host cell response to recombinant membrane protein production. Biochemical Society Transactions, 2011, 39, 719-723.	1.6	13
12	Altering the ribosomal subunit ratio in yeast maximizes recombinant protein yield. Microbial Cell Factories, 2009, 8, 10.	1.9	57
13	Mycetocola reblochoni sp. nov., isolated from the surface microbial flora of Reblochon cheese. International Journal of Systematic and Evolutionary Microbiology, 2008, 58, 2687-2693.	0.8	29
14	Deinococcus aquaticus sp. nov., isolated from fresh water, and Deinococcus caeni sp. nov., isolated from activated sludge. International Journal of Systematic and Evolutionary Microbiology, 2008, 58, 2348-2353.	0.8	79
15	Agrococcus casei sp. nov., isolated from the surfaces of smear-ripened cheeses. International Journal of Systematic and Evolutionary Microbiology, 2007, 57, 92-97.	0.8	40
16	Diversity and biogeography of marine actinobacteria. Current Opinion in Microbiology, 2006, 9, 279-286.	2.3	149