

Matthieu Hein

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9212993/publications.pdf>

Version: 2024-02-01

34
papers

390
citations

759233
12
h-index

839539
18
g-index

52
all docs

52
docs citations

52
times ranked

396
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence and risk factors of type 2 diabetes in insomnia sufferers: a study on 1311 individuals referred for sleep examinations. <i>Sleep Medicine</i> , 2018, 46, 37-45.	1.6	32
2	Prevalence and risk factors of excessive daytime sleepiness in major depression: A study with 703 individuals referred for polysomnography. <i>Journal of Affective Disorders</i> , 2019, 243, 23-32.	4.1	32
3	Similar polysomnographic pattern in primary insomnia and major depression with objective insomnia: a sign of common pathophysiology?. <i>BMC Psychiatry</i> , 2017, 17, 273.	2.6	28
4	Prevalence and risk factors of moderate to severe obstructive sleep apnea syndrome in insomnia sufferers: a study on 1311 subjects. <i>Respiratory Research</i> , 2017, 18, 135.	3.6	27
5	Objective sleep alterations and long-term use of short or intermediate half-life benzodiazepine receptor agonists are risk factors for high blood pressure in individuals with insomnia: a study in 1272 individuals referred for sleep examinations. <i>Sleep Medicine</i> , 2019, 53, 115-123.	1.6	25
6	Prevalence and risk factors of moderate to severe obstructive sleep apnea syndrome in major depression: a observational and retrospective study on 703 subjects. <i>BMC Pulmonary Medicine</i> , 2017, 17, 165.	2.0	24
7	Prevalence and risk factors of excessive daytime sleepiness in insomnia sufferers: A study with 1311 individuals. <i>Journal of Psychosomatic Research</i> , 2017, 103, 63-69.	2.6	20
8	Hyperarousal during sleep in untreated primary insomnia sufferers: A polysomnographic study. <i>Psychiatry Research</i> , 2017, 253, 71-78.	3.3	19
9	Hyperarousal during sleep in untreated, major depressed subjects with prodromal insomnia: A polysomnographic study. <i>Psychiatry Research</i> , 2017, 258, 387-395.	3.3	19
10	Prognostic impact of glioblastoma stem cell markers OLIG2 and CCND2. <i>Cancer Medicine</i> , 2020, 9, 1069-1078.	2.8	18
11	Risk of resistant hypertension associated with restless legs syndrome and periodic limb movements during sleep: a study on 673 treated hypertensive individuals. <i>Sleep Medicine</i> , 2019, 63, 46-56.	1.6	16
12	Prevalence and Risk Factors of Type 2 Diabetes in Major Depression: A Study on 703 Individuals Referred for Sleep Examinations. <i>Psychosomatics</i> , 2018, 59, 144-157.	2.5	14
13	Alterations of neural network organisation during rapid eye movement sleep and slow-wave sleep in major depression: Implications for diagnosis, classification, and treatment. <i>Psychiatry Research - Neuroimaging</i> , 2019, 291, 71-78.	1.8	13
14	Changes in Pro-Inflammatory Markers in Detoxifying Chronic Alcohol Abusers, Divided by Lesch Typology, Reflect Cognitive Dysfunction. <i>Alcohol and Alcoholism</i> , 2017, 52, 529-534.	1.6	12
15	Risk of high blood pressure associated with objective insomnia and self-reported insomnia complaints in major depression: A study on 703 individuals. <i>Clinical and Experimental Hypertension</i> , 2019, 41, 538-547.	1.3	12
16	Impulsivity and its Therapeutic Management in Borderline Personality Disorder: a Systematic Review. <i>Psychiatric Quarterly</i> , 2020, 91, 1333-1362.	2.1	11
17	Nonremission After Electroconvulsive Therapy in Individuals With Major Depression. <i>Journal of ECT</i> , 2022, 38, 238-243.	0.6	11
18	Risk of Excessive Daytime Sleepiness Associated to Major Depression in Adolescents. <i>Psychiatric Quarterly</i> , 2021, 92, 1473-1488.	2.1	8

#	ARTICLE	IF	CITATIONS
19	Sleep ultradian cycling: Statistical distribution and links with other sleep variables, depression, insomnia and sleepiness – A retrospective study on 2,312 polysomnograms. <i>Psychiatry Research</i> , 2019, 279, 140-147.	3.3	7
20	The association between type 2 diabetes and anhedonic subtype of major depression in hypertensive individuals. <i>Journal of Clinical Hypertension</i> , 2022, 24, 156-166.	2.0	6
21	Risk of relapse within 6 months associated with borderline personality disorder in major depressed individuals treated with electroconvulsive therapy. <i>Psychiatry Research</i> , 2022, 314, 114650.	3.3	5
22	The sleep network organization during slow-wave sleep is more stable with age and has small-world characteristics more marked than during REM sleep in healthy men. <i>Neuroscience Research</i> , 2019, 145, 30-38.	1.9	4
23	Impact of number of sleep ultradian cycles on polysomnographic parameters related to REM sleep in major depression: Implications for future sleep research in psychiatry. <i>Psychiatry Research</i> , 2020, 285, 112818.	3.3	4
24	Polysomnographic markers of suicidal ideation in untreated unipolar major depressed individuals. <i>International Journal of Psychophysiology</i> , 2021, 166, 19-24.	1.0	4
25	Excessive daytime sleepiness in adolescents: current treatment strategies. <i>Sleep Science</i> , 2020, 13, 157-171.	1.0	4
26	Risk of current suicidal ideations associated with lifelong anhedonia and recent change of anhedonia in individuals with insomnia: A cross-sectional study. <i>Journal of Psychiatric Research</i> , 2022, 150, 338-345.	3.1	4
27	Alterations of neural network organization during REM sleep in women: implication for sex differences in vulnerability to mood disorders. <i>Biology of Sex Differences</i> , 2020, 11, 22.	4.1	3
28	Cardiovascular risk associated with co-morbid insomnia and sleep apnoea (COMISA) in type 2 diabetics. <i>Sleep Science</i> , 2022, 15, 184-194.	1.0	3
29	The association between type 2 diabetes and major depression in apnoeic individuals. <i>Sleep and Biological Rhythms</i> , 2022, 20, 219.	1.0	2
30	Risk of Cardiovascular Disease in Apnoeic Individuals: Role of Comorbid Insomnia Disorder. <i>Life</i> , 2022, 12, 944.	2.4	2
31	EP-2321: Evolution of OLIG2 expression during radio(chemo)therapy has a prognostic value in glioblastoma. <i>Radiotherapy and Oncology</i> , 2018, 127, S1280-S1281.	0.6	0
32	PO-1067 High expression of CCND2 in glioblastoma is associated with an increased risk of early mortality.. <i>Radiotherapy and Oncology</i> , 2019, 133, S593-S594.	0.6	0
33	Isotoxic high-dose stereotactic body radiotherapy integrated in a total multimodal neoadjuvant strategy for the treatment of localized pancreatic ductal adenocarcinoma. <i>Therapeutic Advances in Medical Oncology</i> , 2021, 13, 17588359211045860.	3.2	0
34	Impact of Alexithymia on the Lipid Profile in Major Depressed Individuals. <i>Journal of Lipids</i> , 2022, 2022, 1-11.	4.8	0