

# Matthieu Hein

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9212993/publications.pdf>

Version: 2024-02-01

34  
papers

390  
citations

758635

12  
h-index

839053

18  
g-index

52  
all docs

52  
docs citations

52  
times ranked

396  
citing authors

#	ARTICLE	IF	CITATIONS
1	Risk of current suicidal ideations associated with lifelong anhedonia and recent change of anhedonia in individuals with insomnia: A cross-sectional study. <i>Journal of Psychiatric Research</i> , 2022, 150, 338-345.	1.5	4
2	The association between type 2 diabetes and anhedonic subtype of major depression in hypertensive individuals. <i>Journal of Clinical Hypertension</i> , 2022, 24, 156-166.	1.0	6
3	The association between type 2 diabetes and major depression in apnoeic individuals. <i>Sleep and Biological Rhythms</i> , 2022, 20, 219.	0.5	2
4	Cardiovascular risk associated with co-morbid insomnia and sleep apnoea (COMISA) in type 2 diabetics. <i>Sleep Science</i> , 2022, 15, 184-194.	0.4	3
5	Nonremission After Electroconvulsive Therapy in Individuals With Major Depression. <i>Journal of ECT</i> , 2022, 38, 238-243.	0.3	11
6	Risk of relapse within 6 months associated with borderline personality disorder in major depressed individuals treated with electroconvulsive therapy. <i>Psychiatry Research</i> , 2022, 314, 114650.	1.7	5
7	Impact of Alexithymia on the Lipid Profile in Major Depressed Individuals. <i>Journal of Lipids</i> , 2022, 2022, 1-11.	1.9	0
8	Risk of Cardiovascular Disease in Apnoeic Individuals: Role of Comorbid Insomnia Disorder. <i>Life</i> , 2022, 12, 944.	1.1	2
9	Risk of Excessive Daytime Sleepiness Associated to Major Depression in Adolescents. <i>Psychiatric Quarterly</i> , 2021, 92, 1473-1488.	1.1	8
10	Polysomnographic markers of suicidal ideation in untreated unipolar major depressed individuals. <i>International Journal of Psychophysiology</i> , 2021, 166, 19-24.	0.5	4
11	Isotoxic high-dose stereotactic body radiotherapy integrated in a total multimodal neoadjuvant strategy for the treatment of localized pancreatic ductal adenocarcinoma. <i>Therapeutic Advances in Medical Oncology</i> , 2021, 13, 17588359211045860.	1.4	0
12	Prognostic impact of glioblastoma stem cell markers OLIG2 and CCND2. <i>Cancer Medicine</i> , 2020, 9, 1069-1078.	1.3	18
13	Impulsivity and its Therapeutic Management in Borderline Personality Disorder: a Systematic Review. <i>Psychiatric Quarterly</i> , 2020, 91, 1333-1362.	1.1	11
14	Impact of number of sleep ultradian cycles on polysomnographic parameters related to REM sleep in major depression: Implications for future sleep research in psychiatry. <i>Psychiatry Research</i> , 2020, 285, 112818.	1.7	4
15	Alterations of neural network organization during REM sleep in women: implication for sex differences in vulnerability to mood disorders. <i>Biology of Sex Differences</i> , 2020, 11, 22.	1.8	3
16	Excessive daytime sleepiness in adolescents: current treatment strategies. <i>Sleep Science</i> , 2020, 13, 157-171.	0.4	4
17	The sleep network organization during slow-wave sleep is more stable with age and has small-world characteristics more marked than during REM sleep in healthy men. <i>Neuroscience Research</i> , 2019, 145, 30-38.	1.0	4
18	Alterations of neural network organisation during rapid eye movement sleep and slow-wave sleep in major depression: Implications for diagnosis, classification, and treatment. <i>Psychiatry Research - Neuroimaging</i> , 2019, 291, 71-78.	0.9	13

#	ARTICLE	IF	CITATIONS
19	PO-1067 High expression of CCND2 in glioblastoma is associated with an increased risk of early mortality.. Radiotherapy and Oncology, 2019, 133, S593-S594.	0.3	0
20	Risk of resistant hypertension associated with restless legs syndrome and periodic limb movements during sleep: a study on 673 treated hypertensive individuals. Sleep Medicine, 2019, 63, 46-56.	0.8	16
21	Prevalence and risk factors of excessive daytime sleepiness in major depression: A study with 703 individuals referred for polysomnography. Journal of Affective Disorders, 2019, 243, 23-32.	2.0	32
22	Sleep ultradian cycling: Statistical distribution and links with other sleep variables, depression, insomnia and sleepinessâ€”A retrospective study on 2,312 polysomnograms. Psychiatry Research, 2019, 279, 140-147.	1.7	7
23	Risk of high blood pressure associated with objective insomnia and self-reported insomnia complaints in major depression: A study on 703 individuals. Clinical and Experimental Hypertension, 2019, 41, 538-547.	0.5	12
24	Objective sleep alterations and long-term use of short or intermediate half-life benzodiazepine receptor agonists are risk factors for high blood pressure in individuals with insomnia: a study in 1272 individuals referred for sleep examinations. Sleep Medicine, 2019, 53, 115-123.	0.8	25
25	Prevalence and risk factors of type 2 diabetes in insomnia sufferers: a study on 1311 individuals referred for sleep examinations. Sleep Medicine, 2018, 46, 37-45.	0.8	32
26	Prevalence and Risk Factors of Type 2 Diabetes in Major Depression: A Study on 703 Individuals Referred for Sleep Examinations. Psychosomatics, 2018, 59, 144-157.	2.5	14
27	EP-2321: Evolution of OLIG2 expression during radio(chemo)therapy has a prognostic value in glioblastoma. Radiotherapy and Oncology, 2018, 127, S1280-S1281.	0.3	0
28	Hyperarousal during sleep in untreated primary insomnia sufferers: A polysomnographic study. Psychiatry Research, 2017, 253, 71-78.	1.7	19
29	Prevalence and risk factors of excessive daytime sleepiness in insomnia sufferers: A study with 1311 individuals. Journal of Psychosomatic Research, 2017, 103, 63-69.	1.2	20
30	Changes in Pro-Inflammatory Markers in Detoxifying Chronic Alcohol Abusers, Divided by Lesch Typology, Reflect Cognitive Dysfunction. Alcohol and Alcoholism, 2017, 52, 529-534.	0.9	12
31	Hyperarousal during sleep in untreated, major depressed subjects with prodromal insomnia: A polysomnographic study. Psychiatry Research, 2017, 258, 387-395.	1.7	19
32	Prevalence and risk factors of moderate to severe obstructive sleep apnea syndrome in major depression: a observational and retrospective study on 703 subjects. BMC Pulmonary Medicine, 2017, 17, 165.	0.8	24
33	Prevalence and risk factors of moderate to severe obstructive sleep apnea syndrome in insomnia sufferers: a study on 1311 subjects. Respiratory Research, 2017, 18, 135.	1.4	27
34	Similar polysomnographic pattern in primary insomnia and major depression with objective insomnia: a sign of common pathophysiology?. BMC Psychiatry, 2017, 17, 273.	1.1	28