

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9213209/publications.pdf Version: 2024-02-01



Yı Vı

#	Article	IF	CITATIONS
1	Fatal attraction of Caenorhabditis elegans to predatory fungi through 6-methyl-salicylic acid. Nature Communications, 2021, 12, 5462.	12.8	34
2	Intercellular communication is required for trap formation in the nematode-trapping fungus Duddingtonia flagrans. PLoS Genetics, 2019, 15, e1008029.	3.5	59
3	A Gene Cluster for the Biosynthesis of Dibenzodioxocinons in the Endophyte Pestalotiopsis microspora, a Taxol Producer. Journal of Microbiology and Biotechnology, 2019, 29, 1570-1579.	2.1	7
4	Activatable Protein Nanoparticles for Targeted Delivery of Therapeutic Peptides. Advanced Materials, 2018, 30, 1705383.	21.0	38
5	Roles of phospholipid methyltransferases in pycnidia development, stress tolerance and secondary metabolism in the taxol-producing fungus Pestalotiopsis microspore. Microbiological Research, 2018, 210, 33-42.	5.3	13
6	Cα-cAMP/PKA pathway positively regulates pigmentation, chaetoglobosin A biosynthesis and sexual development in Chaetomium globosum. PLoS ONE, 2018, 13, e0195553.	2.5	14
7	The Gα1-cAMP signaling pathway controls conidiation, development and secondary metabolism in the taxol-producing fungus Pestalotiopsis microspora. Microbiological Research, 2017, 203, 29-39.	5.3	15
8	Amino Acid Sensor Kinase Gcn2 Is Required for Conidiation, Secondary Metabolism, and Cell Wall Integrity in the Taxol-Producer Pestalotiopsis microspora. Frontiers in Microbiology, 2017, 8, 1879.	3.5	9
9	A Bâ€type histone acetyltransferase Hat1 regulates secondary metabolism, conidiation, and cell wall integrity in the taxolâ€producing fungus <i>Pestalotiopsis microspora</i> . Journal of Basic Microbiology, 2016, 56, 1380-1391.	3.3	14
10	Orotidine 5′â€phosphate decarboxylaseâ€based reusable in situ genetic editing system: Development and application in taxolâ€producing <i>Pestalotiopsis microspora</i> . Engineering in Life Sciences, 2015, 15, 542-549.	3.6	11
11	Melanin is required for the formation of the multi-cellular conidia in the endophytic fungus Pestalotiopsis microspora. Microbiological Research, 2015, 179, 1-11.	5.3	48
12	High frequency of homologous gene disruption by single-stranded DNA in the taxol-producing fungus Pestalotiopsis microspora. Annals of Microbiology, 2015, 65, 2151-2160.	2.6	10
13	A Putative Histone Deacetylase Modulates the Biosynthesis of Pestalotiollide B and Conidiation in Pestalotiopsis microspora. Journal of Microbiology and Biotechnology, 2015, 25, 579-588.	2.1	27