

Jan A Walewski

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9215005/publications.pdf>

Version: 2024-02-01

124
papers

10,703
citations

117453

34
h-index

31759

101
g-index

127
all docs

127
docs citations

127
times ranked

8741
citing authors

#	ARTICLE	IF	CITATIONS
1	CHOP-like chemotherapy plus rituximab versus CHOP-like chemotherapy alone in young patients with good-prognosis diffuse large-B-cell lymphoma: a randomised controlled trial by the MabThera International Trial (MInT) Group. <i>Lancet Oncology, The</i> , 2006, 7, 379-391.	5.1	1,840
2	CVP chemotherapy plus rituximab compared with CVP as first-line treatment for advanced follicular lymphoma. <i>Blood</i> , 2005, 105, 1417-1423.	0.6	896
3	CHOP-like chemotherapy with or without rituximab in young patients with good-prognosis diffuse large-B-cell lymphoma: 6-year results of an open-label randomised study of the MabThera International Trial (MInT) Group. <i>Lancet Oncology, The</i> , 2011, 12, 1013-1022.	5.1	633
4	Brentuximab vedotin as consolidation therapy after autologous stem-cell transplantation in patients with Hodgkin's lymphoma at risk of relapse or progression (AETHERA): a randomised, double-blind, placebo-controlled, phase 3 trial. <i>Lancet, The</i> , 2015, 385, 1853-1862.	6.3	633
5	Brentuximab Vedotin with Chemotherapy for Stage III or IV Hodgkin's Lymphoma. <i>New England Journal of Medicine</i> , 2018, 378, 331-344.	13.9	564
6	Brentuximab vedotin or physician's choice in CD30-positive cutaneous T-cell lymphoma (ALCANZA): an international, open-label, randomised, phase 3, multicentre trial. <i>Lancet, The</i> , 2017, 390, 555-566.	6.3	444
7	Chemotherapy plus Involved-Field Radiation in Early-Stage Hodgkin's Disease. <i>New England Journal of Medicine</i> , 2007, 357, 1916-1927.	13.9	412
8	Belinostat in Patients With Relapsed or Refractory Peripheral T-Cell Lymphoma: Results of the Pivotal Phase II BELIEF (CLN-19) Study. <i>Journal of Clinical Oncology</i> , 2015, 33, 2492-2499.	0.8	394
9	Safety and efficacy of ofatumumab, a fully human monoclonal anti-CD20 antibody, in patients with relapsed or refractory B-cell chronic lymphocytic leukemia: a phase 1-2 study. <i>Blood</i> , 2008, 111, 1094-1100.	0.6	369
10	Addition of high-dose cytarabine to immunochemotherapy before autologous stem-cell transplantation in patients aged 65 years or younger with mantle cell lymphoma (MCL Younger): a randomised, open-label, phase 3 trial of the European Mantle Cell Lymphoma Network. <i>Lancet, The</i> , 2016, 388, 565-575.	6.3	328
11	Rituximab versus a watch-and-wait approach in patients with advanced-stage, asymptomatic, non-bulky follicular lymphoma: an open-label randomised phase 3 trial. <i>Lancet Oncology, The</i> , 2014, 15, 424-435.	5.1	304
12	First clinical use of ofatumumab, a novel fully human anti-CD20 monoclonal antibody in relapsed or refractory follicular lymphoma: results of a phase 1/2 trial. <i>Blood</i> , 2008, 111, 5486-5495.	0.6	247
13	Molecular remission is an independent predictor of clinical outcome in patients with mantle cell lymphoma after combined immunochemotherapy: a European MCL intergroup study. <i>Blood</i> , 2010, 115, 3215-3223.	0.6	243
14	Improved outcome of adult Burkitt lymphoma/leukemia with rituximab and chemotherapy: report of a large prospective multicenter trial. <i>Blood</i> , 2014, 124, 3870-3879.	0.6	236
15	A prospective clinicopathologic study of dose-modified CODOX-M/IVAC in patients with sporadic Burkitt lymphoma defined using cytogenetic and immunophenotypic criteria (MRC/NCRI LY10 trial). <i>Blood</i> , 2008, 112, 2248-2260.	0.6	199
16	Outcome of adult patients with T-lymphoblastic lymphoma treated according to protocols for acute lymphoblastic leukemia. <i>Blood</i> , 2002, 99, 4379-4385.	0.6	195
17	ESMO Guidelines consensus conference on malignant lymphoma 2011 part 1: diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL) and chronic lymphocytic leukemia (CLL). <i>Annals of Oncology</i> , 2013, 24, 561-576.	0.6	193
18	Prognostic significance of maximum tumour (bulk) diameter in young patients with good-prognosis diffuse large-B-cell lymphoma treated with CHOP-like chemotherapy with or without rituximab: an exploratory analysis of the MabThera International Trial Group (MInT) study. <i>Lancet Oncology, The</i> , 2008, 9, 435-444.	5.1	190

#	ARTICLE	IF	CITATIONS
19	Five-year PFS from the AETHERA trial of brentuximab vedotin for Hodgkin lymphoma at high risk of progression or relapse. <i>Blood</i> , 2018, 132, 2639-2642.	0.6	172
20	Confirmation of the Mantle-Cell Lymphoma International Prognostic Index in Randomized Trials of the European Mantle-Cell Lymphoma Network. <i>Journal of Clinical Oncology</i> , 2014, 32, 1338-1346.	0.8	137
21	Lenalidomide versus investigator's choice in relapsed or refractory mantle cell lymphoma (MCL-002); Tj ETQq1 1 0.784314 rgBT /Over 5.1 135	0.8	111
22	Comparison of ABVD and Alternating or Hybrid Multidrug Regimens for the Treatment of Advanced Hodgkin's Lymphoma: Results of the United Kingdom Lymphoma Group LY09 Trial (ISRCTN97144519). <i>Journal of Clinical Oncology</i> , 2005, 23, 9208-9218.	0.8	130
23	Rituximab Purging and/or Maintenance in Patients Undergoing Autologous Transplantation for Relapsed Follicular Lymphoma: A Prospective Randomized Trial From the Lymphoma Working Party of the European Group for Blood and Marrow Transplantation. <i>Journal of Clinical Oncology</i> , 2013, 31, 1624-1630.	0.8	111
24	Bortezomib plus rituximab versus rituximab alone in patients with relapsed, rituximab-naïve or rituximab-sensitive, follicular lymphoma: a randomised phase 3 trial. <i>Lancet Oncology</i> , The, 2011, 12, 773-784.	5.1	98
25	Brentuximab vedotin with chemotherapy for stage III/IV classical Hodgkin lymphoma: 3-year update of the ECHELON-1 study. <i>Blood</i> , 2020, 135, 735-742.	0.6	86
26	Central nervous system involvement by multiple myeloma: A multi-institutional retrospective study of 172 patients in daily clinical practice. <i>American Journal of Hematology</i> , 2016, 91, 575-580.	2.0	83
27	Brentuximab vedotin with chemotherapy for stage III or IV classical Hodgkin lymphoma (ECHELON-1): 5-year update of an international, open-label, randomised, phase 3 trial. <i>Lancet Haematology</i> , the, 2021, 8, e410-e421.	2.2	83
28	Ibrutinib plus Bendamustine and Rituximab in Untreated Mantle-Cell Lymphoma. <i>New England Journal of Medicine</i> , 2022, 386, 2482-2494.	13.9	83
29	Treatment of Older Patients With Mantle Cell Lymphoma (MCL): Long-Term Follow-Up of the Randomized European MCL Elderly Trial. <i>Journal of Clinical Oncology</i> , 2020, 38, 248-256.	0.8	73
30	ESMO Consensus Conference on malignant lymphoma: general perspectives and recommendations for the clinical management of the elderly patient with malignant lymphoma. <i>Annals of Oncology</i> , 2018, 29, 544-562.	0.6	64
31	Partial trisomy 11, dup(11)(q23q13), as a defect characterizing lymphomas with Burkitt pathomorphology without MYC gene rearrangement. <i>Medical Oncology</i> , 2011, 28, 1589-1595.	1.2	46
32	Randomized phase 3 ALCANZA study of brentuximab vedotin vs physician's choice in cutaneous T-cell lymphoma: final data. <i>Blood Advances</i> , 2021, 5, 5098-5106.	2.5	46
33	Frontline low-dose alemtuzumab with fludarabine and cyclophosphamide prolongs progression-free survival in high-risk CLL. <i>Blood</i> , 2014, 123, 3255-3262.	0.6	45
34	A comprehensive flow-cytometry-based immunophenotypic characterization of Burkitt-like lymphoma with 11q aberration. <i>Modern Pathology</i> , 2018, 31, 732-743.	2.9	42
35	Pharmacokinetics and pharmacokinetic/pharmacodynamic associations of ofatumumab, a human monoclonal CD20 antibody, in patients with relapsed or refractory chronic lymphocytic leukaemia: a phase 1 study. <i>British Journal of Haematology</i> , 2010, 150, 58-71.	1.2	37
36	Immunogenicity of Influenza Vaccination in Patients with Non-Hodgkin Lymphoma. <i>Journal of Clinical Immunology</i> , 2007, 27, 339-346.	2.0	34

#	ARTICLE	IF	CITATIONS
37	Effect of Cytochrome P450 3A4 Inducers on the Pharmacokinetic, Pharmacodynamic and Safety Profiles of Bortezomib in Patients with Multiple Myeloma or Non-Hodgkin's Lymphoma. <i>Clinical Pharmacokinetics</i> , 2011, 50, 781-791.	1.6	31
38	Primary Mediastinal Large B-cell Lymphoma. <i>Current Hematologic Malignancy Reports</i> , 2014, 9, 273-283.	1.2	31
39	Quality of life results from a phase 3 study of brentuximab vedotin consolidation following autologous haematopoietic stem cell transplant for persons with Hodgkin lymphoma. <i>British Journal of Haematology</i> , 2016, 175, 860-867.	1.2	30
40	Cutaneous involvement in multiple myeloma: a multi-institutional retrospective study of 53 patients. <i>Leukemia and Lymphoma</i> , 2016, 57, 2071-2076.	0.6	30
41	Update on the molecular pathogenesis and clinical treatment of mantle cell lymphoma: report of the 10th annual conference of the European Mantle Cell Lymphoma Network. <i>Leukemia and Lymphoma</i> , 2011, 52, 2226-2236.	0.6	29
42	Response to brentuximab vedotin versus physician's choice by CD30 expression and large cell transformation status in patients with mycosis fungoides: An ALCANZA sub-analysis. <i>European Journal of Cancer</i> , 2021, 148, 411-421.	1.3	27
43	Alternating Courses of 3x CHOP and 3x DHAP Plus Rituximab Followed by a High Dose ARA-C Containing Myeloablative Regimen and Autologous Stem Cell Transplantation (ASCT) Is Superior to 6 Courses CHOP Plus Rituximab Followed by Myeloablative Radiochemotherapy and ASCT In Mantle Cell Lymphoma: Results of the MCL Younger Trial of the European Mantle Cell Lymphoma Network (MCL Tj ETQq1 1 0.784314 rgBT /Over	0.6	27
44	Individual Quality Assessment of Autografting by Probability Estimation for Clinical Endpoints: A Prospective Validation Study from the European Group for Blood and Marrow Transplantation. <i>Biology of Blood and Marrow Transplantation</i> , 2013, 19, 1670-1676.	2.0	26
45	The addition of rituximab to fludarabine and cyclophosphamide chemotherapy results in a significant improvement in overall survival in patients with newly diagnosed mantle cell lymphoma: results of a randomized UK National Cancer Research Institute trial. <i>Haematologica</i> , 2016, 101, 235-240.	1.7	24
46	Treatment for patients with relapsed/refractory mantle cell lymphoma: European-based recommendations. <i>Leukemia and Lymphoma</i> , 2018, 59, 1814-1828.	0.6	23
47	Mapping cancer research across Central and Eastern Europe, the Russian Federation and Central Asia: Implications for future national cancer control planning. <i>European Journal of Cancer</i> , 2018, 104, 127-136.	1.3	23
48	Randomized comparison of cladribine alone or in combination with cyclophosphamide, and cyclophosphamide, vincristine and prednisone in previously untreated low-grade B-cell non-Hodgkin lymphoma patients. <i>Cancer</i> , 2008, 113, 367-375.	2.0	22
49	PD-L1 Overexpression, SWI/SNF Complex Deregulation, and Profound Transcriptomic Changes Characterize Cancer-Dependent Exhaustion of Persistently Activated CD4+ T Cells. <i>Cancers</i> , 2021, 13, 4148.	1.7	22
50	Patient-reported quality of life in patients with relapsed/refractory cutaneous T-cell lymphoma: Results from the randomised phase III ALCANZA study. <i>European Journal of Cancer</i> , 2020, 133, 120-130.	1.3	21
51	Updated Efficacy and Safety Data from the AETHERA Trial of Consolidation with Brentuximab Vedotin after Autologous Stem Cell Transplant (ASCT) in Hodgkin Lymphoma Patients at High Risk of Relapse. <i>Blood</i> , 2015, 126, 3172-3172.	0.6	20
52	Adjuvant vaccination with melanoma antigen-pulsed dendritic cells in stage III melanoma patients. <i>Medical Oncology</i> , 2012, 29, 2966-2977.	1.2	18
53	Prospective study of brentuximab vedotin in relapsed/refractory Hodgkin lymphoma patients who are not suitable for stem cell transplant or multi-agent chemotherapy. <i>British Journal of Haematology</i> , 2018, 183, 400-410.	1.2	18
54	miR expression in MYC-negative DLBCL/BL with partial trisomy 11 is similar to classical Burkitt lymphoma and different from diffuse large B-cell lymphoma. <i>Tumor Biology</i> , 2015, 36, 5377-5388.	0.8	17

#	ARTICLE	IF	CITATIONS
55	Current Treatment of Chronic Lymphocytic Leukemia. <i>Current Treatment Options in Oncology</i> , 2017, 18, 5.	1.3	17
56	Resminostat in patients with relapsed or refractory Hodgkin lymphoma: results of the phase II SAPHIRE study. <i>Leukemia and Lymphoma</i> , 2019, 60, 675-684.	0.6	17
57	Frequent aberrations of chromosome 8 in aggressive B-cell non-Hodgkin lymphoma. <i>Cancer Genetics and Cytogenetics</i> , 2005, 156, 114-121.	1.0	16
58	Bortezomib plus rituximab versus rituximab in patients with high-risk, relapsed, rituximab-naïve or rituximab-sensitive follicular lymphoma: subgroup analysis of a randomized phase 3 trial. <i>Journal of Hematology and Oncology</i> , 2012, 5, 67.	6.9	16
59	First-line R-CVP versus R-CHOP induction immunochemotherapy for indolent lymphoma with rituximab maintenance. A multicentre, phase III randomized study by the Polish Lymphoma Research Group PLRG4. <i>British Journal of Haematology</i> , 2020, 188, 898-906.	1.2	16
60	FcγRIIA and FcγRIIIA polymorphisms do not influence survival and response to rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone immunochemotherapy in patients with diffuse large B-cell lymphoma. <i>Leukemia and Lymphoma</i> , 2011, 52, 1604-1606.	0.6	15
61	Phase II Randomized, Multicenter Study of Lenalidomide Vs Best Investigator's Choice in Relapsed/Refractory Mantle Cell Lymphoma: Results of the MCL-002 (SPRINT) Study. <i>Blood</i> , 2014, 124, 626-626.	0.6	15
62	Serum macrophage colony-stimulating factor (M-CSF) in patients with Hodgkin lymphoma. <i>Medical Oncology</i> , 2012, 29, 2143-2147.	1.2	14
63	Tumor and Cerebrospinal Fluid microRNAs in Primary Central Nervous System Lymphomas. <i>Cancers</i> , 2019, 11, 1647.	1.7	14
64	Human regulatory T cells suppress proliferation of B lymphoma cells. <i>Leukemia and Lymphoma</i> , 2016, 57, 1903-1920.	0.6	13
65	Primary prophylaxis with G-CSF may improve outcomes in patients with newly diagnosed stage III/IV Hodgkin lymphoma treated with brentuximab vedotin plus chemotherapy. <i>Leukemia and Lymphoma</i> , 2020, 61, 2931-2938.	0.6	13
66	Improving outcomes after autologous transplantation in relapsed/refractory Hodgkin lymphoma: a European expert perspective. <i>BMC Cancer</i> , 2020, 20, 1088.	1.1	12
67	Fine Needle Aspiration Biopsy with Flow Cytometry Is a Reliable Method for Defining Immunophenotype of Prognostic Value in T Lymphoblastic Lymphoma.. <i>Blood</i> , 2009, 114, 3928-3928.	0.6	11
68	Implications of the European Organisation for Research And Treatment Of Cancer (EORTC) Guidelines on the Use of Granulocyte Colony-Stimulating Factor (G-CSF) for Lymphoma Care. <i>Clinical Drug Investigation</i> , 2009, 29, 491-513.	1.1	10
69	Zalecenia Polskiej Grupy Szpiczakowej dotyczÄ...ce rozpoznawania i leczenia szpiczaka plazmocytoowego oraz innych dyskrazji plazmocytoowych na rok 2016. <i>Acta Haematologica Polonica</i> , 2016, 47, 39-85.	0.1	10
70	Lenalidomide potentiates CD4+CD25+Treg-related suppression of lymphoma B-cell proliferation. <i>Clinical and Experimental Medicine</i> , 2017, 17, 193-207.	1.9	10
71	The HOVON68 CLL trial revisited: performance status and comorbidity affect survival in elderly patients with chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2017, 58, 594-600.	0.6	10
72	Prospective subgroup analyses of the randomized MCL-002 (SPRINT) study: lenalidomide versus investigator's choice in relapsed or refractory mantle cell lymphoma. <i>British Journal of Haematology</i> , 2018, 180, 224-235.	1.2	10

#	ARTICLE	IF	CITATIONS
73	Addition of High-Dose Cytarabine to Immunochemotherapy before Autologous Stem-Cell Transplantation in Patients Aged 65 Years or Younger with Mantle Cell Lymphoma (MCL Younger): A Long-Term Follow-up of the Randomized, Open-Label, Phase 3 Trial of the European Mantle Cell Lymphoma Network. <i>Blood</i> , 2021, 138, 380-380.	0.6	10
74	Recovery of dendritic cell counts and function in peripheral blood of cancer patients after chemotherapy. <i>Cytokines, Cellular & Molecular Therapy</i> , 2002, 7, 15-24.	0.3	9
75	Procollagen I amino-terminal propeptide as a potential marker for multiple myeloma. <i>Clinical Biochemistry</i> , 2010, 43, 604-608.	0.8	9
76	Autologous stem cell transplantation as consolidation therapy for patients with peripheral T cell lymphoma in first remission: long-term outcome and risk factors analysis. <i>Annals of Hematology</i> , 2013, 92, 925-933.	0.8	9
77	Polish Lymphoma Research Group Experience With Bexarotene in the Treatment of Cutaneous T-Cell Lymphoma. <i>American Journal of Therapeutics</i> , 2016, 23, e749-e756.	0.5	9
78	Soluble CD52 is an indicator of disease activity in chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2017, 58, 2356-2362.	0.6	9
79	Unusual Cyclin D1 Positive Marginal Zone Lymphoma of Mediastinum. <i>Medical Oncology</i> , 2006, 23, 423-428.	1.2	8
80	Phase 2 Study Evaluating the Efficacy and Safety of Parsaclisib in Patients with Relapsed or Refractory Mantle Cell Lymphoma Not Previously Treated with a BTK Inhibitor (CITADEL-205). <i>Blood</i> , 2020, 136, 22-23.	0.6	8
81	A survey of prognostic value of serum factors in multiple myeloma patients before treatment: macrophage-colony stimulating factor (M-CSF) is a powerful predictor of survival. <i>Medical Oncology</i> , 2011, 28, 194-198.	1.2	7
82	Discriminant analysis involving serum cytokine levels and prediction of the response to therapy of patients with Hodgkin lymphoma. <i>Tumor Biology</i> , 2012, 33, 1733-1738.	0.8	7
83	Population-based epidemiological data of follicular lymphoma in Poland: 15 years of observation. <i>Scientific Reports</i> , 2020, 10, 14610.	1.6	7
84	Higher genome variability within metabolism genes associates with recurrent <i>Clostridium difficile</i> infection. <i>BMC Microbiology</i> , 2021, 21, 36.	1.3	7
85	Mantle Cell Lymphoma Presenting with Paraproteinemia. <i>Medical Oncology</i> , 2005, 22, 319-324.	1.2	6
86	Primary refractory multiple myeloma: a real-world experience with 85 cases. <i>Leukemia and Lymphoma</i> , 2020, 61, 2868-2875.	0.6	6
87	Unusual IgD+/CD38-Follicular Lymphoma with Leukemic Presentation. <i>Medical Oncology</i> , 2006, 23, 131-136.	1.2	5
88	Zalecenia Polskiej Grupy Szpiczakowej dotyczÄ...ce rozpoznawania i leczenia szpiczaka plazmocytoowego na rok 2012. <i>Acta Haematologica Polonica</i> , 2012, 43, 7-47.	0.1	5
89	Treatment strategy based on gemcitabine-containing salvage chemotherapy used with intent to proceed to second stem cell transplant for patients with Hodgkin lymphoma relapsing after a prior autologous transplant. <i>Leukemia and Lymphoma</i> , 2013, 54, 973-978.	0.6	5
90	Zalecenia Polskiej Grupy Szpiczakowej dotyczÄ...ce rozpoznawania i leczenia szpiczaka plazmocytoowego oraz innych dyskrazji plazmocytoowych na rok 2017. <i>Acta Haematologica Polonica</i> , 2017, 48, 55-103.	0.1	5

#	ARTICLE	IF	CITATIONS
91	Aggressive B-Cell Lymphoma: Chasing the Target. <i>Journal of Investigative Medicine</i> , 2020, 68, 331-334.	0.7	5
92	Brentuximab Vedotin with Chemotherapy for Patients with Previously Untreated, Stage III/IV Classical Hodgkin Lymphoma: 5-Year Update of the ECHELON-1 Study. <i>Blood</i> , 2020, 136, 26-28.	0.6	5
93	European MCL Network: An Update on Current First Line Trials.. <i>Blood</i> , 2007, 110, 388-388.	0.6	5
94	Role of rituximab in the first-line therapy of high-risk diffuse large B-cell lymphoma: a retrospective analysis by the Polish Lymphoma Research Group. <i>Polish Archives of Internal Medicine</i> , 2015, 125, 741-748.	0.3	5
95	CHOP-21 for unfavorable Hodgkin's lymphoma. An exploratory study. <i>Medical Oncology</i> , 2010, 27, 262-267.	1.2	4
96	Bendamustine as Monotherapy and in Combination Regimens for the Treatment of Chronic Lymphocytic Leukemia and Non-Hodgkin Lymphoma: A Retrospective Analysis. <i>Chemotherapy</i> , 2013, 59, 280-289.	0.8	4
97	Efficacy of siltuximab in the treatment of idiopathic multicentric castlemans disease, the first Polish, real-world experience with long-term observation. <i>Leukemia and Lymphoma</i> , 2021, 62, 3031-3034.	0.6	4
98	Comparison of benefits of early, delayed, and no administration of G-CSF after autologous peripheral blood stem cell transplantation in lymphoma patients. <i>Annals of Transplantation</i> , 2013, 18, 336-341.	0.5	4
99	The Selective Bruton Tyrosine Kinase (BTK) Inhibitor TG-1701 As Monotherapy and in Combination with Ublituximab and Umbralisib (U2) in Patients with B-Cell Malignancies. <i>Blood</i> , 2021, 138, 1549-1549.	0.6	4
100	Significance of CD10 protein expression in the diagnostics of follicular lymphoma: A comparison of conventional immunohistochemistry with flow cytometry supported by the establishment of BCL2 and BCL6 rearrangements. <i>International Journal of Laboratory Hematology</i> , 2020, 42, 453-463.	0.7	3
101	Plasma alemtuzumab levels in patients with chronic lymphocytic leukemia treated with alemtuzumab combined with chemotherapy reflect the efficacy of the treatment: a hypothesis. <i>Leukemia and Lymphoma</i> , 2013, 54, 790-793.	0.6	2
102	PD1 distribution pattern, regardless of the cell origin, is an independent microenvironmental prognostic factor for progression-free survival in follicular lymphoma. <i>Pathology Research and Practice</i> , 2020, 216, 153096.	1.0	2
103	Prognostic Value of the Immunological Subtypes of Adolescent and Adult T-Cell Lymphoblastic Lymphoma; an Ultra-High-Risk Pro-T/CD2(âˆ†) Subtype. <i>Cancers</i> , 2021, 13, 1911.	1.7	2
104	DA-EPOCH-R Is an Effective Regimen in High Grade B-Cell Lymphoma Defined By Cell-of-Origin, Karyotype and BCL2/MYC/BCL6 Status and Expression. <i>Blood</i> , 2016, 128, 1754-1754.	0.6	2
105	Improved survival of Burkitt lymphoma/leukemia patients: observations from Poland, 1999â€“2020. <i>Annals of Hematology</i> , 2022, 101, 1059-1065.	0.8	2
106	High efficacy of intensive immunochemotherapy for primary mediastinal B-cell lymphoma with prolonged follow up. <i>Scientific Reports</i> , 2022, 12, .	1.6	2
107	Tolerability and efficacy of GM-CSF [Leucomax] in patients with small cell lung cancer treated with intensive chemotherapy. <i>Medical Oncology</i> , 1996, 13, 199-205.	1.2	1
108	EZH2 Expression in Follicular Lymphoma Is Variable and Independent from the Progression of Disease Within 24 Months of First Treatment. <i>Anticancer Research</i> , 2020, 40, 6685-6697.	0.5	1

#	ARTICLE	IF	CITATIONS
109	Isolation and Characteristics of Dendritic Cell Progenitors from the Bone Marrow of the Hodgkinâ€™s Disease Patients. <i>Advances in Experimental Medicine and Biology</i> , 1995, 378, 553-555.	0.8	1
110	CHOP Chemotherapy for Unfavorable or Advanced Hodgkinâ€™s Lymphoma. A Pilot Study.. <i>Blood</i> , 2004, 104, 1311-1311.	0.6	1
111	Flow Cytometry and Cytogenetics of Fine Needle Aspiration Biopsy Samples Is a Reliable Method for Diagnosing Burkitt Lymphoma. Evaluation of 78 Cases from a Single-Institution. <i>Blood</i> , 2014, 124, 1640-1640.	0.6	1
112	Significance of a Critical Set of 11q Chromosome Aberrations for Diagnosis of MYC Negative Burkitt Lymphoma. <i>Blood</i> , 2015, 126, 2679-2679.	0.6	1
113	Serendipity w drodze do skutecznego leczenia: poczÄ™tki i perspektywy immunochemioterapii, czyli od nitrogranulogenu do receptorÄ™w chimerowych. <i>Nowotwory</i> , 2015, 65, 96-102.	0.1	1
114	Ofatumumab with iphosphamide, etoposide and cytarabine for patients with transplantationâ€™ineligible relapsed and refractory diffuse large Bâ€™cell lymphoma. <i>British Journal of Haematology</i> , 2022, , .	1.2	1
115	Monoclonal antibody therapy for classical Hodgkin lymphoma. <i>Clinical Investigation</i> , 2013, 3, 899-910.	0.0	0
116	Follow-up in aggressive lymphomas. <i>Memo - Magazine of European Medical Oncology</i> , 2014, 7, 80-82.	0.3	0
117	ChÄ™niaki agresywne z komÄ™rek B z podwÄ™jnÄ™.../potrÄ™jnÄ™... translokacjÄ™... â€™ double/triple hit lymphoma. <i>Acta Haematologica Polonica</i> , 2015, 46, 263-268.	0.1	0
118	Zalecenia Polskiej Grupy Szpiczakowej dotyczÄ™ce rozpoznawania i leczenia szpiczaka plazmocytoowego oraz innych dyskrazji plazmocytoowych na rok 2015. <i>Acta Haematologica Polonica</i> , 2015, 46, 159-211.	0.1	0
119	MoÅ¼liwoÅ›ci leczenia indukcyjnego chorych na szpiczaka plazmocytoowego kwalifikujÄ™cych siÄ™ do chemioterapii wysokodawkowanej wspomaganÄ™ autologicznÄ™... transplantacjÄ™... komÄ™rek krwiotwÄ™rczych a aktualne zalecenia Polskiej Grupy Szpiczakowej. <i>Acta Haematologica Polonica</i> , 2017, 48, 104-111.	0.1	0
120	Peripheral Blood Cells from Patients with Hodgkinâ€™s and Diffuse Large B Cell Lymphomas May Be a Better Source of Candidate Diagnostic miRNAs Than Circulating miRNAs. <i>BioMed Research International</i> , 2021, 2021, 1-9.	0.9	0
121	Czy naleÅ¼y stosowaÄ™ radioterapiÄ™ w chÄ™niaku rozlanym z duÅ¼ych komÄ™rek B? GÄ™os na nie. <i>Nowotwory</i> , 2014, 64, 81-83.	0.1	0
122	Czynniki pobudzajÄ™ce granulopoezÄ™ sÄ™... naduÅ¼ywane w systemowym leczeniu onkologicznym. <i>Nowotwory</i> , 2016, 65, 529-534.	0.1	0
123	Stanowisko polskich ekspertÄ™w dotyczÄ™ce zastosowania leku brentuksymab vedotin w leczeniu chorych na pierwotne chÄ™niaki skÄ™ry CD30+. <i>Hematologia</i> , 2018, 9, 83-89.	0.0	0
124	Rituximab and Chemotherapy for Newly Diagnosed Follicular Lymphoma - Real World Report of Polish Lymphoma Research Group. <i>Chemotherapy</i> , 2022, , .	0.8	0