Temel Ozek

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9217204/publications.pdf

Version: 2024-02-01

186

all docs

186 3,762 29
papers citations h-index

186

docs citations

h-index g-index

186
3452
times ranked citing authors

205818

48

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Chemical characterization, antioxidant activity, ï†-amylase and acetylcholinesterase inhibitory potential of Angelica pancicii Vandas ex Velen. Boletin Latinoamericano Y Del Caribe De Plantas Medicinales Y Aromaticas, 2022, 21, 418-430. | 0.2 | O |
| 2 | Effects of different nitrogen doses on thymoquinone and fatty acid composition in seed oil of black cumin (<i>Nigella sativa</i> L.). JAOCS, Journal of the American Oil Chemists' Society, 2022, 99, 229-237. | 0.8 | 3 |
| 3 | Neutrophil Immunomodulatory Activity of Farnesene, a Component of ArtemisiaÂdracunculus Essential Oils. Pharmaceuticals, 2022, 15, 642. | 1.7 | 12 |
| 4 | Leaf essential oil analysis and anatomical study of Cordia myxa from Turkey. Plant Biosystems, 2021, 155, 204-210. | 0.8 | 3 |
| 5 | Chemical Composition and Immunomodulatory Activity of Essential Oils from Rhododendron albiflorum. Molecules, 2021, 26, 3652. | 1.7 | 16 |
| 6 | Essential Oil Compositions and Site Characteristics of Sideritis pisidica in Natural Habitat. Contemporary Problems of Ecology, 2021, 14, 675-689. | 0.3 | 1 |
| 7 | Innate Immunomodulatory Activity of Cedrol, a Component of Essential Oils Isolated from Juniperus Species. Molecules, 2021, 26, 7644. | 1.7 | 17 |
| 8 | Essential Oils from Monarda fistulosa: Chemical Composition and Activation of Transient Receptor Potential A1 (TRPA1) Channels. Molecules, 2020, 25, 4873. | 1.7 | 24 |
| 9 | Chemical Composition and Immunomodulatory Activity of Hypericum perforatum Essential Oils. Biomolecules, 2020, 10, 916. | 1.8 | 35 |
| 10 | Caffeoylquinic Acids, Cytotoxic, Antioxidant, Acetylcholinesterase and Tyrosinase Enzyme Inhibitory Activities of Six <i>Inula</i> Species from Bulgaria. Chemistry and Biodiversity, 2020, 17, e2000051. | 1.0 | 31 |
| 11 | Volatile constituents of four Inula species of Bulgarian origin. Biochemical Systematics and Ecology, 2020, 90, 104035. | 0.6 | 4 |
| 12 | Essential oils and lipids from the flowers of two varieties of Ocimum basilicum L. cultivated in Uzbekistan. Journal of Essential Oil Research, 2020, 32, 323-330. | 1.3 | 6 |
| 13 | The Leaf and the Gall Volatiles of Salvia fruticosa Miller from Turkey: Chemical Composition and Biological Activities. Records of Natural Products, 2020, 15, 10-24. | 1.3 | 3 |
| 14 | Essential oil composition of a medicinally important Cape species: Pentzia punctata (Asteraceae). South African Journal of Botany, 2019, 127, 208-212. | 1.2 | 5 |
| 15 | Furanocoumarin Content, Antioxidant Activity, and Inhibitory Potential of Heracleum verticillatum, Heracleum sibiricum, Heracleum angustisectum, and Heracleum ternatum Extracts against Enzymes Involved in Alzheimer's Disease and Type II Diabetes. Chemistry and Biodiversity, 2019, 16, e1800672. | 1.0 | 7 |
| 16 | Glandular trichome structures and chemical composition of the volatiles of five <i>Ribes </i> species from Turkey. Journal of Essential Oil Research, 2019, 31, 111-119. | 1.3 | 3 |
| 17 | Phytochemical Profiling and Evaluation of Marrubium sivasense AytaÃS, Akgül & Ekici for Antioxidant Activity and Inhibition Effects on î±-Amylase, Lipoxygenase, Xanthine Oxidase and Tyrosinase Enzymes. Journal of the Turkish Chemical Society, Section A: Chemistry, 2019, 6, 281-292. | 0.4 | 3 |
| 18 | Assessment of Endemic Cota fulvida (Asteraceae) for Phytochemical Composition and Inhibitory Activities against Oxidation, ï-Amylase, Lipoxygenase, Xanthine Oxidase and Tyrosinase Enzymes. Records of Natural Products, 2019, 13, 333-345. | 1.3 | 8 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | The effect of different nitrogen levels on yield and quality of stevia (<i>Stevia rebaudiana</i> bert.). Journal of Plant Nutrition, 2018, 41, 1130-1137. | 0.9 | 12 |
| 20 | The Effect of the Plant Age and Growth Period on the Nutritional Substance, Chlorophyll and Steviol Glycoside Rates in Stevia (<i>Stevia Rebaudiana</i> Bertoni) Leaves. Communications in Soil Science and Plant Analysis, 2018, 49, 291-302. | 0.6 | 2 |
| 21 | Biological evaluation, overpressured layer chromatography separation, and isolation of a new acetylenic derivative compound from <i>Prangos platychlaena</i> ssp. <i>platychlaena</i> fruit essential oils. Journal of Planar Chromatography - Modern TLC, 2018, 31, 61-71. | 0.6 | 7 |
| 22 | Essential oil composition of <i>Pentzia incana</i> (Asteraceae), an important natural pasture plant in the Karoo region of South Africa. African Journal of Range and Forage Science, 2018, 35, 137-145. | 0.6 | 8 |
| 23 | Isolation of eudesmane type sesquiterpene ketone from Prangos heyniae H.Duman & Duman & M.F.Watson essential oil and mosquitocidal activity of the essential oils. Open Chemistry, 2018, 16, 453-467. | 1.0 | 15 |
| 24 | Chemical Composition and Antibacterial Activity of Essential Oils from Ferula L. Species against Methicillin-Resistant Staphylococcus aureus. Molecules, 2018, 23, 1679. | 1.7 | 46 |
| 25 | Chemical Compositions of Achillea sivasica: Different Plant Part Volatiles, Enantiomers and Fatty Acids. Records of Natural Products, 2018, 12, 142-159. | 1.3 | 9 |
| 26 | Fatty acid composition and anticancer activity in colon carcinoma cell lines of <i>Prunus dulcis</i> seed oil. Pharmaceutical Biology, 2017, 55, 1239-1248. | 1.3 | 53 |
| 27 | Chemical composition and phagocyte immunomodulatory activity of <i>Ferula iliensis</i> oils. Journal of Leukocyte Biology, 2017, 101, 1361-1371. | 1.5 | 30 |
| 28 | Essential oil composition and leaf trichomes of Pegolettia baccharidifolia and Pegolettia retrofracta (Asteraceae). South African Journal of Botany, 2017, 111, 275-282. | 1.2 | 0 |
| 29 | Phytochemicals, antioxidant, and antityrosinase activities of Achillea sivasica Çelik and Akpulat. International Journal of Food Properties, 2017, 20, S693-S706. | 1.3 | 14 |
| 30 | Chemical and Biological Diversity of the Leaf and Rhizome Volatiles of Acorus calamus L. from Turkey. Journal of Essential Oil-bearing Plants: JEOP, 2017, 20, 646-661. | 0.7 | 4 |
| 31 | Composition and potential of <i>Tanacetum haussknechtii</i> Bornm. Grierson as antioxidant and inhibitor of acetylcholinesterase, tyrosinase, and î±-amylase enzymes. International Journal of Food Properties, 2017, 20, S2359-S2378. | 1.3 | 10 |
| 32 | Investigation of <i>Galatella villosa</i> and <i>G. tatarica</i> for Antioxidant, α-Amylase, Tyrosinase, Lipoxygenase and Xanthine Oxidase Inhibitory Activities. Natural Product Communications, 2017, 12, 1934578X1701200. | 0.2 | 1 |
| 33 | Chemical Composition, Antioxidant and Anticholinesterase Activities of the Essential oil of Origanum rotundifolium Boiss. from Turkey. Records of Natural Products, 2017, 11, 485-490. | 1.3 | 7 |
| 34 | 11-Hydroxy-2,4-cycloeudesmane from the Leaf Oil of Juglans regia and Evaluation of its Larvicidal Activity. Natural Product Communications, 2016, 11, 1934578X1601101. | 0.2 | 2 |
| 35 | Preparative Capillary GC for Characterization of Five <i>Dracocephalum</i> Essential Oils from Mongolia, and their Mosquito Larvicidal Activity. Natural Product Communications, 2016, 11, 1934578X1601101. | 0.2 | 3 |
| 36 | Modulation of Human Neutrophil Responses by the Essential Oils from <i>Ferula akitschkensis</i> and Their Constituents. Journal of Agricultural and Food Chemistry, 2016, 64, 7156-7170. | 2.4 | 36 |

3

| # | Article | IF | CITATIONS |
|----|---|-----------|----------------------|
| 37 | Investigation of Essential Oils from Three Natural Populations of Lonicera iliensis. Chemistry of Natural Compounds, 2016, 52, 751-753. | 0.2 | 4 |
| 38 | Essential Oils of Echinophora lamondiana (Apiales: Umbelliferae): A Relationship Between Chemical Profile and Biting Deterrence and Larvicidal Activity Against Mosquitoes (Diptera: Culicidae). Journal of Medical Entomology, 2015, 52, 93-100. | 0.9 | 25 |
| 39 | Inhibition of Human Neutrophil Responses by the Essential Oil of <i>Artemisia kotuchovii</i> and Its Constituents. Journal of Agricultural and Food Chemistry, 2015, 63, 4999-5007. | 2.4 | 28 |
| 40 | Chemical composition and bioactivity studies of Alpinia nigra essential oils. Industrial Crops and Products, 2014, 53, 111-119. | 2.5 | 23 |
| 41 | Essential oils of i> Mentha i> species from Marmara region of Turkey. Journal of Essential Oil Research, 2012, 24, 265-272. | 1.3 | 25 |
| 42 | Isolation of Natural Products by Preparative Gas Chromatography. Methods in Molecular Biology, 2012, 864, 275-300. | 0.4 | 10 |
| 43 | Analysis of Essential Oils and Fragrances by Gas Chromatography. , 2012, , 519-527. | | 2 |
| 44 | Oneâ€step multiple component isolation from the oil of <i><scp>C</scp>rinitaria tatarica</i> (Less.) <scp>S</scp> ojak by preparative capillary gas chromatography with characterization by spectroscopic and spectrometric techniques and evaluation of biological activity. Journal of Separation Science, 2012, 35, 650-660. | 1.3 | 23 |
| 45 | Chemical Diversity of Volatiles of <i>Teucrium orientale</i> L. var. <i>orientale</i> , var. <i>puberulens</i> , and var. <i>glabrescens</i> Determined by Simultaneous GCâ€FID and GC/MS Techniques. Chemistry and Biodiversity, 2012, 9, 1144-1154. | 1.0 | 7 |
| 46 | Gas chromatographic analysis of essential oils., 2012,, 675-682. | | 1 |
| 47 | Rare sesquiterpenes from South African Pteronia species. South African Journal of Botany, 2010, 76, 146-152. | 1.2 | 12 |
| 48 | Chemical composition of the wood and leaf oils from the "Clanwilliam Cedar―(Widdringtonia) Tj ETQq0 0 0 652-654. | rgBT /Ove | erlock 10 Tf 5 28 |
| 49 | Gas chromatographic–mass spectrometric analysis of volatiles obtained by four different techniques from Salvia rosifolia Sm., and evaluation for biological activity. Journal of Chromatography A, 2010, 1217, 741-748. | 1.8 | 83 |
| 50 | Component composition of essential oils of Artemisia lercheana and A. sieversiana of the flora of Kazakhstan. Antimicrobial activity of A. sieversiana essential oil. Chemistry of Natural Compounds, 2009, 45, 120-123. | 0.2 | 12 |
| 51 | Studies on the Volatile Oils of i>Momordica charantia / i>L. (Cucurbitaceae) and i>Phyllanthus amarus / i>Sch. et Thonn (Euphorbiaceae). Journal of Essential Oil Research, 2009, 21, 393-399. | 1.3 | 23 |
| 52 | Composition of the Essential Oils of Calamintha tauricola P.H. Davis. Journal of Essential Oil Research, 2009, 21, 143-145. | 1.3 | 4 |
| 53 | Essential Oil Composition of <i> Gmelina arborea < /i > Roxb., Verbenaceae, From Nigeria. Journal of Essential Oil Research, 2009, 21, 264-266.</i> | 1.3 | 6 |
| 54 | The Needle Oil of <i>Pinus caribaea</i> Morelet From Nigeria. Journal of Essential Oil Research, 2009, 21, 342-344. | 1.3 | 4 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Bioactivity-Guided Fractionation and GC/MS Fingerprinting of <i>Angelica sinensis</i> and <i>Angelica archangelica</i> Root Components for Antifungal and Mosquito Deterrent Activity. Journal of Agricultural and Food Chemistry, 2009, 57, 464-470. | 2.4 | 95 |
| 56 | Component composition and antimicrobial activity of essential oil from Artemisia kasakorum. Chemistry of Natural Compounds, 2008, 44, 263-265. | 0.2 | 4 |
| 57 | Head-space volatiles of Gethyllis afra and G. ciliaris fruits ("kukumakrankaâ€). South African Journal of Botany, 2008, 74, 768-770. | 1.2 | 9 |
| 58 | Essential Oil of the Leaves of Ribes nigrum L. from Turkey. Journal of Essential Oil Research, 2008, 20, 512-514. | 1.3 | 10 |
| 59 | Composition and Antimicrobial Activity of the Oils of <i>Ferula szowitsiana </i> DC. from Turkey. Journal of Essential Oil Research, 2008, 20, 186-190. | 1.3 | 17 |
| 60 | Composition of the Essential Oils of <i>Angelica sylvestris</i> L. var. <i>sylvestris</i> Isolated from the Fruits by Different Isolation Techniques. Journal of Essential Oil Research, 2008, 20, 408-411. | 1.3 | 12 |
| 61 | Composition and Antimicrobial Activity of the Essential Oils of <i>Calamintha betulifolia </i> Boiss. et Bal Journal of Essential Oil Research, 2007, 19, 285-287. | 1.3 | 5 |
| 62 | Composition of the Essential Oils of <i>Tordylium trachycarpum</i> (Boiss.) Al-Eisawi et Jury and <i>Tordylium hasselquistiae</i> DC. Growing in Turkey. Journal of Essential Oil Research, 2007, 19, 410-412. | 1.3 | 3 |
| 63 | Composition and Antimicrobial Activity of the Essential Oil of <i>Tanacetum cadmeum </i> (Boiss.) Heywood subsp. <i>orientale </i> (I) Grierson. Journal of Essential Oil Research, 2007, 19, 392-395. | 1.3 | 14 |
| 64 | Comparison of Microwave-Assisted Hydrodistillation and Hydrodistillation Methods for the Fruit Essential Oils of <i>Foeniculum vulgare </i> . Journal of Essential Oil Research, 2007, 19, 426-429. | 1.3 | 21 |
| 65 | Composition of the Essential Oil ofHippomarathrum cristatum(DC.) Boiss Journal of Essential Oil Research, 2007, 19, 540-542. | 1.3 | 5 |
| 66 | Antimicrobial activities of methanol extracts and essential oils of Rosmarinus officinalis, depending on location and seasonal variations. Food Chemistry, 2007, 100, 553-559. | 4.2 | 421 |
| 67 | Seasonal and geographical variation of Heteropyxis natalensis essential oil and the effect thereof on the antimicrobial activity. South African Journal of Botany, 2007, 73, 441-448. | 1.2 | 44 |
| 68 | Comparison of hydrodistillation and microdistillation methods for the analysis of fruit volatiles of Prangos pabularia Lindl., and evaluation of its antimicrobial activity. South African Journal of Botany, 2007, 73, 563-569. | 1.2 | 27 |
| 69 | Composition of essential oils from Salvia anatolica, a new species endemic from Turkey. Chemistry of Natural Compounds, 2007, 43, 667-671. | 0.2 | 7 |
| 70 | Composition of the Essential Oil of Chaerophyllum macropodum Boiss. Fruits Obtained by Microdistillation. Journal of Essential Oil Research, 2006, 18, 515-517. | 1.3 | 15 |
| 71 | Comparison of Essential Oil ofXanthogalum purpurascensLallem. Obtained Via Different Isolation Techniques. Journal of Essential Oil Research, 2006, 18, 181-184. | 1.3 | 3 |
| 72 | Comparison of the Essential Oils of <i>Prangos turcica</i> A. Duran, M. Sagiroglu et H. Duman Fruits Obtained by Different Isolation Techniques. Journal of Essential Oil Research, 2006, 18, 511-514. | 1.3 | 15 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Composition of the Essential Oil of Diosma prama I. Williams. Journal of Essential Oil Research, 2006, 18, 17-18. | 1.3 | 0 |
| 74 | The Essential Oil Composition and Chemotaxonomical Appraisal of South African Pelargoniums (Geraniaceae). Journal of Essential Oil Research, 2006, 18, 89-105. | 1.3 | 26 |
| 75 | A Seasonal Variation Study of the Chemical Composition and Antimicrobial Activity of the Essential Oil of Agathosma ovata (Thunb.) Pillans (Rutaceae). Journal of Essential Oil Research, 2006, 18, 30-36. | 1.3 | 7 |
| 76 | Composition of the Essential Oils of Two <i>Adenandra</i> Species from South Africa. Journal of Essential Oil Research, 2006, 18, 46-47. | 1.3 | 5 |
| 77 | Composition of the Essential Oil of Euchaetis albertiniana I.J.M. Williams. Journal of Essential Oil Research, 2006, 18, 122-123. | 1.3 | 0 |
| 78 | Composition of the Essential Oils of Five <i>Coleonema</i> Species from South Africa. Journal of Essential Oil Research, 2006, 18, 26-29. | 1.3 | 7 |
| 79 | Composition of the Essential Oils of Three Acmadenia Species from South Africa. Journal of Essential Oil Research, 2006, 18, 54-56. | 1.3 | 1 |
| 80 | The Biological Activity and Essential Oil Composition of 17 <i>Agathosma</i> (Rutaceae) Species. Journal of Essential Oil Research, 2006, 18, 2-16. | 1.3 | 28 |
| 81 | Gas chromatographic–mass spectrometric analysis of essential oils from Pimpinella species gathered from Central and Northern Turkey. Journal of Chromatography A, 2006, 1117, 194-205. | 1.8 | 93 |
| 82 | Composition of the essential oil of Centaurea huber-morathii Wagenitz isolated from seeds by microdistillation. Flavour and Fragrance Journal, 2006, 21, 568-570. | 1.2 | 16 |
| 83 | Composition of the essential oils ofRhabdosciadium oligocarpum (Post ex Boiss.) Hedge et Lamond andRhabdosciadium microcalycinum HandMazz Flavour and Fragrance Journal, 2006, 21, 650-655. | 1.2 | 10 |
| 84 | A Preliminary Examination of the Composition of the Seed Oil of <i>Matthiola anchoniifolia </i> HubMor. Obtained by Microdistillation. Journal of Essential Oil Research, 2006, 18, 602-603. | 1.3 | 4 |
| 85 | A Simple Method to Obtain Essential Oils from Salvia triloba L. and Laurus nobilis L. by Using Microwave-assisted Hydrodistillation. Zeitschrift Fur Naturforschung - Section C Journal of Biosciences, 2005, 60, 501-504. | 0.6 | 34 |
| 86 | Composition of the Essential Oils of Juniperus oxycedrus subsp. Macrocarpa from Turkey. Chemistry of Natural Compounds, 2005, 41, 352-354. | 0.2 | 19 |
| 87 | Composition of the Essential Oil of <i>Cymbopogon afronardus</i> Stapf from Uganda. Journal of Essential Oil Research, 2005, 17, 139-140. | 1.3 | 2 |
| 88 | Enantiomeric Distribution of Linalool, Linalyl Acetate and Camphor in Bulgarian Lavender Oil. Journal of Essential Oil Research, 2005, 17, 135-136. | 1.3 | 9 |
| 89 | Comparison of Microwave-Assisted Hydrodistillation and Hydrodistillation Methods for the Analysis of Volatile Secondary Metabolites. Pharmaceutical Biology, 2005, 43, 491-495. | 1.3 | 31 |
| 90 | Comparison of the Essential Oils of Three Endemic TurkishHeracleumSpecies Obtained by Different Isolation Techniques. Journal of Essential Oil Research, 2005, 17, 605-610. | 1.3 | 23 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Composition of the Essential Oil of <i> Achillea sieheana </i> Stapf and the Enantiomeric Distribution of Camphor. Journal of Essential Oil Research, 2004, 16, 180-181. | 1.3 | 6 |
| 92 | A Comparative Study of the Essential Oils of Wild and Cultivated <i>Satureja hor tensis </i> L Journal of Essential Oil Research, 2004, 16, 422-424. | 1.3 | 48 |
| 93 | Essential oil composition and in vitro antimicrobial and anti-inflammatory activity of South African Vitex species. South African Journal of Botany, 2004, 70, 611-617. | 1.2 | 22 |
| 94 | Essential oils of three species of Heracleum. Anticandidal activity. Chemistry of Natural Compounds, 2004, 40, 544-547. | 0.2 | 32 |
| 95 | Comparison of the Essential Oils ofOriganum majoranaL. andOriganumxmajoricumCambess Journal of Essential Oil Research, 2004, 16, 248-252. | 1.3 | 33 |
| 96 | Composition of the Essential Oils of <i>Galium aparine</i> L. and <i>Galium odoratum</i> (L.) Scop. from Turkey. Journal of Essential Oil Research, 2004, 16, 305-307. | 1.3 | 17 |
| 97 | Isolation of Some Soluble and Dispersed Materials of Oregano Water. Chemistry of Natural Compounds, 2003, 39, 465-469. | 0.2 | 4 |
| 98 | The essential oil ofOriganum syriacumL. var.sinaicum(Boiss.) letswaart. Flavour and Fragrance Journal, 2003, 18, 98-99. | 1.2 | 31 |
| 99 | The essential oil ofAchillea falcataL Flavour and Fragrance Journal, 2003, 18, 192-194. | 1.2 | 15 |
| 100 | Composition of Essential Oils from Five Endemic <i>Sideritis</i> Species. Journal of Essential Oil Research, 2003, 15, 221-225. | 1.3 | 15 |
| 101 | Composition of the Microdistilled Essential Oils of Tordylium apulum L. and T. pustulosum Boiss Journal of Essential Oil Research, 2002, 14, 353-354. | 1.3 | 13 |
| 102 | Micro-Distilled Volatile Compounds from Ferulago Species Growing in Western Turkey. Pharmaceutical Biology, 2002, 40, 466-471. | 1.3 | 40 |
| 103 | Essential Oil of Arischrada korolkowii from the Chatkal Mountains of Uzbekistan. Chemistry of Natural Compounds, 2002, 38, 51-53. | 0.2 | 10 |
| 104 | Title is missing!. Chemistry of Natural Compounds, 2002, 38, 48-50. | 0.2 | 13 |
| 105 | Essential Oils of Two Hypericum Species from Uzbekistan. Chemistry of Natural Compounds, 2002, 38, 54-57. | 0.2 | 51 |
| 106 | Composition of the Essential Oil ofXanthogalum purpurascensLallem. Journal of Essential Oil Research, 2001, 13, 206-207. | 1.3 | 5 |
| 107 | Composition of the essential oils ofTanacetum armenum(DC.) Schultz Bip.,T. balsamitaL.,T. chiliophyllum(Fisch. & Mey.) Schultz Bip. var.chiliophyllumandT. haradjani(Rech. fil.) Grierson and the enantiomeric distribution of camphor and carvoneâ€. Flavour and Fragrance Journal, 2001, 16, 195-200. | 1.2 | 69 |
| 108 | ESSENTIAL OIL COMPOSITION OF THREE SPECIES OF Achillea FROM KAZAKHSTAN. Chemistry of Natural Compounds, 2001, 37, 447-450. | 0.2 | 22 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 109 | Title is missing!. Chemistry of Natural Compounds, 2001, 37, 238-241. | 0.2 | 17 |
| 110 | Composition of the Essential Oil ofNepeta betonicifoliaC.A. Meyer from Turkey. Journal of Essential Oil Research, 2001, 13, 35-36. | 1.3 | 16 |
| 111 | Composition of the essential oil ofGlaucosciadium cordifolium (Boiss.) Burtt et Davis from Turkey. Flavour and Fragrance Journal, 2000, 15, 45-46. | 1.2 | 11 |
| 112 | Composition of the essential oil of Prangos heyniae H. Duman et M. F. Watson, a new endemic from Turkey. Flavour and Fragrance Journal, 2000, 15, 47-49. | 1,2 | 24 |
| 113 | Composition of the essential oils of Zosima absinthifolia (Vent.) Link and Ferula elaeochytris Korovin from Turkey. Flavour and Fragrance Journal, 2000, 15, 371-372. | 1.2 | 44 |
| 114 | Lipids of Origanum tytthanthum. Chemistry of Natural Compounds, 2000, 36, 124-127. | 0.2 | 6 |
| 115 | Lipids and essential oil ofOriganum onites. Chemistry of Natural Compounds, 2000, 36, 132-136. | 0.2 | 15 |
| 116 | Production of essential oil from Cumin seeds. Chemistry of Natural Compounds, 2000, 36, 265-268. | 0.2 | 38 |
| 117 | Composition of the Essential Oil of <i>Nepeta fissa</i> C.A.Meyer. Journal of Essential Oil Research, 2000, 12, 27-28. | 1.3 | 13 |
| 118 | Essential Oils of Annual Sideritis Species Growing in Turkey. Pharmaceutical Biology, 2000, 38, 106-111. | 1.3 | 35 |
| 119 | Betulenols from Betula Species. Planta Medica, 2000, 66, 490-493. | 0.7 | 16 |
| 120 | Chemical Composition of Santolina chamaecyparissus L. Essential Oil. Journal of Essential Oil Research, 2000, 12, 625-627. | 1.3 | 24 |
| 121 | Steam Volatiles of <i>Lallemantia peltata </i> (L.) Fisch. et Mey. from Turkey. Journal of Essential Oil Research, 2000, 12, 689-690. | 1.3 | 9 |
| 122 | Chemical Composition of Turkish Myrtle Oil. Journal of Essential Oil Research, 2000, 12, 541-544. | 1.3 | 69 |
| 123 | Essential Oil of <i>Crithmum maritimum </i> L. from Turkey. Journal of Essential Oil Research, 2000, 12, 424-426. | 1.3 | 22 |
| 124 | Essential Oil of Hippomarathrum boissieri Reuter et Hausskn. Journal of Essential Oil Research, 2000, 12, 231-232. | 1.3 | 11 |
| 125 | The Analysis of Essential Oil and Headspace Volatiles of the Flowers ofPelargonium endlicherianumused as an Anthelmintic in Folk Medicine. Planta Medica, 1999, 65, 781-782. | 0.7 | 22 |
| 126 | Essential Oil ofPimpinella anisetumBoiss. et Bal Journal of Essential Oil Research, 1999, 11, 445-446. | 1.3 | 8 |

| # | Article | IF | CITATIONS |
|-----|--|------------------|-------------------|
| 127 | Composition of Essential Oils from Two Varieties of Ajuga chamaepityssubsp.chiafrom Turkey. Journal of Essential Oil Research, 1999, 11, 203-205. | 1.3 | 16 |
| 128 | Composition of the Essential Oils ofThymus leucostomusHausskn. et Velen var.gypsaceusJalas andThymus pubescensBoiss. et Kotschy ex Celak var.cratericolaJalas. Journal of Essential Oil Research, 1999, 11, 776-778. | 1.3 | 7 |
| 129 | Essential Oil ofCymbocarpum wiedemanniiBoiss Journal of Essential Oil Research, 1999, 11, 679-680. | 1.3 | 2 |
| 130 | Composition of essential oils from two endemicSideritis species of Turkey. Chemistry of Natural Compounds, 1999, 35, 61-64. | 0.2 | 11 |
| 131 | Composition of the essential oil ofPlumeria obtusa L Flavour and Fragrance Journal, 1999, 14, 237-240. | 1.2 | 5 |
| 132 | The Composition of Essential Oils from Tilial. Species Growing in Turkey. Journal of Essential Oil Research, 1999, 11, 369-374. | 1.3 | 20 |
| 133 | Composition of the Essential Oil of (i>Bunium persicum (i> (Boiss.) B. Fedtsch. from Tajikistan. Journal of Essential Oil Research, 1997, 9, 597-598. | 1.3 | 40 |
| 134 | Essential Oils ofMediasia macrophylla(Regel et Schmalh.) Pimen. andFoeniculum vulgareMill. from Uzbekistan. Journal of Essential Oil Research, 1997, 9, 249-250. | 1.3 | 16 |
| 135 | Essential Oil ofThymbra sintenisiiBornm. et Aznav. subsp.sintenisii. Journal of Essential Oil Research, 1997, 9, 355-356. | 1.3 | 4 |
| 136 | Composition of the Essential Oil ofThymus subcollinusKlokov from Turkey. Journal of Essential Oil Research, 1997, 9, 105-106. | 1.3 | 2 |
| 137 | Essential Oils ofCalamintha pamphylicaBoiss. et Heldr. subsp.pamphylicaand subsp.davisii(Quezel et) Tj ETQq1 1 (|).784314 | rgBT /Overl |
| 138 | Essential oils of some artemisia species from Central Asia. Chemistry of Natural Compounds, 1997, 33, 293-295. | 0.2 | 14 |
| 139 | Essential Oil of <i>Cyclotrichium origanifolium </i> (Labill.) Manden. et Scheng. from Turkey. Journal of Essential Oil Research, 1996, 8, 569-570. | 1.3 | 20 |
| 140 | Composition of Essential Oils from Three Varieties of <i>Thymus praecox</i> Opiz Growing in Turkey. Journal of Essential Oil Research, 1996, 8, 319-321. | 1.3 | 15 |
| 141 | Constituents of the Essential Oil ofRuta chalepensisL. from Turkey. Journal of Essential Oil Research, 1996, 8, 413-414. | 1.3 | 32 |
| 142 | Essential Oil Composition of Three Labiatae Endemic to Turkey (<i>Micromeria fruticosa</i> (L.) Druce) Tj ETQq0 C | 0 rgBT /0 1.3 | Overlock 10 18 |
| 143 | Essential Oil of <i>Echinophora chrysantha </i> Freyn et Sint Journal of Essential Oil Research, 1996, 8, 433-434. | 1.3 | 9 |
| 144 | Essential Oils ofThymbra sintenisiiBornm. et Aznav. subsp.isauricaP.H. Davis andOriganum leptocladumBoiss Journal of Essential Oil Research, 1996, 8, 675-676. | 1.3 | 10 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 145 | Composition of the Essential Oil of (i) Aloysia triphylla (li) (L'Herit.) Britton Grown in Turkey. Journal of Essential Oil Research, 1996, 8, 581-583. | 1.3 | 23 |
| 146 | Essential Oil ofPimpinella aromaticaBieb. from Turkey. Journal of Essential Oil Research, 1996, 8, 463-464. | 1.3 | 18 |
| 147 | Essential Oil of <i>Origanum micranthum </i> Vogel Journal of Essential Oil Research, 1996, 8, 203-204. | 1.3 | 11 |
| 148 | Essential Oil of <i>Origanum laevigatum < /i> Boiss Journal of Essential Oil Research, 1996, 8, 185-186.</i> | 1.3 | 13 |
| 149 | Composition of the Essential Oil from Fruits of <i>Scaligeria lazica </i> Boiss Journal of Essential Oil Research, 1995, 7, 557-558. | 1.3 | 6 |
| 150 | Composition of the Essential Oil ofHeracleum platytaeniumBoiss. from Turkey. Journal of Essential Oil Research, 1995, 7, 69-70. | 1.3 | 9 |
| 151 | Essential Oil of <i>Origanum saccatum</i> P. H. Davis. Journal of Essential Oil Research, 1995, 7, 175-176. | 1.3 | 14 |
| 152 | Constituents of the Essential Oil of Achillea biebersteinii Afan Journal of Essential Oil Research, 1995, 7, 527-528. | 1.3 | 29 |
| 153 | Composition of the Essential Oil of (i) Ocimum basilicum (i) L. Cultivated in Turkey. Journal of Essential Oil Research, 1995, 7, 203-205. | 1.3 | 22 |
| 154 | Composition of the Essential Oil ofCoridothymus capitatus(L.) Reichb. fil. from Turkey. Journal of Essential Oil Research, 1995, 7, 309-312. | 1.3 | 10 |
| 155 | Constituents of the Essential Oil from the Hulls of <i>Pistacia vera < /i>L Journal of Essential Oil Research, 1995, 7, 441-442.</i> | 1.3 | 31 |
| 156 | Essential Oil ofThymus thracicusVelen var.longidens(Velen) Jalas. Journal of Essential Oil Research, 1995, 7, 661-662. | 1.3 | 11 |
| 157 | Composition of Cold-Pressed Bergamot Oil from Turkey. Journal of Essential Oil Research, 1995, 7, 341-342. | 1.3 | 5 |
| 158 | Composition of the Essential Oil of <i>Nepeta viscida</i> Boiss. from Turkey. Journal of Essential Oil Research, 1995, 7, 569-570. | 1.3 | 12 |
| 159 | Essential Oil ofSalvia caespitosaMontbret et Aucher ex Benth Journal of Essential Oil Research, 1995, 7, 229-230. | 1.3 | 13 |
| 160 | Essential Oil of <i>Origanum rotundifolium </i> Boiss Journal of Essential Oil Research, 1995, 7, 95-96. | 1.3 | 16 |
| 161 | Essential Oil of <i>Thymus sipyleus</i> Boiss. subsp. <i>sipyleus</i> Var. <i>sipyleus</i> Journal of Essential Oil Research, 1995, 7, 411-413. | 1.3 | 7 |
| 162 | Essential Oil of <i>Micromeria carminea </i> P.H. Davis. Journal of Essential Oil Research, 1995, 7, 457-458. | 1.3 | 13 |

| # | Article | IF | CITATIONS |
|-----|---|-----|------------|
| 163 | Composition of the Essential Oil of (i) Salvia cryptantha (i) Montbret et Aucher ex Benth. from Turkey. Journal of Essential Oil Research, 1995, 7, 113-114. | 1.3 | 31 |
| 164 | Composition of the Essential Oil from Viburnum orientale Pallas Leaves. Journal of Essential Oil Research, 1995, 7, 321-323. | 1.3 | 3 |
| 165 | The Essential Oil ofOriganum vulgaresubsp.hirtumof Turkish Origin. Journal of Essential Oil Research, 1994, 6, 31-36. | 1.3 | 7 3 |
| 166 | Composition of the Root Oil of (i) Orthurus heterocarpus (i) (Boiss.) Juz Journal of Essential Oil Research, 1994, 6, 349-351. | 1.3 | 0 |
| 167 | Essential Oil of <i>Sideritis hispida</i> P. H. Davis, an Endemic Species from Turkey. Journal of Essential Oil Research, 1994, 6, 435-436. | 1.3 | 12 |
| 168 | Composition of the Essential Oil ofNepeta caesareaBoiss. from Turkey. Journal of Essential Oil Research, 1994, 6, 645-646. | 1.3 | 16 |
| 169 | Essential Oil ofOriganum solymicumP. H. Davis. Journal of Essential Oil Research, 1994, 6, 503-504. | 1.3 | 10 |
| 170 | Essential Oil of Echinophora tenuifolial. subsp.sibthorpiana (Guss.) Tutin. Journal of Essential Oil Research, 1994, 6, 399-400. | 1.3 | 20 |
| 171 | The Essential Oil Composition ofDictamnus albusfrom Turkey. Planta Medica, 1994, 60, 481-482. | 0.7 | 9 |
| 172 | Composition of Essential Oils from Two Varieties of <i>Thymbra spicata </i> L Journal of Essential Oil Research, 1994, 6, 463-468. | 1.3 | 36 |
| 173 | The composition of Manila elemi oil. Flavour and Fragrance Journal, 1993, 8, 35-37. | 1.2 | 26 |
| 174 | Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. Journal of Essential Oil Research, 1993, 5, 425-431. | 1.3 | 62 |
| 175 | The Essential Oil of <i>Scaligeria lazica</i> Boiss Journal of Essential Oil Research, 1993, 5, 463-464. | 1.3 | 8 |
| 176 | The Essential Oil of <i>Laser trilobum</i> Fruit of Turkish Origin. Journal of Essential Oil Research, 1993, 5, 365-369. | 1.3 | 14 |
| 177 | The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl. subsp.longicaulis in the Same Population. Journal of Essential Oil Research, 1993, 5, 291-295. | 1.3 | 23 |
| 178 | The Essential Oil of Thymus bornmuelleri Velen Journal of Essential Oil Research, 1993, 5, 691-692. | 1.3 | 4 |
| 179 | Composition of the Essential Oil ofCalamintha grandiflora. Planta Medica, 1993, 59, 390-390. | 0.7 | 12 |
| 180 | The Essential Oil of <i>Salvia pomifera</i> L Journal of Essential Oil Research, 1993, 5, 347-348. | 1.3 | 32 |

| # | Article | IF | CITATION |
|-----|---|-----|----------|
| 181 | The Essential Oil of <i>Sideritis athoa</i> Papanikolaou et Kokkini. Journal of Essential Oil Research, 1993, 5, 669-670. | 1.3 | 14 |
| 182 | The Essential Oil of <i>Micromeria fruticosa</i> (L.) Druce subsp. <i>serpyllifolia</i> (Bieb.) P. H. Davis. Journal of Essential Oil Research, 1993, 5, 199-200. | 1.3 | 11 |
| 183 | Composition of the Essential Oil of Nepeta racemosa Lam Journal of Essential Oil Research, 1993, 5, 215-217. | 1.3 | 30 |
| 184 | The Essential Oils of <i>Rhus coriaria </i> L. (Sumac). Journal of Essential Oil Research, 1993, 5, 481-486. | 1.3 | 32 |
| 185 | Composition of the Essential Oils of TurkishOriganumSpecies with Commercial Importance. Journal of Essential Oil Research, 1993, 5, 619-623. | 1.3 | 118 |
| 186 | BIOLOGICAL ACTIVITY DETERMINATION OF BLACK AND WHITE CHIA SEED EXTRACTS OBTAINED BY DIFFERENT EXTRACTION METHODS. Ankara Universitesi Eczacilik Fakultesi Dergisi, 0, , . | 0.2 | 0 |