Masayuki Murata

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9221750/publications.pdf

Version: 2024-02-01

	759233	752698
537	12	20
citations	h-index	g-index
50	50	870
30	30	670
docs citations	times ranked	citing authors
	citations 50	537 12 citations h-index 50 50

#	Article	IF	CITATIONS
1	A comparison of the antitumor effects of interferon- $\hat{l}\pm$ and \hat{l}^2 on human hepatocellular carcinoma cell lines. Cytokine, 2006, 33, 121-128.	3.2	52
2	Glycated albumin as a diagnostic tool for diabetes in a general Japanese population. Metabolism: Clinical and Experimental, 2015, 64, 698-705.	3.4	42
3	Study Profile of the Japan Multi-institutional Collaborative Cohort (J-MICC) Study. Journal of Epidemiology, 2021, 31, 660-668.	2.4	41
4	Potential risk of <scp>HBV</scp> reactivation in patients with resolved <scp>HBV</scp> infection undergoing directâ€acting antiviral treatment for <scp>HCV</scp> . Liver International, 2018, 38, 76-83.	3.9	37
5	Subclinical carotid atherosclerosis and triglycerides predict the incidence of chronic kidney disease in the Japanese general population: Results from the Kyushu and Okinawa Population Study (KOPS). Atherosclerosis, 2015, 238, 207-212.	0.8	33
6	Predictors of kidney tubular dysfunction induced by adefovir treatment for chronic hepatitis B. World Journal of Gastroenterology, 2015, 21, 2116-2123.	3.3	25
7	Higher dietary cholesterol and ï‰-3 fatty acid intakes are associated with a lower success rate of Helicobacter pylori eradication therapy in Japan. American Journal of Clinical Nutrition, 2017, 106, 581-588.	4.7	23
8	Association of Dietary Acid Load with the Prevalence of Metabolic Syndrome among Participants in Baseline Survey of the Japan Multi-Institutional Collaborative Cohort Study. Nutrients, 2020, 12, 1605.	4.1	23
9	Clinical and pathogenic features of SCCmec type II and IV methicillin-resistant Staphylococcus aureus in Japan. Journal of Infection and Chemotherapy, 2017, 23, 90-95.	1.7	20
10	Secondary Syphilis with Pulmonary Involvement. Internal Medicine, 2018, 57, 121-126.	0.7	19
11	The utility of urinary myo-inositol as a marker of glucose intolerance. Diabetes Research and Clinical Practice, 2014, 103, 88-96.	2.8	14
12	Sedentary Time is Associated with Cardiometabolic Diseases in A Large Japanese Population: A Cross-Sectional Study. Journal of Atherosclerosis and Thrombosis, 2020, 27, 1097-1107.	2.0	14
13	Correlation between Thyroid Stimulating Hormone and Renal Function in Euthyroid Residents of Japan: Results from the Kyushu and Okinawa Population Study (KOPS). Journal of Atherosclerosis and Thrombosis, 2018, 25, 335-343.	2.0	13
14	Small Dense Low-Density Lipoprotein Cholesterol and Carotid Intimal Medial Thickness Progression. Journal of Atherosclerosis and Thrombosis, 2020, 27, 1108-1122.	2.0	13
15	Association of genetic risk score and chronic kidney disease in a Japanese population. Nephrology, 2019, 24, 670-673.	1.6	12
16	HIV infection with concomitant cerebral toxoplasmosis and disseminated histoplasmosis in a 45-year-old man. Journal of Infection and Chemotherapy, 2007, 13, 51-55.	1.7	11
17	Association of IL28B rs8099917 genotype and female sex with spontaneous clearance of hepatitis C virus infection: a Japanese cross-sectional study. Archives of Virology, 2016, 161, 641-648.	2.1	11
18	The Expression Level of Neutrophil CD64 Is a Useful Marker of Systemic Inflammation Associated with HIV Infection. AIDS Research and Human Retroviruses, 2017, 33, 147-156.	1.1	10

#	Article	IF	CITATIONS
19	The Relation between Serum Endostatin Level and Carotid Atherosclerosis in Healthy Residents of Japan: Results from the Kyushu and Okinawa Population Study (KOPS). Journal of Atherosclerosis and Thrombosis, 2017, 24, 1023-1030.	2.0	9
20	Prevalence and characteristics of occult hepatitis B virus infection in Japanese human immunodeficiency virus-infected patients. Journal of Infection and Chemotherapy, 2020, 26, 28-32.	1.7	9
21	Impact of <i>PSCA</i> Polymorphisms on the Risk of Duodenal Ulcer. Journal of Epidemiology, 2021, 31, 12-20.	2.4	9
22	Comparison of the Abbott RealTime HCV and Roche COBAS Ampliprep/COBAS TaqMan HCV Assays for the Monitoring of Sofosbuvir-Based Therapy. Antiviral Therapy, 2017, 22, 61-70.	1.0	8
23	A genome-wide association study in Japanese identified one variant associated with a preference for a Japanese dietary pattern. European Journal of Clinical Nutrition, 2021, 75, 937-945.	2.9	8
24	Moderate-to-vigorous Physical Activity and Sedentary Behavior Are Independently Associated With Renal Function: A Cross-sectional Study. Journal of Epidemiology, 2023, 33, 285-293.	2.4	7
25	A genome-wide association study on confection consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort Study. British Journal of Nutrition, 2021, 126, 1843-1851.	2.3	6
26	A case of successful treatment with telaprevir-based triple therapy for hepatitis C infection after treatment failure with vaniprevir-based triple therapy. Journal of Infection and Chemotherapy, 2014, 20, 577-581.	1.7	5
27	The kinetics of the hepatitis B surface antigen level after the initiation of antiretroviral therapy for hepatitis B virus and human immunodeficiency virus coinfected patients. Journal of Infection and Chemotherapy, 2015, 21, 264-271.	1.7	5
28	Toxocariasis Suspected of Having Infiltrated Directly from the Liver to the Lung through the Diaphragm. Internal Medicine, 2019, 58, 2737-2741.	0.7	5
29	Differential Effect of Polymorphisms on Body Mass Index Across the Life Course of Japanese: The Japan Multi-Institutional Collaborative Cohort Study. Journal of Epidemiology, 2021, 31, 172-179.	2.4	5
30	A genome-wide association study on fish consumption in a Japanese populationâ€"the Japan Multi-Institutional Collaborative Cohort study. European Journal of Clinical Nutrition, 2021, 75, 480-488.	2.9	5
31	Long-term effects of lamivudine treatment in Japanese chronic hepatitis B patients. World Journal of Gastroenterology, 2011, 17, 2945.	3.3	5
32	Associations of breastfeeding history with metabolic syndrome and cardiovascular risk factors in community-dwelling parous women: The Japan Multi-Institutional Collaborative Cohort Study. PLoS ONE, 2022, 17, e0262252.	2.5	5
33	A case of successful hepatitis C virus eradication by 24 weeks of telaprevir-based triple therapy for a hemophilia patient with hepatitis C virus/human immunodeficiency virus co-infection who previously failed pegylated interferon-l± and ribavirin therapy. Journal of Infection and Chemotherapy, 2014, 20, 320-324.	1.7	4
34	Impact of HCV kinetics on treatment outcome differs by the type of real-time HCV assay in NS3/4A protease inhibitor-based triple therapy. Antiviral Research, 2016, 126, 35-42.	4.1	4
35	Direct-acting antiviral-based triple therapy on alpha-fetoprotein level in chronic hepatitis C patients. World Journal of Gastroenterology, 2015, 21, 4696-4706.	3.3	4
36	The relation of postprandial plasma glucose and serum endostatin to the urinary albumin excretion of residents with prediabetes: results from the Kyushu and Okinawa Population Study (KOPS). International Urology and Nephrology, 2016, 48, 851-857.	1.4	3

#	Article	IF	CITATIONS
37	Influence of insulin resistance on the development of hepatocellular carcinoma after antiviral treatment for non-cirrhotic patients with chronic hepatitis C. Infectious Agents and Cancer, 2016, 11, 9.	2.6	3
38	Association of the Serum Endostatin Level, Renal Function, and Carotid Atherosclerosis of Healthy Residents of Japan: Results from the Kyushu and Okinawa Population Study (KOPS). Journal of Atherosclerosis and Thrombosis, 2018, 25, 829-835.	2.0	3
39	The efficacy and safety of hachimijiogan for mild Alzheimer disease in an exploratory, open standard treatment controlled, randomized allocation, multicenter trial. Medicine (United States), 2020, 99, e22370.	1.0	3
40	A case of severe COVID-19 with pulmonary thromboembolism related to heparin-induced thrombocytopenia during prophylactic anticoagulation therapy. Journal of Infection and Chemotherapy, 2022, , .	1.7	3
41	A case of lobar pneumonia and sepsis with death caused by invasive Klebsiella rhinoscleromatis infection. Journal of Infection and Chemotherapy, 2016, 22, 707-711.	1.7	2
42	Kyushu and Okinawa Population Study (KOPS): a large prospective cohort study in Japan. BMJ Open, 2021, 11, e053763.	1.9	2
43	Genome-wide association study of serum prostate-specific antigen levels based on 1000 Genomes imputed data in Japanese: the Japan Multi-Institutional Collaborative Cohort Study. Nagoya Journal of Medical Science, 2021, 83, 183-194.	0.3	1
44	Impact of the PNPLA3 genotype on the risk of hepatocellular carcinoma after hepatitis C virus eradication. Journal of Medical Virology, 2022, 94, 5007-5014.	5.0	1
45	Nocturnal Difference in Ultra-Low Frequency Band of the Heart Rate Variability of Patients Stratified by Kampo Medicine Prescription. Journal of Alternative and Complementary Medicine, 2014, 20, A91-A91.	2.1	0
46	A case of granulomatosis with polyangiitis preceded by subacute thyroiditis. Clinical Case Reports (discontinued), 2015, 3, 139-144.	0.5	0
47	Impact of Food Consumption on Helicobacter pylori Eradication Therapy: A Japanese Community-Based Observational Study. Open Forum Infectious Diseases, 2016, 3, .	0.9	0
48	A case of amebiasis with negative serologic markers that caused intra-abdominal abscess. Journal of Infection and Chemotherapy, 2017, 23, 778-781.	1.7	0
49	Population-Based Impact of Smoking, Drinking, and Genetic Factors on HDL-Cholesterol Levels in J-MICC Study Participants. Journal of Epidemiology, 2021, , .	2.4	0
50	Assessing the Relationship Between High-sensitivity C-reactive Protein and Kidney Function Employing Mendelian Randomization in the Japanese Community-based J-MICC Study. Journal of Epidemiology, 2021, , .	2.4	0

4