

Tsutomu Nakazawa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9234027/publications.pdf>

Version: 2024-02-01

17
papers

597
citations

840776

11
h-index

940533

16
g-index

17
all docs

17
docs citations

17
times ranked

842
citing authors

#	ARTICLE	IF	CITATIONS
1	AML1-ETO rapidly induces acute myeloblastic leukemia in cooperation with the Wilms tumor gene, WT1. <i>Blood</i> , 2006, 107, 3303-3312.	1.4	111
2	Overexpression of the Wilms' tumor gene WT1 in primary astrocytic tumors. <i>Cancer Science</i> , 2004, 95, 822-827.	3.9	110
3	Novel Human NK Cell Line Carrying CAR Targeting EGFRvIII Induces Antitumor Effects in Glioblastoma Cells. <i>Anticancer Research</i> , 2018, 38, 5049-5056.	1.1	82
4	Effect of CRISPR/Cas9-Mediated PD-1-Disrupted Primary Human Third-Generation CAR-T Cells Targeting EGFRvIII on In Vitro Human Glioblastoma Cell Growth. <i>Cells</i> , 2020, 9, 998.	4.1	64
5	Wilms's tumor gene WT1 17AA(-)/KTS(-) isoform induces morphological changes and promotes cell migration and invasion in vitro. <i>Cancer Science</i> , 2006, 97, 259-270.	3.9	61
6	CRISPR-Cas9-Mediated TIM3 Knockout in Human Natural Killer Cells Enhances Growth Inhibitory Effects on Human Glioma Cells. <i>International Journal of Molecular Sciences</i> , 2021, 22, 3489.	4.1	32
7	Ex vivo-expanded highly purified natural killer cells in combination with temozolomide induce antitumor effects in human glioblastoma cells in vitro. <i>PLoS ONE</i> , 2019, 14, e0212455.	2.5	31
8	Cytotoxic human peripheral blood-derived β^2 microglobulin T cells kill glioblastoma cell lines: implications for cell-based immunotherapy for patients with glioblastoma. <i>Journal of Neuro-Oncology</i> , 2014, 116, 31-39.	2.9	21
9	Antitumor effects of minodronate, a third-generation nitrogen-containing bisphosphonate, in synergy with β^2 microglobulin T cells in human glioblastoma in vitro and in vivo. <i>Journal of Neuro-Oncology</i> , 2016, 129, 231-241.	2.9	15
10	KHYG-1 Cells With EGFRvIII-specific CAR Induced a Pseudoprogression-like Feature in Subcutaneous Tumours Derived from Glioblastoma-like Cells. <i>Anticancer Research</i> , 2020, 40, 3231-3237.	1.1	15
11	Expression of peptide transporter 1 has a positive correlation in protoporphyrin IX accumulation induced by 5-aminolevulinic acid with photodynamic detection of non-small cell lung cancer and metastatic brain tumor specimens originating from non-small cell lung cancer. <i>Photodiagnosis and Photodynamic Therapy</i> , 2019, 25, 309-316.	2.6	12
12	Evaluation of Comprehensive Gene Expression and NK Cell-Mediated Killing in Glioblastoma Cell Line-Derived Spheroids. <i>Cancers</i> , 2021, 13, 4896.	3.7	12
13	The Wilms's tumor gene WT1 is a common marker of progenitor cells in fetal liver. <i>Biochemical and Biophysical Research Communications</i> , 2005, 326, 836-843.	2.1	11
14	Ex Vivo Expanded and Activated Natural Killer Cells Prolong the Overall Survival of Mice with Glioblastoma-like Cell-Derived Tumors. <i>International Journal of Molecular Sciences</i> , 2021, 22, 9975.	4.1	10
15	Risk of brain herniation after craniotomy with lumbar spinal drainage: a propensity score analysis. <i>Journal of Neurosurgery</i> , 2019, 130, 1710-1720.	1.6	8
16	Minodronic Acid in Combination with β^2 microglobulin T Cells Induces Apoptosis of Non-small Cell Lung Carcinoma Cell Lines. <i>Anticancer Research</i> , 2016, 36, 5883-5886.	1.1	1
17	Combined Treatment With Radiotherapy and Immunotherapy for Isocitrate Dehydrogenase Mutant Brainstem Glioma in Adult: A Case Report. <i>Brain Tumor Research and Treatment</i> , 2022, 10, 129.	1.0	1