

Takafumi Nakano

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9239448/publications.pdf>

Version: 2024-02-01

18
papers

371
citations

933447

10
h-index

839539

18
g-index

18
all docs

18
docs citations

18
times ranked

598
citing authors

#	ARTICLE	IF	CITATIONS
1	<i>GET4</i> is a novel driver gene in colorectal cancer that regulates the localization of BAG6, a nucleocytoplasmic shuttling protein. <i>Cancer Science</i> , 2022, 113, 156-169.	3.9	7
2	Retrospective Study of Cisplatin/Carboplatin, 5-Fluorouracil Plus Cetuximab (EXTREME) for Advanced-stage Salivary Gland Cancer. <i>In Vivo</i> , 2022, 36, 979-984.	1.3	4
3	Pan-tropomyosin receptor kinase immunoreactivity, ETV6-NTRK3 fusion subtypes, and RET rearrangement in salivary secretory carcinoma. <i>Human Pathology</i> , 2021, 109, 37-44.	2.0	9
4	The novel driver gene <i>ASAP2</i> is a potential druggable target in pancreatic cancer. <i>Cancer Science</i> , 2021, 112, 1655-1668.	3.9	18
5	p16 overexpression and Rb loss correlate with high-risk HPV infection in oropharyngeal squamous cell carcinoma. <i>Histopathology</i> , 2021, 79, 358-369.	2.9	12
6	Programmed Death-Ligand 1 Expression and Tumor-Infiltrating Lymphocytes in Temporal Bone Squamous Cell Carcinoma. <i>Laryngoscope</i> , 2021, 131, 2674-2683.	2.0	4
7	YAP1 is a potent driver of the onset and progression of oral squamous cell carcinoma. <i>Science Advances</i> , 2020, 6, eaay3324.	10.3	75
8	The clinical value of serum squamous cell carcinoma antigens 1 and 2 in head and neck squamous cell carcinoma. <i>Auris Nasus Larynx</i> , 2019, 46, 135-140.	1.2	9
9	Prognostic value of programmed death ligand-1 and ligand-2 co-expression in salivary gland carcinomas. <i>Oral Oncology</i> , 2019, 90, 30-37.	1.5	43
10	The treatment and outcome analysis of primary squamous cell carcinoma of the thyroid. <i>Auris Nasus Larynx</i> , 2018, 45, 553-557.	1.2	17
11	Combination of serum squamous cell carcinoma antigens 1 and 2 as potential diagnostic marker for sinonasal squamous cell carcinoma and inverted papilloma. <i>Head and Neck</i> , 2018, 40, 2583-2589.	2.0	6
12	Clinical management of squamous cell carcinoma associated with sinonasal inverted papilloma. <i>Auris Nasus Larynx</i> , 2017, 44, 98-103.	1.2	26
13	Low-grade intraductal carcinoma (low-grade cribriform cystadenocarcinoma) with tumor-associated lymphoid proliferation of parotid gland. <i>Pathology Research and Practice</i> , 2017, 213, 706-709.	2.3	14
14	Syndrome of inappropriate antidiuretic hormone secretion in a case of olfactory neuroblastoma without anti-diuretic hormone immunoreactivity: A case report and review of the literature. <i>Auris Nasus Larynx</i> , 2017, 44, 771-774.	1.2	10
15	Primary combined small cell carcinoma and squamous cell carcinoma of the oropharynx with special reference to EGFR status of small cell carcinoma component: Case report and review of the literature. <i>Auris Nasus Larynx</i> , 2017, 44, 472-478.	1.2	10
16	Molecular subclassification determined by human papillomavirus and epidermal growth factor receptor status is associated with the prognosis of oropharyngeal squamous cell carcinoma. <i>Human Pathology</i> , 2016, 50, 51-61.	2.0	28
17	Hyalinizing clear cell carcinoma with EWSR1-ATF1 fusion gene: report of three cases with molecular analyses. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2015, 466, 37-43.	2.8	32
18	<i>HER2</i> and <i>EGFR</i> gene copy number alterations are predominant in high-grade salivary mucoepidermoid carcinoma irrespective of <i>MAML2</i> fusion status. <i>Histopathology</i> , 2013, 63, 378-392.	2.9	47