

Hans J ,rgen Aarstad

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9246705/publications.pdf>

Version: 2024-02-01

39
papers

674
citations

567281

15
h-index

610901

24
g-index

39
all docs

39
docs citations

39
times ranked

845
citing authors

#	ARTICLE	IF	CITATIONS
1	Generic quality of life in persons with hearing loss: a systematic literature review. <i>BMC Ear, Nose and Throat Disorders</i> , 2018, 18, 1.	2.6	92
2	Exercise-induced laryngeal obstruction: natural history and effect of surgical treatment. <i>European Archives of Oto-Rhino-Laryngology</i> , 2011, 268, 1485-1492.	1.6	83
3	Mood, anxiety and sense of humor in head and neck cancer patients in relation to disease stage, prognosis and quality of life. <i>Acta Oto-Laryngologica</i> , 2005, 125, 557-565.	0.9	48
4	Congenital laryngomalacia is related to exercise-induced laryngeal obstruction in adolescence. <i>Archives of Disease in Childhood</i> , 2016, 101, 443-448.	1.9	41
5	Cross-Cultural Adaption and Translation of the Voice Handicap Index into Norwegian. <i>Folia Phoniatica Et Logopaedica</i> , 2012, 64, 234-240.	1.1	30
6	The impact of HPV infection on survival in a geographically defined cohort of oropharynx squamous cell carcinoma (OPSCC) patients in whom surgical treatment has been one main treatment. <i>Acta Oto-Laryngologica</i> , 2014, 134, 636-645.	0.9	29
7	Hearing as an Independent Predictor of Postural Balance in 1075 Patients Evaluated for Dizziness. <i>Otolaryngology - Head and Neck Surgery</i> , 2019, 161, 478-484.	1.9	27
8	Acoustic Voice Analysis and Maximum Phonation Time in Relation to Voice Handicap Index Score and Larynx Disease. <i>Journal of Voice</i> , 2020, 34, 161.e27-161.e35.	1.5	25
9	Personality Traits in Head and Neck Squamous Cell Carcinoma Patients in Relation to the Disease State, Disease Extent and Prognosis. <i>Acta Oto-Laryngologica</i> , 2002, 122, 892-899.	0.9	21
10	Presence of activated T lymphocytes in peripheral blood of head and neck squamous cell carcinoma patients predicts impaired prognosis. <i>Acta Oto-Laryngologica</i> , 2006, 126, 1326-1333.	0.9	19
11	Functional-structural reorganisation of the neuronal network for auditory perception in subjects with unilateral hearing loss: Review of neuroimaging studies. <i>Hearing Research</i> , 2016, 332, 73-79.	2.0	19
12	The impact of HPV infection, smoking history, age and operability of the patient on disease-specific survival in a geographically defined cohort of patients with oropharyngeal squamous cell carcinoma. <i>Acta Oto-Laryngologica</i> , 2014, 134, 964-973.	0.9	18
13	Health-related Quality of Life as Studied by EORTC QLQ and Voice Handicap Index Among Various Patients With Laryngeal Disease. <i>Journal of Voice</i> , 2017, 31, 251.e17-251.e26.	1.5	17
14	Primary surgery results in no survival benefit compared to primary radiation for oropharyngeal cancer patients stratified by high-risk human papilloma virus status. <i>European Archives of Oto-Rhino-Laryngology</i> , 2017, 274, 477-487.	1.6	17
15	Head and neck squamous cell carcinoma spheroid- and monocyte spheroid-stimulated IL-6 and monocyte chemotactic protein-1 secretion are related to TNM stage, inflammatory state and tumor macrophage density. <i>Acta Oto-Laryngologica</i> , 2005, 125, 1097-1104.	0.9	15
16	Level of distress predicts subsequent survival in successfully treated head and neck cancer patients: a prospective cohort study. <i>Acta Oto-Laryngologica</i> , 2014, 134, 211-219.	0.9	14
17	Mononuclear phagocytes in head and neck squamous cell carcinoma. <i>European Archives of Oto-Rhino-Laryngology</i> , 2010, 267, 335-344.	1.6	13
18	Peripheral blood monocyte and T lymphocyte activation levels at diagnosis predict long-term survival in head and neck squamous cell carcinoma patients. <i>Apmis</i> , 2015, 123, 305-314.	2.0	13

#	ARTICLE	IF	CITATIONS
19	Prediction of 5 year survival from level of perceived distress in newly diagnosed head and neck squamous cell carcinoma patients. <i>Oral Oncology</i> , 2013, 49, 964-969.	1.5	12
20	General health-related quality of life scores from head and neck squamous cell carcinoma patients obtained throughout the first year following diagnosis predicted up to 10-year overall survival. <i>European Archives of Oto-Rhino-Laryngology</i> , 2018, 275, 207-217.	1.6	12
21	In vitro cholesteatoma growth and secretion of cytokines. <i>Acta Oto-Laryngologica</i> , 2010, 130, 815-819.	0.9	11
22	Tumor stromal desmoplasia and inflammatory response uniquely predict survival with and without stratification for HPV tumor infection in OPSCC patients. <i>Acta Oto-Laryngologica</i> , 2018, 138, 1035-1042.	0.9	11
23	The Norwegian Voice Handicap Index (VHI-N) patient scores are dependent on voice-related disease group. <i>European Archives of Oto-Rhino-Laryngology</i> , 2015, 272, 2897-2905.	1.6	9
24	The Acute Phase Reaction and Its Prognostic Impact in Patients with Head and Neck Squamous Cell Carcinoma: Single Biomarkers Including C-Reactive Protein Versus Biomarker Profiles. <i>Biomedicines</i> , 2020, 8, 418.	3.2	9
25	Tumor Infiltration Levels of CD3, Foxp3 (+) Lymphocytes and CD68 Macrophages at Diagnosis Predict 5-Year Disease-Specific Survival in Patients with Oropharynx Squamous Cell Carcinoma. <i>Cancers</i> , 2022, 14, 1508.	3.7	9
26	Clinical Application and Psychometric Properties of a Norwegian Questionnaire for the Self-Assessment of Communication in Quiet and Adverse Conditions Using Two Revised APHAB Subscales. <i>Journal of the American Academy of Audiology</i> , 2018, 29, 025-034.	0.7	8
27	An fMRI-study on single-sided deafness: Spectral-temporal properties and side of stimulation modulates hemispheric dominance. <i>NeuroImage: Clinical</i> , 2019, 24, 101969.	2.7	8
28	EORTC Quality of Life Questionnaire Head and Neck (H&N)-35 scores from H&N squamous cell carcinoma patients obtained at diagnosis and at 6, 9 and 12 months following diagnosis predict 10-year overall survival. <i>European Archives of Oto-Rhino-Laryngology</i> , 2019, 276, 3495-3505.	1.6	8
29	Importance of personality and coping expectancy on patient-reported hearing disability, quality of life and distress level: a study of patients referred to an audiology service. <i>Health and Quality of Life Outcomes</i> , 2021, 19, 168.	2.4	6
30	In vitro Monocyte Secretion Levels Following Stimulation with Autologous Spheroids Derived from Tumour or Benign Mucosa Predict Long-term Survival in Head and Neck Squamous Cell Carcinoma Patients. <i>Scandinavian Journal of Immunology</i> , 2017, 85, 211-219.	2.7	5
31	Reduced grey- and white matter volumes due to unilateral hearing loss following treatment for vestibular schwannoma. <i>Heliyon</i> , 2020, 6, e05658.	3.2	5
32	Quality of life in persons with hearing loss: a study of patients referred to an audiological service. <i>International Journal of Audiology</i> , 2019, 58, 696-703.	1.7	4
33	Periodontal status at diagnosis predicts non-disease-specific survival in a geographically defined cohort of patients with oropharynx squamous cell carcinoma. <i>Acta Oto-Laryngologica</i> , 2019, 139, 309-315.	0.9	4
34	Evaluation of Self-reported Symptoms in 1,457 Dizzy Patients and Associations With Caloric Testing and Posturography. <i>Otology and Neurotology</i> , 2020, 41, 956-963.	1.3	4
35	Perioperative Cytokine Response to Infection Associated With Elective Arterial Surgery. <i>Vascular and Endovascular Surgery</i> , 2014, 48, 116-122.	0.7	3
36	Psychometric properties of the Norwegian translation of the Tinnitus Handicap Inventory (THI-NOR). <i>International Journal of Audiology</i> , 2021, , 1-6.	1.7	3

#	ARTICLE	IF	CITATIONS
37	Long-term Survival in 1,931 Patients With Dizziness: Disease- and Symptom-Specific Mortality. Laryngoscope, 2021, 131, E2031-E2037.	2.0	2
38	Psychometric properties for the Norwegian translations of two revised APHAB-subcales and an adapted IOI-HA (IOI-CI) in patients with cochlear implants. International Journal of Audiology, 2021, , 1-8.	1.7	0
39	Abstract 2520: Detection of transcriptional active HPV 16/18 in patients with oropharyngeal squamous cell carcinoma by dual immunohistochemistry p16INK4 and in situ hybridization E6/E7 mRNA in archival material older than 25 years. , 2021, , .		0