
## Hans JÃ, rgen Aarstad

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9246705/publications.pdf

Version: 2024-02-01



#	Article	IF	CITATIONS
1	Tumor Infiltration Levels of CD3, Foxp3 (+) Lymphocytes and CD68 Macrophages at Diagnosis Predict 5-Year Disease-Specific Survival in Patients with Oropharynx Squamous Cell Carcinoma. Cancers, 2022, 14, 1508.	3.7	9
2	Longâ€Term Survival in 1,931 Patients With Dizziness: Disease―and Symptomâ€&pecific Mortality. Laryngoscope, 2021, 131, E2031-E2037.	2.0	2
3	Psychometric properties of the Norwegian translation of the Tinnitus Handicap Inventory (THI-NOR). International Journal of Audiology, 2021, , 1-6.	1.7	3
4	Psychometric properties for the Norwegian translations of two revised APHAB-subscales and an adapted IOI-HA (IOI-CI) in patients with cochlear implants. International Journal of Audiology, 2021, , 1-8.	1.7	0
5	Importance of personality and coping expectancy on patient-reported hearing disability, quality of life and distress level: a study of patients referred to an audiology service. Health and Quality of Life Outcomes, 2021, 19, 168.	2.4	6
6	Abstract 2520: Detection of transcriptional active HPV 16/18 in patients with oropharyngeal squamous cell carcinoma by dual immunohistochemistry p16INK4and in situ hybridization E6/E7 mRNA in archival material older than 25 years. , 2021, , .		0
7	Acoustic Voice Analysis and Maximum Phonation Time in Relation to Voice Handicap Index Score and Larynx Disease. Journal of Voice, 2020, 34, 161.e27-161.e35.	1.5	25
8	The Acute Phase Reaction and Its Prognostic Impact in Patients with Head and Neck Squamous Cell Carcinoma: Single Biomarkers Including C-Reactive Protein Versus Biomarker Profiles. Biomedicines, 2020, 8, 418.	3.2	9
9	Evaluation of Self-reported Symptoms in 1,457 Dizzy Patients and Associations With Caloric Testing and Posturography. Otology and Neurotology, 2020, 41, 956-963.	1.3	4
10	Reduced grey- and white matter volumes due to unilateral hearing loss following treatment for vestibular schwannoma. Heliyon, 2020, 6, e05658.	3.2	5
11	An fMRI-study on single-sided deafness: Spectral-temporal properties and side of stimulation modulates hemispheric dominance. NeuroImage: Clinical, 2019, 24, 101969.	2.7	8
12	EORTC Quality of Life Questionnaire Head and Neck (H&N)-35 scores from H&N squamous cell carcinoma patients obtained at diagnosis and at 6, 9 and 12Amonths following diagnosis predict 10-year overall survival. European Archives of Oto-Rhino-Laryngology, 2019, 276, 3495-3505.	1.6	8
13	Quality of life in persons with hearing loss: a study of patients referred to an audiological service. International Journal of Audiology, 2019, 58, 696-703.	1.7	4
14	Hearing as an Independent Predictor of Postural Balance in 1075ÂPatients Evaluated for Dizziness. Otolaryngology - Head and Neck Surgery, 2019, 161, 478-484.	1.9	27
15	Periodontal status at diagnosis predicts non-disease-specific survival in a geographically defined cohort of patients with oropharynx squamous cell carcinoma. Acta Oto-Laryngologica, 2019, 139, 309-315.	0.9	4
16	General health-related quality of life scores from head and neck squamous cell carcinoma patients obtained throughout the first year following diagnosis predicted up to 10-year overall survival. European Archives of Oto-Rhino-Laryngology, 2018, 275, 207-217.	1.6	12
17	Tumor stromal desmoplasia and inflammatory response uniquely predict survival with and without stratification for HPV tumor infection in OPSCC patients. Acta Oto-Laryngologica, 2018, 138, 1035-1042.	0.9	11
18	Generic quality of life in persons with hearing loss: a systematic literature review. BMC Ear, Nose and Throat Disorders, 2018, 18, 1.	2.6	92

HANS JÃ, RGEN AARSTAD

#	Article	IF	CITATIONS
19	Clinical Application and Psychometric Properties of a Norwegian Questionnaire for the Self-Assessment of Communication in Quiet and Adverse Conditions Using Two Revised APHAB Subscales. Journal of the American Academy of Audiology, 2018, 29, 025-034.	0.7	8
20	<i>In vitro</i> Monocyte <scp>IL</scp> â€6 Secretion Levels Following Stimulation with Autologous Spheroids Derived from Tumour or Benign Mucosa Predict Longâ€ŧerm Survival in Head and Neck Squamous Cell Carcinoma Patients. Scandinavian Journal of Immunology, 2017, 85, 211-219.	2.7	5
21	Health-related Quality of Life as Studied by EORTC QLQ and Voice Handicap Index Among Various Patients With Laryngeal Disease. Journal of Voice, 2017, 31, 251.e17-251.e26.	1.5	17
22	Primary surgery results in no survival benefit compared to primary radiation for oropharyngeal cancer patients stratified by high-risk human papilloma virus status. European Archives of Oto-Rhino-Laryngology, 2017, 274, 477-487.	1.6	17
23	Congenital laryngomalacia is related to exercise-induced laryngeal obstruction in adolescence. Archives of Disease in Childhood, 2016, 101, 443-448.	1.9	41
24	Functional-structural reorganisation of the neuronal network for auditory perception in subjects with unilateral hearing loss: Review of neuroimaging studies. Hearing Research, 2016, 332, 73-79.	2.0	19
25	The Norwegian Voice Handicap Index (VHI-N) patient scores are dependent on voice-related disease group. European Archives of Oto-Rhino-Laryngology, 2015, 272, 2897-2905.	1.6	9
26	Peripheral blood monocyte and T″ymphocyte activation levels at diagnosis predict longâ€ŧerm survival in head and neck squamous cell carcinoma patients. Apmis, 2015, 123, 305-314.	2.0	13
27	The impact of HPV infection, smoking history, age and operability of the patient on disease-specific survival in a geographically defined cohort of patients with oropharyngeal squamous cell carcinoma. Acta Oto-Laryngologica, 2014, 134, 964-973.	0.9	18
28	Perioperative Cytokine Response to Infection Associated With Elective Arterial Surgery. Vascular and Endovascular Surgery, 2014, 48, 116-122.	0.7	3
29	Level of distress predicts subsequent survival in successfully treated head and neck cancer patients: a prospective cohort study. Acta Oto-Laryngologica, 2014, 134, 211-219.	0.9	14
30	The impact of HPV infection on survival in a geographically defined cohort of oropharynx squamous cell carcinoma (OPSCC) patients in whom surgical treatment has been one main treatment. Acta Oto-Laryngologica, 2014, 134, 636-645.	0.9	29
31	Prediction of 5 year survival from level of perceived distress in newly diagnosed head and neck squamous cell carcinoma patients. Oral Oncology, 2013, 49, 964-969.	1.5	12
32	Cross-Cultural Adaption and Translation of the Voice Handicap Index into Norwegian. Folia Phoniatrica Et Logopaedica, 2012, 64, 234-240.	1.1	30
33	Exercise-induced laryngeal obstruction: natural history and effect of surgical treatment. European Archives of Oto-Rhino-Laryngology, 2011, 268, 1485-1492.	1.6	83
34	Mononuclear phagocytes in head and neck squamous cell carcinoma. European Archives of Oto-Rhino-Laryngology, 2010, 267, 335-344.	1.6	13
35	In vitro cholesteatoma growth and secretion of cytokines. Acta Oto-Laryngologica, 2010, 130, 815-819.	0.9	11
36	Presence of activated T lymphocytes in peripheral blood of head and neck squamous cell carcinoma patients predicts impaired prognosis. Acta Oto-Laryngologica, 2006, 126, 1326-1333.	0.9	19

#	Article	IF	CITATIONS
37	Mood, anxiety and sense of humor in head and neck cancer patients in relation to disease stage, prognosis and quality of life. Acta Oto-Laryngologica, 2005, 125, 557-565.	0.9	48
38	Head and neck squamous cell carcinoma spheroid- and monocyte spheroid-stimulated IL-6 and monocyte chemotactic protein-1 secretion are related to TNM stage, inflammatory state and tumor macrophage density. Acta Oto-Laryngologica, 2005, 125, 1097-1104.	0.9	15
39	Personality Traits in Head and Neck Squamous Cell Carcinoma Patients in Relation to the Disease State, Disease Extent and Prognosis. Acta Oto-Laryngologica, 2002, 122, 892-899.	0.9	21