

Motohiro Tani

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9248358/publications.pdf>

Version: 2024-02-01

64
papers

2,267
citations

218381

26
h-index

223531

46
g-index

66
all docs

66
docs citations

66
times ranked

2029
citing authors

#	ARTICLE	IF	CITATIONS
1	Regulation of sphingolipid biosynthesis in the endoplasmic reticulum via signals from the plasma membrane in budding yeast. <i>FEBS Journal</i> , 2022, 289, 457-472.	2.2	8
2	Crosstalk between protein kinase A and the HOG pathway under impaired biosynthesis of complex sphingolipids in budding yeast. <i>FEBS Journal</i> , 2022, 289, 766-786.	2.2	5
3	Involvement of the mitochondrial retrograde pathway in dihydrosphingosine-induced cytotoxicity in budding yeast. <i>Biochemical and Biophysical Research Communications</i> , 2022, 605, 63-69.	1.0	0
4	Identification of <i>sur2</i> mutation affecting the lifespan of fission yeast. <i>FEMS Microbiology Letters</i> , 2021, 368, .	0.7	2
5	ROS-mediated synthetic growth defect caused by impaired metabolism of sphingolipids and phosphatidylserine in budding yeast. <i>Bioscience, Biotechnology and Biochemistry</i> , 2020, 84, 2529-2532.	0.6	2
6	Proper regulation of inositolphosphorylceramide levels is required for acquirement of low pH resistance in budding yeast. <i>Scientific Reports</i> , 2020, 10, 10792.	1.6	10
7	Mitochondrial reactive oxygen species-mediated cytotoxicity of intracellularly accumulated dihydrosphingosine in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2020, 287, 3427-3448.	2.2	9
8	Overexpression of PDR16 confers resistance to complex sphingolipid biosynthesis inhibitor aureobasidin A in yeast <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 2018, 365, .	0.7	13
9	Protection mechanisms against aberrant metabolism of sphingolipids in budding yeast. <i>Current Genetics</i> , 2018, 64, 1021-1028.	0.8	15
10	Protective role of the HOG pathway against the growth defect caused by impaired biosynthesis of complex sphingolipids in yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2018, 107, 363-386.	1.2	21
11	Mannosylinositol phosphorylceramides and ergosterol coordinately maintain cell wall integrity in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2018, 285, 2405-2427.	2.2	26
12	Neutral Ceramidase. , 2018, , 3450-3457.		0
13	Structure-Function Relationship of Complex Sphingolipids in Yeast. <i>Trends in Glycoscience and Glycotechnology</i> , 2016, 28, E109-E116.	0.0	14
14	Structure-Function Relationship of Complex Sphingolipids in Yeast. <i>Trends in Glycoscience and Glycotechnology</i> , 2016, 28, J107-J114.	0.0	0
15	Yeast lacking the amphiphysin family protein Rvs167 is sensitive to disruptions in sphingolipid levels. <i>FEBS Journal</i> , 2016, 283, 2911-2928.	2.2	11
16	Neutral Ceramidase. , 2016, , 1-8.		0
17	VID22 is required for transcriptional activation of the PSD2 gene in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 2015, 472, 319-328.	1.7	3
18	Phospholipid methylation controls Atg32-mediated mitophagy and Atg8 recycling. <i>EMBO Journal</i> , 2015, 34, 2703-2719.	3.5	39

#	ARTICLE	IF	CITATIONS
19	Ergosteryl- β -glucosidase (Egh1) involved in sterylglucoside catabolism and vacuole formation in <i>Saccharomyces cerevisiae</i> . <i>Glycobiology</i> , 2015, 25, 1079-1089.	1.3	20
20	COX assembly factor cdc56 regulates mitochondrial morphology by affecting mitochondrial recruitment of Drp1. <i>FEBS Letters</i> , 2015, 589, 3126-3132.	1.3	8
21	Synthesis of mannosylinositol phosphorylceramides is involved in maintenance of cell integrity of yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2015, 95, 706-722.	1.2	24
22	Alteration of complex sphingolipid composition and its physiological significance in yeast <i>Saccharomyces cerevisiae</i> lacking vacuolar ATPase. <i>Microbiology (United Kingdom)</i> , 2015, 161, 2369-2383.	0.7	19
23	Involvement of Sac1 phosphoinositide phosphatase in the metabolism of phosphatidylserine in the yeast <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2014, 31, 145-158.	0.8	23
24	Change in activity of serine palmitoyltransferase affects sensitivity to syringomycin E in yeast <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 2014, 358, 64-71.	0.7	15
25	Loss of hydroxyl groups from the ceramide moiety can modify the lateral diffusion of membrane proteins in <i>S. cerevisiae</i> . <i>Journal of Lipid Research</i> , 2014, 55, 1343-1356.	2.0	23
26	New insight into the structure, reaction mechanism, and biological functions of neutral ceramidase. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2014, 1841, 682-691.	1.2	30
27	Hydroxylation state of fatty acid and long-chain base moieties of sphingolipid determine the sensitivity to growth inhibition due to AUR1 repression in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2012, 417, 673-678.	1.0	19
28	Involvement of complex sphingolipids and phosphatidylserine in endosomal trafficking in yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2012, 86, 1262-1280.	1.2	20
29	Expression of budding yeast IPT1 produces mannosyl-diinositol phosphorylceramide in fission yeast and inhibits cell growth. <i>Microbiology (United Kingdom)</i> , 2012, 158, 1219-1228.	0.7	6
30	FMP30 is required for the maintenance of a normal cardiolipin level and mitochondrial morphology in the absence of mitochondrial phosphatidylethanolamine synthesis. <i>Molecular Microbiology</i> , 2011, 80, 248-265.	1.2	34
31	The Mucin Box and Signal/Anchor Sequence of Rat Neutral Ceramidase Recruit Bacterial Sphingomyelinase to the Plasma Membrane. <i>Bioscience, Biotechnology and Biochemistry</i> , 2011, 75, 987-990.	0.6	2
32	Requirement of a specific group of sphingolipid-metabolizing enzyme for growth of yeast <i>Saccharomyces cerevisiae</i> under impaired metabolism of glycerophospholipids. <i>Molecular Microbiology</i> , 2010, 78, 395-413.	1.2	24
33	Defect of synthesis of very long-chain fatty acids confers resistance to growth inhibition by inositol phosphorylceramide synthase repression in yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Biochemistry</i> , 2010, 148, 565-571.	0.9	27
34	Mannosylinositol phosphorylceramide is a major sphingolipid component and is required for proper localization of plasma-membrane proteins in <i>Schizosaccharomyces pombe</i> . <i>Journal of Cell Science</i> , 2010, 123, 1578-1587.	1.2	47
35	Mechanistic Insights into the Hydrolysis and Synthesis of Ceramide by Neutral Ceramidase. <i>Journal of Biological Chemistry</i> , 2009, 284, 9566-9577.	1.6	30
36	Sphingomyelin synthase 2 is palmitoylated at the COOH-terminal tail, which is involved in its localization in plasma membranes. <i>Biochemical and Biophysical Research Communications</i> , 2009, 381, 328-332.	1.0	28

#	ARTICLE	IF	CITATIONS
37	Klotho-related Protein Is a Novel Cytosolic Neutral β -Glycosylceramidase. <i>Journal of Biological Chemistry</i> , 2007, 282, 30889-30900.	1.6	84
38	Neutral Sphingomyelinase 2 Is Palmitoylated on Multiple Cysteine Residues. <i>Journal of Biological Chemistry</i> , 2007, 282, 10047-10056.	1.6	63
39	Large-scale purification and characterization of recombinant <i>Pseudomonas</i> ceramidase: regulation by calcium. <i>Journal of Lipid Research</i> , 2007, 48, 600-608.	2.0	18
40	Lack of sphingosine 1-phosphate-degrading enzymes in erythrocytes. <i>Biochemical and Biophysical Research Communications</i> , 2007, 357, 212-217.	1.0	166
41	Analysis of membrane topology of neutral sphingomyelinase 2. <i>FEBS Letters</i> , 2007, 581, 1323-1328.	1.3	57
42	Ceramide/sphingosine/sphingosine 1-phosphate metabolism on the cell surface and in the extracellular space. <i>Cellular Signalling</i> , 2007, 19, 229-237.	1.7	140
43	The Extended Family of Neutral Sphingomyelinases. <i>Biochemistry</i> , 2006, 45, 11247-11256.	1.2	156
44	Rescue of cell growth by sphingosine with disruption of lipid microdomain formation in <i>Saccharomyces cerevisiae</i> deficient in sphingolipid biosynthesis. <i>Biochemical Journal</i> , 2006, 394, 237-242.	1.7	27
45	Neutral Ceramidase as an Integral Modulator for the Generation of S1P and S1P-Mediated Signaling. , 2006, , 183-196.		0
46	C18:3-GM1a induces apoptosis in Neuro2a cells: enzymatic remodeling of fatty acyl chains of glycosphingolipids. <i>Journal of Lipid Research</i> , 2005, 46, 1103-1112.	2.0	11
47	Mechanisms of sphingosine and sphingosine 1-phosphate generation in human platelets. <i>Journal of Lipid Research</i> , 2005, 46, 2458-2467.	2.0	114
48	Involvement of Neutral Ceramidase in Ceramide Metabolism at the Plasma Membrane and in Extracellular Milieu. <i>Journal of Biological Chemistry</i> , 2005, 280, 36592-36600.	1.6	46
49	Subcellular localization of human neutral ceramidase expressed in HEK293 cells. <i>Biochemical and Biophysical Research Communications</i> , 2005, 331, 37-42.	1.0	58
50	Conserved Amino Acid Residues in the COOH-terminal Tail Are Indispensable for the Correct Folding and Localization and Enzyme Activity of Neutral Ceramidase. <i>Journal of Biological Chemistry</i> , 2004, 279, 29351-29358.	1.6	8
51	Molecular Cloning and Functional Analysis of Zebrafish Neutral Ceramidase. <i>Journal of Biological Chemistry</i> , 2004, 279, 44012-44022.	1.6	46
52	O-Glycosylation of Mucin-like Domain Retains the Neutral Ceramidase on the Plasma Membranes as a Type II Integral Membrane Protein. <i>Journal of Biological Chemistry</i> , 2003, 278, 10523-10530.	1.6	56
53	A neutral ceramidase homologue from <i>Dictyostelium discoideum</i> exhibits an acidic pH optimum. <i>Biochemical Journal</i> , 2003, 376, 473-479.	1.7	22
54	Molecular Cloning and Characterization of a Secretory Neutral Ceramidase of <i>Drosophila melanogaster</i> . <i>Journal of Biochemistry</i> , 2002, 132, 229-236.	0.9	49

#	ARTICLE	IF	CITATIONS
55	Molecular Evolution of Neutral Ceramidase: From Bacteria to Mammals. <i>Molecular Biology Intelligence Unit</i> , 2002, , 41-48.	0.2	6
56	Purification, Characterization, Molecular Cloning, and Subcellular Distribution of Neutral Ceramidase of Rat Kidney. <i>Journal of Biological Chemistry</i> , 2001, 276, 26249-26259.	1.6	107
57	[¹⁴ C] Enzymatic synthesis of [¹⁴ C]ceramide, [¹⁴ C]glycosphingolipids, and [¹⁴ C]-aminoceramide. <i>Methods in Enzymology</i> , 2000, 311, 682-689.	0.4	12
58	Purification and Characterization of a Neutral Ceramidase from Mouse Liver. <i>Journal of Biological Chemistry</i> , 2000, 275, 3462-3468.	1.6	92
59	Neutral/Alkaline and Acid Ceramidase Activities Are Actively Released by Murine Endothelial Cells. <i>Biochemical and Biophysical Research Communications</i> , 2000, 275, 746-751.	1.0	50
60	Molecular Cloning of the Full-length cDNA Encoding Mouse Neutral Ceramidase. <i>Journal of Biological Chemistry</i> , 2000, 275, 11229-11234.	1.6	102
61	Specific and Sensitive Assay for Alkaline and Neutral Ceramidases Involving C12-NBD-Ceramide. <i>Journal of Biochemistry</i> , 1999, 125, 746-749.	0.9	41
62	Preparation of Fluorescence-Labeled GM1 and Sphingomyelin by the Reverse Hydrolysis Reaction of Sphingolipid Ceramide N-Deacylase as Substrates for Assay of Sphingolipid-Degrading Enzymes and for Detection of Sphingolipid-Binding Proteins. <i>Journal of Biochemistry</i> , 1999, 126, 604-611.	0.9	17
63	Enzymatic Synthesis of [¹⁴ C]-Amino-Ceramide: Preparation of a Sensitive Fluorescent Substrate for Ceramidase. <i>Analytical Biochemistry</i> , 1998, 263, 183-188.	1.1	40
64	Purification and Characterization of a Novel Ceramidase from <i>Pseudomonas aeruginosa</i> . <i>Journal of Biological Chemistry</i> , 1998, 273, 14368-14373.	1.6	101