

Karien Bloem

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9252656/publications.pdf>

Version: 2024-02-01

13
papers

403
citations

933264

10
h-index

1125617

13
g-index

13
all docs

13
docs citations

13
times ranked

565
citing authors

#	ARTICLE	IF	CITATIONS
1	Comprehensive evaluation of microneedle-based intradermal adalimumab delivery vs. subcutaneous administration: results of a randomized controlled clinical trial. <i>British Journal of Clinical Pharmacology</i> , 2021, 87, 3162-3176.	1.1	11
2	Infliximab Level Between Venous and Capillary Blood Using Novel Device Strongly Correlate in Paediatric Inflammatory Bowel Disease Patients. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2021, 72, 56-60.	0.9	2
3	Monitoring of Adalimumab Concentrations at Home in Patients with Inflammatory Bowel Disease Using Dried Blood Samples. <i>Therapeutic Drug Monitoring</i> , 2020, 42, 289-294.	1.0	13
4	Clinical Impact of Antibodies against Ustekinumab in Psoriasis: An Observational, Cross-Sectional, Multicenter Study. <i>Journal of Investigative Dermatology</i> , 2020, 140, 2129-2137.	0.3	6
5	Association of Serum Ustekinumab Levels With Clinical Response in Psoriasis. <i>JAMA Dermatology</i> , 2019, 155, 1235.	2.0	30
6	Dynamics of circulating TNF during adalimumab treatment using a drug-tolerant TNF assay. <i>Science Translational Medicine</i> , 2019, 11, .	5.8	41
7	Dried blood samples can support monitoring of infliximab concentrations in patients with inflammatory bowel disease: A clinical validation. <i>British Journal of Clinical Pharmacology</i> , 2019, 85, 1544-1551.	1.1	26
8	Defining the Therapeutic Range for Adalimumab and Predicting Response in Psoriasis: A Multicenter Prospective Observational Cohort Study. <i>Journal of Investigative Dermatology</i> , 2019, 139, 115-123.	0.3	60
9	Capillary blood microsampling to determine serum biopharmaceutical concentration: Mitra [®] microampler vs dried blood spot. <i>Bioanalysis</i> , 2018, 10, 815-823.	0.6	41
10	Immunogenicity of Therapeutic Antibodies: Monitoring Antidrug Antibodies in a Clinical Context. <i>Therapeutic Drug Monitoring</i> , 2017, 39, 327-332.	1.0	56
11	Dried blood spots from finger prick facilitate therapeutic drug monitoring of adalimumab and anti-adalimumab in patients with inflammatory diseases. <i>British Journal of Clinical Pharmacology</i> , 2017, 83, 2474-2484.	1.1	31
12	Predictive Value of Serum Infliximab Levels at Induction Phase in Rheumatoid Arthritis Patients. <i>Open Rheumatology Journal</i> , 2017, 11, 75-87.	0.1	16
13	Systematic comparison of drug-tolerant assays for anti-drug antibodies in a cohort of adalimumab-treated rheumatoid arthritis patients. <i>Journal of Immunological Methods</i> , 2015, 418, 29-38.	0.6	70