
## Charlotte Hellgren

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9254500/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Biological Aspects of Postpartum Depression. Women's Health, 2012, 8, 659-672.	0.7	112
2	Low Serum Allopregnanolone Is Associated with Symptoms of Depression in Late Pregnancy. Neuropsychobiology, 2014, 69, 147-153.	0.9	87
3	Prenatal and Postpartum Evening Salivary Cortisol Levels in Association with Peripartum Depressive Symptoms. PLoS ONE, 2015, 10, e0135471.	1.1	69
4	Cortisol awakening response in late pregnancy in women with previous or ongoing depression. Psychoneuroendocrinology, 2013, 38, 3150-3154.	1.3	51
5	Inflammatory markers in late pregnancy in association with postpartum depression—A nested case-control study. Psychoneuroendocrinology, 2017, 79, 146-159.	1.3	51
6	Lower inflammatory markers in women with antenatal depression brings the M1/M2 balance into focus from a new direction. Psychoneuroendocrinology, 2017, 80, 15-25.	1.3	48
7	Allopregnanolone levels and depressive symptoms during pregnancy in relation to single nucleotide polymorphisms in the allopregnanolone synthesis pathway. Hormones and Behavior, 2017, 94, 106-113.	1.0	36
8	Sleep duration, depression, and oxytocinergic genotype influence prepulse inhibition of the startle reflex in postpartum women. European Neuropsychopharmacology, 2016, 26, 767-776.	0.3	30
9	Cohort profile: the Biology, Affect, Stress, Imaging and Cognition (BASIC) study on perinatal depression in a population-based Swedish cohort. BMJ Open, 2019, 9, e031514.	0.8	30
10	MID-PREGNANCY CORTICOTROPIN-RELEASING HORMONE LEVELS IN ASSOCIATION WITH POSTPARTUM DEPRESSIVE SYMPTOMS. Depression and Anxiety, 2016, 33, 1023-1030.	2.0	23
11	Tandem mass spectrometry determined maternal cortisone to cortisol ratio and psychiatric morbidity during pregnancyâ^'interaction with birth weight. Psychoneuroendocrinology, 2016, 69, 142-149.	1.3	23
12	Stress-related genetic polymorphisms in association with peripartum depression symptoms and stress hormones: A longitudinal population-based study. Psychoneuroendocrinology, 2019, 103, 296-305.	1.3	18
13	Supraphysiological hormonal status, anxiety disorders, and COMT Val/Val genotype are associated with reduced sensorimotor gating in women. Psychoneuroendocrinology, 2015, 60, 217-223.	1.3	17
14	Decreased startle modulation during anticipation in the postpartum period in comparison to late pregnancy. Archives of Women's Mental Health, 2012, 15, 87-94.	1.2	15
15	The effect of antenatal depression and antidepressant treatment on placental tissue: a protein-validated gene expression study. BMC Pregnancy and Childbirth, 2019, 19, 479.	0.9	14
16	Different patterns of attentional bias in antenatal and postpartum depression. Brain and Behavior, 2017, 7, e00844.	1.0	13
17	Sympathetic reactivity in late pregnancy is related to labour onset in women. Stress, 2011, 14, 627-633.	0.8	12
18	Influence of catechol-O-methyltransferase Val158Met polymorphism on startle response in the presence of high estradiol levels. European Neuropsychopharmacology, 2013, 23, 629-635.	0.3	11

#	Article	lF	CITATIONS
19	Hypothalamic-pituitary-adrenal axis responsiveness, startle response, and sensorimotor gating in late pregnancy. Psychoneuroendocrinology, 2019, 106, 1-8.	1.3	7
20	Neuroticism is associated with higher antenatal care utilization in obstetric lowâ€risk women. Acta Obstetricia Et Gynecologica Scandinavica, 2019, 98, 470-478.	1.3	2
21	Skin conductance activity in post-term pregnancies. Journal of Maternal-Fetal and Neonatal Medicine, 2015, 28, 1912-1916.	0.7	Ο