

# Charlotte Hellgren

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9254500/publications.pdf>

Version: 2024-02-01

21  
papers

678  
citations

623188

14  
h-index

713013

21  
g-index

22  
all docs

22  
docs citations

22  
times ranked

870  
citing authors

#	ARTICLE	IF	CITATIONS
1	Neuroticism is associated with higher antenatal care utilization in obstetric low-risk women. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2019, 98, 470-478.	1.3	2
2	Hypothalamic-pituitary-adrenal axis responsiveness, startle response, and sensorimotor gating in late pregnancy. <i>Psychoneuroendocrinology</i> , 2019, 106, 1-8.	1.3	7
3	Stress-related genetic polymorphisms in association with peripartum depression symptoms and stress hormones: A longitudinal population-based study. <i>Psychoneuroendocrinology</i> , 2019, 103, 296-305.	1.3	18
4	Cohort profile: the Biology, Affect, Stress, Imaging and Cognition (BASIC) study on perinatal depression in a population-based Swedish cohort. <i>BMJ Open</i> , 2019, 9, e031514.	0.8	30
5	The effect of antenatal depression and antidepressant treatment on placental tissue: a protein-validated gene expression study. <i>BMC Pregnancy and Childbirth</i> , 2019, 19, 479.	0.9	14
6	Lower inflammatory markers in women with antenatal depression brings the M1/M2 balance into focus from a new direction. <i>Psychoneuroendocrinology</i> , 2017, 80, 15-25.	1.3	48
7	Inflammatory markers in late pregnancy in association with postpartum depression – A nested case-control study. <i>Psychoneuroendocrinology</i> , 2017, 79, 146-159.	1.3	51
8	Allopregnanolone levels and depressive symptoms during pregnancy in relation to single nucleotide polymorphisms in the allopregnanolone synthesis pathway. <i>Hormones and Behavior</i> , 2017, 94, 106-113.	1.0	36
9	Different patterns of attentional bias in antenatal and postpartum depression. <i>Brain and Behavior</i> , 2017, 7, e00844.	1.0	13
10	MID-PREGNANCY CORTICOTROPIN-RELEASING HORMONE LEVELS IN ASSOCIATION WITH POSTPARTUM DEPRESSIVE SYMPTOMS. <i>Depression and Anxiety</i> , 2016, 33, 1023-1030.	2.0	23
11	Tandem mass spectrometry determined maternal cortisone to cortisol ratio and psychiatric morbidity during pregnancy – interaction with birth weight. <i>Psychoneuroendocrinology</i> , 2016, 69, 142-149.	1.3	23
12	Sleep duration, depression, and oxytocinergic genotype influence prepulse inhibition of the startle reflex in postpartum women. <i>European Neuropsychopharmacology</i> , 2016, 26, 767-776.	0.3	30
13	Supraphysiological hormonal status, anxiety disorders, and COMT Val/Val genotype are associated with reduced sensorimotor gating in women. <i>Psychoneuroendocrinology</i> , 2015, 60, 217-223.	1.3	17
14	Skin conductance activity in post-term pregnancies. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2015, 28, 1912-1916.	0.7	0
15	Prenatal and Postpartum Evening Salivary Cortisol Levels in Association with Peripartum Depressive Symptoms. <i>PLoS ONE</i> , 2015, 10, e0135471.	1.1	69
16	Low Serum Allopregnanolone Is Associated with Symptoms of Depression in Late Pregnancy. <i>Neuropsychobiology</i> , 2014, 69, 147-153.	0.9	87
17	Cortisol awakening response in late pregnancy in women with previous or ongoing depression. <i>Psychoneuroendocrinology</i> , 2013, 38, 3150-3154.	1.3	51
18	Influence of catechol-O-methyltransferase Val158Met polymorphism on startle response in the presence of high estradiol levels. <i>European Neuropsychopharmacology</i> , 2013, 23, 629-635.	0.3	11

#	ARTICLE	IF	CITATIONS
19	Biological Aspects of Postpartum Depression. <i>Women's Health</i> , 2012, 8, 659-672.	0.7	112
20	Decreased startle modulation during anticipation in the postpartum period in comparison to late pregnancy. <i>Archives of Women's Mental Health</i> , 2012, 15, 87-94.	1.2	15
21	Sympathetic reactivity in late pregnancy is related to labour onset in women. <i>Stress</i> , 2011, 14, 627-633.	0.8	12