

Daniel E Francs

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9254537/daniel-e-frances-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

33
papers

688
citations

16
h-index

25
g-index

34
ext. papers

788
ext. citations

4.6
avg, IF

3.44
L-index

#	Paper	IF	Citations
33	Diethylnitrosamine enhances hepatic tumorigenic pathways in mice fed with high fat diet (Hfd). <i>Chemico-Biological Interactions</i> , 2019 , 303, 70-78	5	6
32	Protective Role of Hepatocyte Cyclooxygenase-2 Expression Against Liver Ischemia-Reperfusion Injury in Mice. <i>Hepatology</i> , 2019 , 70, 650-665	11.2	34
31	Heme oxygenase-1 induction by hemin prevents oxidative stress-induced acute cholestasis in the rat. <i>Clinical Science</i> , 2019 , 133, 117-134	6.5	10
30	Liver carbohydrates metabolism: A new islet-neogenesis associated protein peptide (INGAP-PP) target. <i>Peptides</i> , 2018 , 101, 44-50	3.8	
29	Disruption of tumor necrosis factor alpha receptor 1 signaling accelerates NAFLD progression in mice upon a high-fat diet. <i>Journal of Nutritional Biochemistry</i> , 2018 , 58, 17-27	6.3	16
28	PGE induces apoptosis of hepatic stellate cells and attenuates liver fibrosis in mice by downregulating miR-23a-5p and miR-28a-5p. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018 , 1864, 325-337	6.9	27
27	Diethylnitrosamine Increases Proliferation in Early Stages of Hepatic Carcinogenesis in Insulin-Treated Type 1 Diabetic Mice. <i>BioMed Research International</i> , 2018 , 2018, 9472939	3	13
26	Benznidazole, the trypanocidal drug used for Chagas disease, induces hepatic NRF2 activation and attenuates the inflammatory response in a murine model of sepsis. <i>Toxicology and Applied Pharmacology</i> , 2017 , 315, 12-22	4.6	12
25	The Enriched Proanthocyanidin Extract of <i>Ligaria cuneifolia</i> Shows a Marked Hypocholesterolemic Effect in Rats Fed with Cholesterol-Enriched Diet. <i>Recent Patents on Endocrine, Metabolic & Immune Drug Discovery</i> , 2017 , 11, 47-53		3
24	Cyclooxygenase-2 expression in hepatocytes attenuates non-alcoholic steatohepatitis and liver fibrosis in mice. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2016 , 1862, 1710-23	6.9	27
23	The trypanocidal benznidazole promotes adaptive response to oxidative injury: Involvement of the nuclear factor-erythroid 2-related factor-2 (Nrf2) and multidrug resistance associated protein 2 (MRP2). <i>Toxicology and Applied Pharmacology</i> , 2016 , 304, 90-8	4.6	16
22	Regulation of MicroRNA 183 by Cyclooxygenase 2 in Liver Is DEAD-Box Helicase p68 (DDX5) Dependent: Role in Insulin Signaling. <i>Molecular and Cellular Biology</i> , 2015 , 35, 2554-67	4.8	30
21	Hepatic cyclooxygenase-2 expression protects against diet-induced steatosis, obesity, and insulin resistance. <i>Diabetes</i> , 2015 , 64, 1522-31	0.9	30
20	Quercetin prevents liver carcinogenesis by inducing cell cycle arrest, decreasing cell proliferation and enhancing apoptosis. <i>Molecular Nutrition and Food Research</i> , 2014 , 58, 289-300	5.9	42
19	FoxO3a nuclear localization and its association with E-catenin and Smads in IFN- β -treated hepatocellular carcinoma cell lines. <i>Journal of Interferon and Cytokine Research</i> , 2014 , 34, 858-69	3.5	6
18	NOD1 receptor is up-regulated in diabetic human and murine myocardium. <i>Clinical Science</i> , 2014 , 127, 665-77	6.5	16
17	FoxO3a modulation and promotion of apoptosis by interferon- α 2b in rat preneoplastic liver. <i>Liver International</i> , 2014 , 34, 1566-77	7.9	8

16	Evidence for necrosis, but not apoptosis, in human hepatoma cells with knockdown of mitochondrial aquaporin-8. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2014 , 19, 851-94	5.4	13
15	Cyclooxygenase-2 over-expression inhibits liver apoptosis induced by hyperglycemia. <i>Journal of Cellular Biochemistry</i> , 2013 , 114, 669-80	4.7	16
14	Diabetes, an inflammatory process: Oxidative Stress and TNF-alpha involved in hepatic complication. <i>Journal of Biomedical Science and Engineering</i> , 2013 , 06, 645-653	0.7	36
13	Mitochondrial aquaporin-8 knockdown in human hepatoma HepG2 cells causes ROS-induced mitochondrial depolarization and loss of viability. <i>Toxicology and Applied Pharmacology</i> , 2012 , 264, 246-54	4.6	82
12	Benznidazole treatment attenuates liver NF- κ B activity and MAPK in a cecal ligation and puncture model of sepsis. <i>Molecular Immunology</i> , 2011 , 48, 867-73	4.3	22
11	Tumor necrosis factor alpha pathways develops liver apoptosis in type 1 diabetes mellitus. <i>Molecular Immunology</i> , 2011 , 48, 1397-407	4.3	43
10	ROS formation and antioxidant status in brain areas of rats exposed to sodium metavanadate. <i>Neurotoxicology and Teratology</i> , 2011 , 33, 297-302	3.9	26
9	Role of reactive oxygen species in the early stages of liver regeneration in streptozotocin-induced diabetic rats. <i>Free Radical Research</i> , 2011 , 45, 1143-53	4	9
8	Attenuation of the Wnt/beta-catenin/TCF pathway by in vivo interferon-alpha2b (IFN-alpha2b) treatment in preneoplastic rat livers. <i>Growth Factors</i> , 2010 , 28, 166-77	1.6	7
7	Hyperglycemia induces apoptosis in rat liver through the increase of hydroxyl radical: new insights into the insulin effect. <i>Journal of Endocrinology</i> , 2010 , 205, 187-200	4.7	55
6	Tumor necrosis factor alpha induced by Trypanosoma cruzi infection mediates inflammation and cell death in the liver of infected mice. <i>Cytokine</i> , 2010 , 49, 64-72	4	16
5	Cross-talk between IFN-alpha and TGF-beta1 signaling pathways in preneoplastic rat liver. <i>Growth Factors</i> , 2009 , 27, 1-11	1.6	13
4	Involvement of reactive oxygen species on the apoptotic mechanism induced by IFN-alpha2b in rat preneoplastic liver. <i>Biochemical Pharmacology</i> , 2007 , 73, 1776-85	6	18
3	Hepatocytes isolated from preneoplastic rat livers are resistant to ethacrynic acid cytotoxicity. <i>Archives of Toxicology</i> , 2007 , 81, 565-73	5.8	6
2	Vascular endothelial growth factor and nitric oxide in rat liver regeneration. <i>Life Sciences</i> , 2007 , 81, 750-5.8	5.8	15
1	Oxidative stress in primary culture hepatocytes isolated from partially hepatectomized rats. <i>Canadian Journal of Physiology and Pharmacology</i> , 2007 , 85, 1047-51	2.4	15