

Dai-Hui Peng

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9258983/publications.pdf>

Version: 2024-02-01

74
papers

2,162
citations

257450

24
h-index

265206

42
g-index

74
all docs

74
docs citations

74
times ranked

3185
citing authors

#	ARTICLE	IF	CITATIONS
1	Difference in the regulation of biological rhythm symptoms of Major depressive disorder between escitalopram and mirtazapine. <i>Journal of Affective Disorders</i> , 2022, 296, 258-264.	4.1	3
2	Impaired robust interhemispheric function integration of depressive brain from REST-meta-MDD database in China. <i>Bipolar Disorders</i> , 2022, 24, 400-411.	1.9	8
3	Evaluating the efficacy and moderators of algorithm-guided antidepressant treatments of major depressive disorder. <i>Journal of Affective Disorders</i> , 2022, 297, 68-75.	4.1	1
4	A Preliminary Study of Different Treatment Strategies for Anxious Depression. <i>Neuropsychiatric Disease and Treatment</i> , 2022, Volume 18, 11-18.	2.2	0
5	Bilateral Habenula deep brain stimulation for treatment-resistant depression: clinical findings and electrophysiological features. <i>Translational Psychiatry</i> , 2022, 12, 52.	4.8	21
6	Probing the clinical and brain structural boundaries of bipolar and major depressive disorder. <i>Translational Psychiatry</i> , 2021, 11, 48.	4.8	9
7	Disturbances of affective cognition in mood disorders. <i>Science China Life Sciences</i> , 2021, 64, 938-941.	4.9	3
8	Disrupted hemispheric connectivity specialization in patients with major depressive disorder: Evidence from the REST-meta-MDD Project. <i>Journal of Affective Disorders</i> , 2021, 284, 217-228.	4.1	23
9	Gastrointestinal Symptoms During Depressive Episodes in 3256 Patients with Major Depressive Disorders: Findings from the NSSD. <i>Journal of Affective Disorders</i> , 2021, 286, 27-32.	4.1	15
10	Clinical features of the patients with major depressive disorder co-occurring insomnia and hypersomnia symptoms: a report of NSSD study. <i>Sleep Medicine</i> , 2021, 81, 375-381.	1.6	13
11	Predictors and moderators of quality of life in patients with major depressive disorder: An AGTs-MDD study report. <i>Journal of Psychiatric Research</i> , 2021, 138, 96-102.	3.1	5
12	Reduced NLRP3 inflammasome expression in the brain is associated with stress resilience. <i>Psychoneuroendocrinology</i> , 2021, 128, 105211.	2.7	9
13	A Preliminary Randomized Controlled Trial of Different Treatment Regimens for Melancholic Depression. <i>Neuropsychiatric Disease and Treatment</i> , 2021, Volume 17, 2441-2449.	2.2	0
14	Disrupted intrinsic functional brain topology in patients with major depressive disorder. <i>Molecular Psychiatry</i> , 2021, 26, 7363-7371.	7.9	82
15	Reward mechanism of depressive episodes in bipolar disorder: Enhanced theta power in feedback-related negativity. <i>Journal of Affective Disorders</i> , 2021, 292, 217-222.	4.1	2
16	Neural biomarker of functional disability in major depressive disorder: A structural neuroimaging study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 111, 110337.	4.8	2
17	Brain structural alterations in MDD patients with gastrointestinal symptoms: Evidence from the REST-meta-MDD project. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 111, 110386.	4.8	18
18	Signatures of 4 autophagy-related genes as diagnostic markers of MDD and their correlation with immune infiltration. <i>Journal of Affective Disorders</i> , 2021, 295, 11-20.	4.1	20

#	ARTICLE	IF	CITATIONS
19	Symptomatology differences of major depression in psychiatric versus general hospitals: A machine learning approach. <i>Journal of Affective Disorders</i> , 2020, 260, 349-360.	4.1	7
20	Causes of drug discontinuation in patients with major depressive disorder in China. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2020, 96, 109755.	4.8	23
21	Altered resting-state dynamic functional brain networks in major depressive disorder: Findings from the REST-meta-MDD consortium. <i>NeuroImage: Clinical</i> , 2020, 26, 102163.	2.7	76
22	Biotypes of major depressive disorder: Neuroimaging evidence from resting-state default mode network patterns. <i>NeuroImage: Clinical</i> , 2020, 28, 102514.	2.7	51
23	Altered resting-state fMRI signals and network topological properties of bipolar depression patients with anxiety symptoms. <i>Journal of Affective Disorders</i> , 2020, 277, 358-367.	4.1	12
24	Challenges and opportunities in mental health services during the COVID-19 pandemic. <i>Annals of General Psychiatry</i> , 2020, 33, e100275.	3.1	19
25	<p>The Developmental and Translational Study on Biomarkers and Clinical Characteristics-based Diagnostic and Therapeutic Identification of Major Depressive Disorder: Study Protocol for a Multicenter Randomized Controlled Trial in China</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 2343-2351.	2.2	1
26	<p>Individual Perceived Stress Mediates Psychological Distress in Medical Workers During COVID-19 Epidemic Outbreak in Wuhan</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2020, Volume 16, 2529-2537.	2.2	18
27	The concurrent disturbance of dynamic functional and structural brain connectome in major depressive disorder: the prefronto-insular pathway. <i>Journal of Affective Disorders</i> , 2020, 274, 1084-1090.	4.1	10
28	The Insular Subregions Topological Characteristics of Patients With Bipolar Depressive Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 253.	2.6	7
29	Cortical thickness and subcortical volumes alterations in euthymic bipolar I patients treated with different mood stabilizers. <i>Brain Imaging and Behavior</i> , 2019, 13, 1255-1264.	2.1	8
30	Clinical characteristics associated with therapeutic nonadherence of the patients with major depressive disorder: A report on the National Survey on Symptomatology of Depression in China. <i>CNS Neuroscience and Therapeutics</i> , 2019, 25, 215-222.	3.9	8
31	Disagreement and factors between symptom on self-report and clinician rating of major depressive disorder: A report of a national survey in China. <i>Journal of Affective Disorders</i> , 2019, 253, 141-146.	4.1	10
32	Intrinsic gray-matter connectivity of the brain in major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 251, 78-85.	4.1	17
33	Reduced default mode network functional connectivity in patients with recurrent major depressive disorder. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 9078-9083.	7.1	441
34	Subtypes of treatment-resistant depression determined by a latent class analysis in a Chinese clinical population. <i>Journal of Affective Disorders</i> , 2019, 249, 82-89.	4.1	12
35	HMGB1 involved in stress-induced depression and its neuroinflammatory priming role: a systematic review. <i>Annals of General Psychiatry</i> , 2019, 32, e100084.	3.1	46
36	The association between somatic symptoms and suicidal ideation in Chinese first-episode major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 245, 17-21.	4.1	30

#	ARTICLE	IF	CITATIONS
37	Neuroimaging Advance in Depressive Disorder. <i>Advances in Experimental Medicine and Biology</i> , 2019, 1180, 59-83.	1.6	4
38	Neuroimaging genomic studies in major depressive disorder: A systematic review. <i>CNS Neuroscience and Therapeutics</i> , 2018, 24, 1020-1036.	3.9	13
39	Prevalence, risk factors and clinical characteristics of suicidal ideation in Chinese patients with depression. <i>Journal of Affective Disorders</i> , 2018, 235, 135-141.	4.1	40
40	The Metabolic Factor Kynurenic Acid of Kynurenine Pathway Predicts Major Depressive Disorder. <i>Frontiers in Psychiatry</i> , 2018, 9, 552.	2.6	62
41	Effects of tumor necrosis factor- $\hat{\pm}$ polymorphism on the brain structural changes of the patients with major depressive disorder. <i>Translational Psychiatry</i> , 2018, 8, 217.	4.8	25
42	Somatic symptoms vary in major depressive disorder in China. <i>Comprehensive Psychiatry</i> , 2018, 87, 32-37.	3.1	37
43	The clinical correlates of comorbid anxiety symptoms and syndromal anxiety in patients with major depressive disorder. <i>Psychiatry Research</i> , 2018, 269, 251-257.	3.3	18
44	Aberrant Neural Activity in Patients With Bipolar Depressive Disorder Distinguishing to the Unipolar Depressive Disorder: A Resting-State Functional Magnetic Resonance Imaging Study. <i>Frontiers in Psychiatry</i> , 2018, 9, 238.	2.6	28
45	Different levels of pro- and anti-inflammatory cytokines in patients with unipolar and bipolar depression. <i>Journal of Affective Disorders</i> , 2018, 237, 65-72.	4.1	47
46	Abnormal white matter integrity in Chinese young adults with first-episode medication-free anxious depression: a possible neurological biomarker of subtype major depressive disorder. <i>Neuropsychiatric Disease and Treatment</i> , 2018, Volume 14, 2017-2026.	2.2	13
47	Reduced ENA78 levels as novel biomarker for major depressive disorder and venlafaxine efficiency: Result from a prospective longitudinal study. <i>Psychoneuroendocrinology</i> , 2017, 81, 113-121.	2.7	21
48	Identification of plasma biomarkers for distinguishing bipolar depression from major depressive disorder by iTRAQ-coupled LC-MS/MS and bioinformatics analysis. <i>Psychoneuroendocrinology</i> , 2017, 86, 17-24.	2.7	51
49	Ratio of mBDNF to proBDNF for Differential Diagnosis of Major Depressive Disorder and Bipolar Depression. <i>Molecular Neurobiology</i> , 2017, 54, 5573-5582.	4.0	62
50	Complement factor H and susceptibility to major depressive disorder in Han Chinese. <i>British Journal of Psychiatry</i> , 2016, 208, 446-452.	2.8	21
51	Alterations of microRNA-124 expression in peripheral blood mononuclear cells in pre- and post-treatment patients with major depressive disorder. <i>Journal of Psychiatric Research</i> , 2016, 78, 65-71.	3.1	74
52	Evaluating the association between the SHANK3 gene and bipolar disorder. <i>Psychiatry Research</i> , 2016, 244, 284-288.	3.3	10
53	ZNF804A Genetic Variation Confers Risk to Bipolar Disorder. <i>Molecular Neurobiology</i> , 2016, 53, 2936-2943.	4.0	21
54	Atypical Features and Bipolar Disorder. <i>Shanghai Archives of Psychiatry</i> , 2016, 28, 166-168.	0.7	1

#	ARTICLE	IF	CITATIONS
55	Increased Cognition Connectivity Network in Major Depression Disorder: A fMRI Study. <i>Psychiatry Investigation</i> , 2015, 12, 227.	1.6	40
56	Atypical features and treatment choices in bipolar disorders: a result of the National Bipolar Mania Pathway Survey in China. <i>Neuroscience Bulletin</i> , 2015, 31, 22-30.	2.9	4
57	Down-regulation of PRKCB1 expression in Han Chinese patients with subsyndromal symptomatic depression. <i>Journal of Psychiatric Research</i> , 2015, 69, 1-6.	3.1	6
58	Guidelines concordance of maintenance treatment in euthymic patients with bipolar disorder: Data from the national bipolar mania pathway survey (BIPAS) in mainland China. <i>Journal of Affective Disorders</i> , 2015, 182, 101-105.	4.1	6
59	Dissociated large-scale functional connectivity networks of the precuneus in medication-naïve first-episode depression. <i>Psychiatry Research - Neuroimaging</i> , 2015, 232, 250-256.	1.8	65
60	Comorbid bipolar disorder and obsessive-compulsive disorder. <i>Shanghai Archives of Psychiatry</i> , 2015, 27, 246-8.	0.7	4
61	Surface Vulnerability of Cerebral Cortex to Major Depressive Disorder. <i>PLoS ONE</i> , 2015, 10, e0120704.	2.5	62
62	Nerve growth factor variations in patients with mood disorders: no changes in eight weeks of clinical treatment. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 835.	2.2	11
63	Alterations in effective connectivity anchored on the insula in major depressive disorder. <i>European Neuropsychopharmacology</i> , 2014, 24, 1784-1792.	0.7	58
64	Altered brain network modules induce helplessness in major depressive disorder. <i>Journal of Affective Disorders</i> , 2014, 168, 21-29.	4.1	57
65	Decreased serum fibroblast growth factor - 2 levels in pre- and post-treatment patients with major depressive disorder. <i>Neuroscience Letters</i> , 2014, 579, 168-172.	2.1	39
66	Influence of BCL2 gene in major depression susceptibility and antidepressant treatment outcome. <i>Journal of Affective Disorders</i> , 2014, 155, 288-294.	4.1	27
67	Elevated serum levels of FGF-2, NGF and IGF-1 in patients with manic episode of bipolar disorder. <i>Psychiatry Research</i> , 2014, 218, 54-60.	3.3	58
68	Guidelines Disconcordance in Acute Bipolar Depression: Data from the National Bipolar Mania Pathway Survey (BIPAS) in Mainland China. <i>PLoS ONE</i> , 2014, 9, e96096.	2.5	11
69	Evaluation of antidepressant polypharmacy and other interventions for treatment-resistant depression. <i>Shanghai Archives of Psychiatry</i> , 2014, 26, 365-7.	0.7	0
70	Difference in remission in a Chinese population with anxious versus nonanxious treatment-resistant depression: A report of OPERATION study. <i>Journal of Affective Disorders</i> , 2013, 150, 834-839.	4.1	58
71	The extracellular signal-regulated kinase pathway may play an important role in mediating antidepressant-stimulated hippocampus neurogenesis in depression. <i>Medical Hypotheses</i> , 2012, 79, 87-91.	1.5	15
72	Symptom severity is more closely associated with social functioning status in inpatients with schizophrenia than cognitive deficits. <i>Shanghai Archives of Psychiatry</i> , 2012, 24, 83-90.	0.7	0

#	ARTICLE	IF	CITATIONS
73	Abnormal functional connectivity with mood regulating circuit in unmedicated individual with major depression: a resting-state functional magnetic resonance study. Chinese Medical Journal, 2012, 125, 3701-6.	2.3	18
74	Decreased regional homogeneity in major depression as revealed by resting-state functional magnetic resonance imaging. Chinese Medical Journal, 2011, 124, 369-73.	2.3	45