

# MarÃ-a I Gil

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9259817/publications.pdf>

Version: 2024-02-01

224  
papers

16,587  
citations

14644

66  
h-index

17090

122  
g-index

231  
all docs

231  
docs citations

231  
times ranked

13309  
citing authors

#	ARTICLE	IF	CITATIONS
1	Strategies for mitigating chlorinated disinfection byproducts in wastewater treatment plants. <i>Chemosphere</i> , 2022, 288, 132583.	4.2	28
2	Bioactive compounds in lettuce: Highlighting the benefits to human health and impacts of preharvest and postharvest practices. <i>Comprehensive Reviews in Food Science and Food Safety</i> , 2022, 21, 4-45.	5.9	41
3	Frozen Vegetable Processing Plants Can Harbour Diverse <i>Listeria monocytogenes</i> Populations: Identification of Critical Operations by WGS. <i>Foods</i> , 2022, 11, 1546.	1.9	8
4	Practical applications of sensor-based methodologies for monitoring peracetic acid (PAA) as a disinfectant of fresh produce wash water. <i>Food Control</i> , 2021, 121, 107632.	2.8	5
5	Recent progress on the management of the industrial washing of fresh produce with a focus on microbiological risks. <i>Current Opinion in Food Science</i> , 2021, 38, 46-51.	4.1	24
6	Occurrence and Accumulation of Human Enteric Viruses and Phages in Process Water from the Fresh Produce Industry. <i>Foods</i> , 2021, 10, 1853.	1.9	5
7	UPLC-QTOF-MS metabolomics reveals biomarkers related to browning susceptibility of fresh-cut lettuce. <i>Acta Horticulturae</i> , 2021, , 43-46.	0.1	0
8	Monitoring of human enteric virus and coliphages throughout water reuse system of wastewater treatment plants to irrigation endpoint of leafy greens. <i>Science of the Total Environment</i> , 2021, 782, 146837.	3.9	21
9	Management of preharvest and postharvest factors related to quality and safety aspects of leafy vegetables. <i>Acta Horticulturae</i> , 2021, , 1-12.	0.1	1
10	Monitoring and control of wash water sanitation. <i>Acta Horticulturae</i> , 2021, , 75-80.	0.1	0
11	New standards at European Union level on water reuse for agricultural irrigation: Are the Spanish wastewater treatment plants ready to produce and distribute reclaimed water within the minimum quality requirements?. <i>International Journal of Food Microbiology</i> , 2021, 356, 109352.	2.1	18
12	Peroxyacetic acid and chlorine dioxide unlike chlorine induce viable but non-culturable (VBNC) stage of <i>Listeria monocytogenes</i> and <i>Escherichia coli</i> O157:H7 in wash water. <i>Food Microbiology</i> , 2021, 100, 103866.	2.1	14
13	Post-process treatments are effective strategies to reduce <i>Listeria monocytogenes</i> on the surface of leafy greens: A pilot study. <i>International Journal of Food Microbiology</i> , 2020, 313, 108390.	2.1	19
14	Critical points affecting the microbiological safety of bell peppers washed with peroxyacetic acid in a commercial packinghouse. <i>Food Microbiology</i> , 2020, 88, 103409.	2.1	14
15	La importancia del agua en la industria de alimentos vegetales. <i>Arbor</i> , 2020, 196, 547.	0.1	1
16	CA/MA on bioactive compounds. , 2020, , 131-146.		2
17	The impact of light on modified atmosphere storage and quality of fresh produce. , 2020, , 167-184.		1
18	Leafy vegetables: Fresh-cut lettuce. , 2020, , 545-550.		0

#	ARTICLE	IF	CITATIONS
19	Leafy vegetables: Fresh and fresh-cut mature spinach. , 2020, , 551-555.		3
20	Mushrooms. , 2020, , 577-581.		1
21	Leafy vegetables: Baby leaves. , 2020, , 527-536.		3
22	Use of Chlorine Dioxide to Treat Recirculated Process Water in a Commercial Tomato Packinghouse: Microbiological and Chemical Risks. <i>Frontiers in Sustainable Food Systems</i> , 2020, 4, .	1.8	2
23	Detection and Quantification Methods for Viable but Non-culturable (VBNC) Cells in Process Wash Water of Fresh-Cut Produce: Industrial Validation. <i>Frontiers in Microbiology</i> , 2020, 11, 673.	1.5	27
24	Chlorate accumulation in commercial lettuce cultivated in open field and irrigated with reclaimed water. <i>Food Control</i> , 2020, 114, 107283.	2.8	11
25	Chlorinated wash water and pH regulators affect chlorine gas emission and disinfection by-products. <i>Innovative Food Science and Emerging Technologies</i> , 2020, 66, 102533.	2.7	18
26	Chlorination management in commercial fresh produce processing lines. <i>Food Control</i> , 2019, 106, 106760.	2.8	28
27	Suitability of centrifuge water for detecting the presence of <i>Escherichia coli</i> versus finished fresh-cut lettuce testing. <i>Food Microbiology</i> , 2019, 84, 103271.	2.1	2
28	Chemical risks associated with ready-to-eat vegetables: quantitative analysis to estimate formation and/or accumulation of disinfection byproducts during washing. <i>EFSA Journal</i> , 2019, 17, e170913.	0.9	10
29	Operational limits of sodium hypochlorite for different fresh produce wash water based on microbial inactivation and disinfection by-products (DBPs). <i>Food Control</i> , 2019, 104, 300-307.	2.8	31
30	Targeted Metabolomics Analysis and Identification of Biomarkers for Predicting Browning of Fresh-Cut Lettuce. <i>Journal of Agricultural and Food Chemistry</i> , 2019, 67, 5908-5917.	2.4	24
31	Chlorate uptake during washing is influenced by product type and cut piece size, as well as washing time and wash water content. <i>Postharvest Biology and Technology</i> , 2019, 151, 45-52.	2.9	34
32	Postharvest research and industry implications. <i>Acta Horticulturae</i> , 2019, , 1-8.	0.1	1
33	Phyllosphere microbial communities of leafy vegetables affected by irrigation water sanitation. <i>Acta Horticulturae</i> , 2019, , 393-398.	0.1	1
34	Suitability of chlorine dioxide as a tertiary treatment for municipal wastewater and use of reclaimed water for overhead irrigation of baby lettuce. <i>Food Control</i> , 2019, 96, 186-193.	2.8	18
35	Disinfection by-products generated by sodium hypochlorite and electrochemical disinfection in different process wash water and fresh-cut products and their reduction by activated carbon. <i>Food Control</i> , 2019, 100, 46-52.	2.8	28
36	Impact of weather conditions, leaf age and irrigation water disinfection on the major epiphytic bacterial genera of baby spinach grown in an open field. <i>Food Microbiology</i> , 2019, 78, 46-52.	2.1	17

#	ARTICLE	IF	CITATIONS
37	Microbial and chemical characterization of commercial washing lines of fresh produce highlights the need for process water control. <i>Innovative Food Science and Emerging Technologies</i> , 2019, 51, 211-219.	2.7	46
38	Untargeted metabolomics to explain browning of fresh-cut lettuce. <i>Acta Horticulturae</i> , 2019, , 653-657.	0.1	0
39	Water and Wastewater Use in the Fresh Produce Industry: Food Safety and Environmental Implications. , 2018, , 59-76.		2
40	Disinfection byâ€products in baby lettuce irrigated with electrolysed water. <i>Journal of the Science of Food and Agriculture</i> , 2018, 98, 2981-2988.	1.7	20
41	Impact of relative humidity, inoculum carrier and size, and native microbiota on <i>Salmonella ser. Typhimurium</i> survival in baby lettuce. <i>Food Microbiology</i> , 2018, 70, 155-161.	2.1	29
42	Correlation between <i>E.Âcoli</i> levels and the presence of foodborne pathogens in surface irrigation water: Establishment of a sampling program. <i>Water Research</i> , 2018, 128, 226-233.	5.3	39
43	Demonstration tests of irrigation water disinfection with chlorine dioxide in open field cultivation of baby spinach. <i>Journal of the Science of Food and Agriculture</i> , 2018, 98, 2973-2980.	1.7	21
44	Effect of calcium and anti-browning agents on total phenols and antioxidant capability of â€Packhamâ€™s Triumphâ€™ pears packed in modified atmosphere. <i>Acta Horticulturae</i> , 2018, , 291-300.	0.1	0
45	Impact of climate change and global trends on the microbial quality of leafy greens. <i>Acta Horticulturae</i> , 2018, , 51-56.	0.1	0
46	Electrochemical disinfection of process wash water for the fresh-cut industry. <i>Acta Horticulturae</i> , 2018, , 371-378.	0.1	3
47	Impact of chlorine dioxide disinfection of irrigation water on the epiphytic bacterial community of baby spinach and underlying soil. <i>PLoS ONE</i> , 2018, 13, e0199291.	1.1	38
48	Irrigating Lettuce with Wastewater Effluent: Does Disinfection with Chlorine Dioxide Inactivate Viruses?. <i>Journal of Environmental Quality</i> , 2018, 47, 1139-1145.	1.0	23
49	LCâ€™MS untargeted metabolomics reveals early biomarkers to predict browning of fresh-cut lettuce. <i>Postharvest Biology and Technology</i> , 2018, 146, 9-17.	2.9	20
50	LC-MS Untargeted Metabolomics To Explain the Signal Metabolites Inducing Browning in Fresh-Cut Lettuce. <i>Journal of Agricultural and Food Chemistry</i> , 2017, 65, 4526-4535.	2.4	45
51	Ready-to-eat vegetables: Current problems and potential solutions to reduce microbial risk in the production chain. <i>LWT - Food Science and Technology</i> , 2017, 85, 284-292.	2.5	90
52	Off-odor compounds responsible for quality loss of minimally processed baby spinach stored under MA of low O2 and high CO2 using GCâ€™MS and olfactometry techniques. <i>Postharvest Biology and Technology</i> , 2017, 129, 129-135.	2.9	12
53	Quality and safety of fresh horticultural commodities: Recent advances and future perspectives. <i>Food Packaging and Shelf Life</i> , 2017, 14, 2-11.	3.3	51
54	Modelling of <i>E. coli</i> inactivation by chlorine dioxide in irrigation water. <i>Agricultural Water Management</i> , 2017, 192, 98-102.	2.4	20

#	ARTICLE	IF	CITATIONS
55	Impact of solar radiation exposure on phyllosphere bacterial community of red-pigmented baby leaf lettuce. <i>Food Microbiology</i> , 2017, 66, 77-85.	2.1	30
56	Quantitative contamination assessment of <i>Escherichia coli</i> in baby spinach primary production in Spain: Effects of weather conditions and agricultural practices. <i>International Journal of Food Microbiology</i> , 2017, 257, 238-246.	2.1	37
57	Influence of water stress and storage time on preservation of the fresh volatile profile of three basil genotypes. <i>Food Chemistry</i> , 2017, 221, 169-177.	4.2	14
58	Growing season climates affect quality of fresh-cut lettuce. <i>Postharvest Biology and Technology</i> , 2017, 123, 60-68.	2.9	32
59	A novel electrochemical device as a disinfection system to maintain water quality during washing of ready to eat fresh produce. <i>Food Control</i> , 2017, 71, 242-247.	2.8	27
60	Hyperspectral Imaging to Evaluate the Effect of Irrigation Water Salinity in Lettuce. <i>Applied Sciences (Switzerland)</i> , 2016, 6, 412.	1.3	17
61	Food safety management system (FSMS) adjusted to the characteristics of the leafy greens production chain context in Spain. <i>Acta Horticulturae</i> , 2016, , 219-224.	0.1	0
62	Preharvest factors and fresh-cut quality of leafy vegetables. <i>Acta Horticulturae</i> , 2016, , 57-64.	0.1	10
63	Suitability of different <i>Escherichia coli</i> enumeration techniques to assess the microbial quality of different irrigation water sources. <i>Food Microbiology</i> , 2016, 58, 29-35.	2.1	17
64	Untargeted metabolomics approach using UPLC-ESI-QTOF-MS to explore the metabolome of fresh-cut iceberg lettuce. <i>Metabolomics</i> , 2016, 12, 1.	1.4	66
65	Comprehensive evaluation of different storage conditions for the varietal screening of lettuce for fresh-cut performance. <i>Postharvest Biology and Technology</i> , 2016, 120, 36-44.	2.9	16
66	Occurrence of enteric viruses in reclaimed and surface irrigation water: relationship with microbiological and physicochemical indicators. <i>Journal of Applied Microbiology</i> , 2016, 121, 1180-1188.	1.4	37
67	Modified atmosphere (MA) prevents browning of fresh-cut romaine lettuce through multi-target effects related to phenolic metabolism. <i>Postharvest Biology and Technology</i> , 2016, 119, 84-93.	2.9	40
68	Should chlorate residues be of concern in fresh-cut salads?. <i>Food Control</i> , 2016, 60, 416-421.	2.8	70
69	Modified atmosphere generated during storage under light conditions is the main factor responsible for the quality changes of baby spinach. <i>Postharvest Biology and Technology</i> , 2016, 114, 45-53.	2.9	23
70	Monitoring generic <i>Escherichia coli</i> in reclaimed and surface water used in hydroponically cultivated greenhouse peppers and the influence of fertilizer solutions. <i>Food Control</i> , 2016, 67, 90-95.	2.8	19
71	Optimization and validation of a PMA qPCR method for <i>Escherichia coli</i> quantification in primary production. <i>Food Control</i> , 2016, 62, 150-156.	2.8	50
72	Identification of sampling points suitable for the detection of microbial contamination in fresh-cut processing lines. <i>Food Control</i> , 2016, 59, 841-848.	2.8	15

#	ARTICLE	IF	CITATIONS
73	PRE- AND POSTHARVEST STRATEGIES TO ENHANCE BIOACTIVE CONSTITUENTS OF FRUITS AND VEGETABLES. <i>Acta Horticulturae</i> , 2015, , 95-106.	0.1	2
74	Effects of salt stress on physiological and postharvest quality characteristics of different Iranian genotypes of basil. <i>Horticulture Environment and Biotechnology</i> , 2015, 56, 777-785.	0.7	42
75	Comparison of industrial precooling systems for minimally processed baby spinach. <i>Postharvest Biology and Technology</i> , 2015, 102, 1-8.	2.9	34
76	Climatic variations influence the dynamic of epiphyte bacteria of baby lettuce. <i>Food Research International</i> , 2015, 68, 54-61.	2.9	14
77	Water reconditioning by high power ultrasound combined with residual chemical sanitizers to inactivate foodborne pathogens associated with fresh-cut products. <i>Food Control</i> , 2015, 53, 29-34.	2.8	19
78	Weather Variability Influences Color and Phenolic Content of Pigmented Baby Leaf Lettuces throughout the Season. <i>Journal of Agricultural and Food Chemistry</i> , 2015, 63, 1673-1681.	2.4	62
79	Effects of oxygen-depleted atmospheres on survival and growth of <i>Listeria monocytogenes</i> on fresh-cut Iceberg lettuce stored at mild abuse commercial temperatures. <i>Food Microbiology</i> , 2015, 48, 17-21.	2.1	12
80	Ultrasound treatments improve the microbiological quality of water reservoirs used for the irrigation of fresh produce. <i>Food Research International</i> , 2015, 75, 140-147.	2.9	9
81	Potential of Electrolyzed Water as an Alternative Disinfectant Agent in the Fresh-Cut Industry. <i>Food and Bioprocess Technology</i> , 2015, 8, 1336-1348.	2.6	75
82	Postharvest treatment of table grapes with ultraviolet-C and chitosan coating preserves quality and increases stilbene content. <i>Postharvest Biology and Technology</i> , 2015, 105, 51-57.	2.9	38
83	Effect of Water Stress and Storage Time on Anthocyanins and Other Phenolics of Different Genotypes of Fresh Sweet Basil. <i>Journal of Agricultural and Food Chemistry</i> , 2015, 63, 9223-9231.	2.4	19
84	Time of day for harvest and delay before processing affect the quality of minimally processed baby spinach. <i>Postharvest Biology and Technology</i> , 2015, 110, 9-17.	2.9	17
85	Assessment of microbial risk factors and impact of meteorological conditions during production of baby spinach in the Southeast of Spain. <i>Food Microbiology</i> , 2015, 49, 173-181.	2.1	56
86	Microbial safety considerations of flooding in primary production of leafy greens: A case study. <i>Food Research International</i> , 2015, 68, 62-69.	2.9	44
87	Effect of deficit irrigation on the postharvest quality of different genotypes of basil including purple and green Iranian cultivars and a Genovese variety. <i>Postharvest Biology and Technology</i> , 2015, 100, 127-135.	2.9	29
88	Cross-contamination of <i>Escherichia coli</i> O157:H7 is inhibited by electrolyzed water combined with salt under dynamic conditions of increasing organic matter. <i>Food Microbiology</i> , 2015, 46, 471-478.	2.1	25
89	Pre- and Postharvest Preventive Measures and Intervention Strategies to Control Microbial Food Safety Hazards of Fresh Leafy Vegetables. <i>Critical Reviews in Food Science and Nutrition</i> , 2015, 55, 453-468.	5.4	226
90	Modeling growth of <i>Escherichia coli</i> O157:H7 in fresh-cut lettuce treated with neutral electrolyzed water and under modified atmosphere packaging. <i>International Journal of Food Microbiology</i> , 2014, 177, 1-8.	2.1	42

#	ARTICLE	IF	CITATIONS
91	Corrigendum to "Modelling growth of Escherichia coli O157:H7 in fresh-cut lettuce submitted to commercial process conditions: Chlorine washing and modified atmosphere packaging" [YFMIC 33 (2013) 131-138]. Food Microbiology, 2014, 41, 96.	2.1	0
92	Terminal pro-brain natriuretic peptide level determined at different times identifies transient ischaemic attack patients with atrial fibrillation. European Journal of Neurology, 2014, 21, 679-683.	1.7	15
93	Physiological, phytochemical and structural changes of multi-leaf lettuce caused by salt stress. Journal of the Science of Food and Agriculture, 2014, 94, 1592-1599.	1.7	53
94	Safety assessment of greenhouse hydroponic tomatoes irrigated with reclaimed and surface water. International Journal of Food Microbiology, 2014, 191, 97-102.	2.1	52
95	Disinfection Capacity of High-Power Ultrasound Against E. coli O157:H7 in Process Water of the Fresh-Cut Industry. Food and Bioprocess Technology, 2014, 7, 3390-3397.	2.6	17
96	Minimum free chlorine residual level required for the inactivation of Escherichia coli O157:H7 and trihalomethane generation during dynamic washing of fresh-cut spinach. Food Control, 2014, 42, 132-138.	2.8	92
97	Influence of nutrient solutions in an open-field soilless system on the quality characteristics and shelf life of fresh-cut red and green lettuces ( <i>Lactuca sativa</i> L.) in different seasons. Journal of the Science of Food and Agriculture, 2013, 93, 415-421.	1.7	21
98	Modelling growth of Escherichia coli O157:H7 in fresh-cut lettuce submitted to commercial process conditions: Chlorine washing and modified atmosphere packaging. Food Microbiology, 2013, 33, 131-138.	2.1	38
99	Operating conditions for the electrolytic disinfection of process wash water from the fresh-cut industry contaminated with E. coli o157:H7. Food Control, 2013, 29, 42-48.	2.8	38
100	Generation of trihalomethanes with chlorine-based sanitizers and impact on microbial, nutritional and sensory quality of baby spinach. Postharvest Biology and Technology, 2013, 85, 210-217.	2.9	101
101	Preharvest and postharvest factors related to off-odours of fresh-cut iceberg lettuce. Postharvest Biology and Technology, 2013, 86, 463-471.	2.9	42
102	Optimizing water management to control respiration rate and reduce browning and microbial load of fresh-cut romaine lettuce. Postharvest Biology and Technology, 2013, 80, 9-17.	2.9	21
103	Off-odour development in modified atmosphere packaged baby spinach is an unresolved problem. Postharvest Biology and Technology, 2013, 75, 75-85.	2.9	69
104	Postharvest Handling Conditions Affect Internalization of Salmonella in Baby Spinach during Washing. Journal of Food Protection, 2013, 76, 1145-1151.	0.8	19
105	Soil chemical properties, leaf mineral status and crop production in a lemon tree orchard irrigated with two types of wastewater. Agricultural Water Management, 2012, 109, 54-60.	2.4	49
106	Electrochemical disinfection: An efficient treatment to inactivate Escherichia coli O157:H7 in process wash water containing organic matter. Food Microbiology, 2012, 30, 146-156.	2.1	85
107	Baby-leaf and multi-leaf of green and red lettuces are suitable raw materials for the fresh-cut industry. Postharvest Biology and Technology, 2012, 63, 1-10.	2.9	95
108	Sensory quality, bioactive constituents and microbiological quality of green and red fresh-cut lettuces ( <i>Lactuca sativa</i> L.) are influenced by soil and soilless agricultural production systems. Postharvest Biology and Technology, 2012, 63, 16-24.	2.9	77

#	ARTICLE	IF	CITATIONS
109	Short postharvest storage under low relative humidity improves quality and shelf life of minimally processed baby spinach ( <i>Spinacia oleracea</i> L.). <i>Postharvest Biology and Technology</i> , 2012, 67, 1-9.	2.9	69
110	Influence of preharvest application of fungicides on the postharvest quality of tomato ( <i>Solanum</i> ) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 7	2.9	36
111	Long-term deficit and excess of irrigation influences quality and browning related enzymes and phenolic metabolism of fresh-cut iceberg lettuce ( <i>Lactuca sativa</i> L.). <i>Postharvest Biology and Technology</i> , 2012, 73, 37-45.	2.9	47
112	EFFECT OF IRRIGATION PRACTICES ON THE QUALITY OF FRESH-CUT LETTUCE. <i>Acta Horticulturae</i> , 2012, , 511-514.	0.1	2
113	Optimum controlled atmospheres minimise respiration rate and quality losses while increase phenolic compounds of baby carrots. <i>LWT - Food Science and Technology</i> , 2011, 44, 277-283.	2.5	26
114	Effects of water stress and rootstocks on fruit phenolic composition and physical/chemical quality in Suncrest peach. <i>Annals of Applied Biology</i> , 2011, 158, 226-233.	1.3	54
115	Low oxygen levels and light exposure affect quality of fresh-cut Romaine lettuce. <i>Postharvest Biology and Technology</i> , 2011, 59, 34-42.	2.9	131
116	HPLC-MS Analysis of Proanthocyanidin Oligomers and Other Phenolics in 15 Strawberry Cultivars. <i>Journal of Agricultural and Food Chemistry</i> , 2010, 58, 3916-3926.	2.4	226
117	Suitability of aqueous chlorine dioxide versus sodium hypochlorite as an effective sanitizer for preserving quality of fresh-cut lettuce while avoiding by-product formation. <i>Postharvest Biology and Technology</i> , 2010, 55, 53-60.	2.9	132
118	Cross-contamination of fresh-cut lettuce after a short-term exposure during pre-washing cannot be controlled after subsequent washing with chlorine dioxide or sodium hypochlorite. <i>Food Microbiology</i> , 2010, 27, 199-204.	2.1	131
119	Two-Season Study of the Influence of Regulated Deficit Irrigation and Reflective Mulch on Individual and Total Phenolic Compounds of Nectarines at Harvest and during Storage. <i>Journal of Agricultural and Food Chemistry</i> , 2010, 58, 11783-11789.	2.4	14
120	Impact of Organic Soil Amendments on Phytochemicals and Microbial Quality of Rocket Leaves ( <i>Eruca</i> ) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 7	2.4	40
121	Analysis of methodologies for the study of composition and biochemical carbohydrate changes in harvest and postharvest onion bulbs. <i>Phyton</i> , 2010, 79, 123-132.	0.4	6
122	The California, ABCD, and Unified ABCD2 Risk Scores and the Presence of Acute Ischemic Lesions on Diffusion-Weighted Imaging in TIA Patients. <i>Stroke</i> , 2009, 40, 2229-2232.	1.0	36
123	Edible coatings containing chitosan and moderate modified atmospheres maintain quality and enhance phytochemicals of carrot sticks. <i>Postharvest Biology and Technology</i> , 2009, 51, 364-370.	2.9	94
124	Prevention of <i>Escherichia coli</i> cross-contamination by different commercial sanitizers during washing of fresh-cut lettuce. <i>International Journal of Food Microbiology</i> , 2009, 133, 167-171.	2.1	161
125	Fresh-cut product sanitation and wash water disinfection: Problems and solutions. <i>International Journal of Food Microbiology</i> , 2009, 134, 37-45.	2.1	649
126	Antioxidant compounds in green and red peppers as affected by irrigation frequency, salinity and nutrient solution composition. <i>Journal of the Science of Food and Agriculture</i> , 2009, 89, 1352-1359.	1.7	50



#	ARTICLE	IF	CITATIONS
127	Quorum sensing inhibitory and antimicrobial activities of honeys and the relationship with individual phenolics. <i>Food Chemistry</i> , 2009, 115, 1337-1344.	4.2	83
128	Characterisation of polyphenols and antioxidant properties of five lettuce varieties and escarole. <i>Food Chemistry</i> , 2008, 108, 1028-1038.	4.2	427
129	Disinfection potential of ozone, ultraviolet-C and their combination in wash water for the fresh-cut vegetable industry. <i>Food Microbiology</i> , 2008, 25, 809-814.	2.1	141
130	Respiration rate response of four baby leaf Brassica species to cutting at harvest and fresh-cut washing. <i>Postharvest Biology and Technology</i> , 2008, 47, 382-388.	2.9	57
131	Role of commercial sanitizers and washing systems on epiphytic microorganisms and sensory quality of fresh-cut escarole and lettuce. <i>Postharvest Biology and Technology</i> , 2008, 49, 155-163.	2.9	162
132	A Comparative Study of Flavonoid Compounds, Vitamin C, and Antioxidant Properties of Baby Leaf <i>Brassicaceae</i> Species. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 2330-2340.	2.4	162
133	Ultraviolet-C and Induced Stilbenes Control Ochratoxigenic <i>Aspergillus</i> in Grapes. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 9990-9996.	2.4	16
134	Microbial Quality and Bioactive Constituents of Sweet Peppers from Sustainable Production Systems. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 11334-11341.	2.4	24
135	Effect of Regulated Deficit Irrigation and Crop Load on the Antioxidant Compounds of Peaches. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 3601-3608.	2.4	68
136	Fresh-cut fruit and vegetables. , 2008, , 475-504.		11
137	Impact of Wash Water Quality on Sensory and Microbial Quality, Including <i>Escherichia coli</i> Cross-Contamination, of Fresh-Cut Escarole. <i>Journal of Food Protection</i> , 2008, 71, 2514-2518.	0.8	100
138	Heterogeneous Photocatalytic Disinfection of Wash Waters from the Fresh-Cut Vegetable Industry. <i>Journal of Food Protection</i> , 2008, 71, 286-292.	0.8	36
139	Identification of New Flavonoid Glycosides and Flavonoid Profiles To Characterize Rocket Leafy Salads ( <i>Eruca vesicaria</i> and <i>Diplotaxis tenuifolia</i> ). <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 1356-1363.	2.4	64
140	Potential microbial risk factors related to soil amendments and irrigation water of potato crops. <i>Journal of Applied Microbiology</i> , 2007, 103, 2542-2549.	1.4	22
141	Elimination by ozone of <i>Shigella sonnei</i> in shredded lettuce and water. <i>Food Microbiology</i> , 2007, 24, 492-499.	2.1	108
142	Growth and bacteriocin production by lactic acid bacteria in vegetable broth and their effectiveness at reducing <i>Listeria monocytogenes</i> in vitro and in fresh-cut lettuce. <i>Food Microbiology</i> , 2007, 24, 759-766.	2.1	134
143	Impact of combined postharvest treatments (UV-C light, gaseous O <sub>3</sub> , superatmospheric O <sub>2</sub> and high T <sub>j</sub> ) on the microbial quality of fresh-cut lettuce. <i>Food Microbiology and Technology</i> , 2007, 46, 201-211.	2.9	112
144	Quality Changes and Nutrient Retention in Fresh-Cut versus Whole Fruits during Storage. <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 4284-4296.	2.4	290

#	ARTICLE	IF	CITATIONS
145	Comparison of Ozone and UV-C Treatments on the Postharvest Stilbenoid Monomer, Dimer, and Trimer Induction in Var. 'Superior' White Table Grapes. <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 4222-4228.	2.4	108
146	Minimal processing for healthy traditional foods. <i>Trends in Food Science and Technology</i> , 2006, 17, 513-519.	7.8	194
147	Effect of Ozone on the Inactivation of <i>Yersinia enterocolitica</i> and the Reduction of Natural Flora on Potatoes. <i>Journal of Food Protection</i> , 2006, 69, 2357-2363.	0.8	43
148	PHYTONUTRIENT CONTENT IN NEW APRICOT ( <i>PRUNUS ARMENIACA</i> L.) VARIETIES. <i>Acta Horticulturae</i> , 2006, , 363-368.	0.1	11
149	Controlled atmosphere preserves quality and phytonutrients in wild rocket ( <i>Diplotaxis tenuifolia</i> ). <i>Postharvest Biology and Technology</i> , 2006, 40, 26-33.	2.9	91
150	Microbial, nutritional and sensory quality of rocket leaves as affected by different sanitizers. <i>Postharvest Biology and Technology</i> , 2006, 42, 86-97.	2.9	165
151	Effect of different levels of CO <sub>2</sub> on the antioxidant content and the polyphenol oxidase activity of 'Rocha'™ pears during cold storage. <i>Journal of the Science of Food and Agriculture</i> , 2006, 86, 509-517.	1.7	15
152	Effect of different sanitizers on microbial and sensory quality of fresh-cut potato strips stored under modified atmosphere or vacuum packaging. <i>Postharvest Biology and Technology</i> , 2005, 37, 37-46.	2.9	136
153	Overview of Hazards in Fresh-Cut Produce Production: Control and Management of Food Safety Hazards. , 2005, , 155-219.		6
154	Antioxidant Capacity and Phenolic Content of Spinach As Affected by Genetics and Maturation. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 8618-8623.	2.4	89
155	Ozonated Water Extends the Shelf Life of Fresh-Cut Lettuce. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 5654-5663.	2.4	217
156	Carotenoids from New Apricot ( <i>Prunus armeniaca</i> L.) Varieties and Their Relationship with Flesh and Skin Color. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 6368-6374.	2.4	161
157	Characterization and Quantitation of Phenolic Compounds in New Apricot ( <i>Prunus armeniaca</i> L.) Varieties. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 9544-9552.	2.4	118
158	Polyphenolic compounds of Mediterranean Lamiaceae and investigation of orientational effects on <i>Acanthoscelides obtectus</i> (Say). <i>Journal of Stored Products Research</i> , 2004, 40, 395-408.	1.2	59
159	Characterization and Quantitation of Antioxidant Constituents of Sweet Pepper ( <i>Capsicum annum</i> L.). <i>Journal of Agricultural and Food Chemistry</i> , 2004, 52, 3861-3869.	2.4	417
160	Quality improvement of <i>Pleurotus</i> mushrooms by modified atmosphere packaging and moisture absorbers. <i>Postharvest Biology and Technology</i> , 2003, 28, 169-179.	2.9	106
161	Comparative study of six pear cultivars in terms of their phenolic and vitamin C contents and antioxidant capacity. <i>Journal of the Science of Food and Agriculture</i> , 2003, 83, 995-1003.	1.7	128
162	l-Galactono-1 $\beta$ -Lactone Dehydrogenase Activity and Vitamin C Content in Fresh-Cut Potatoes Stored under Controlled Atmospheres. <i>Journal of Agricultural and Food Chemistry</i> , 2003, 51, 4296-4302.	2.4	16

#	ARTICLE	IF	CITATIONS
163	Influence of Industrial Processing on Orange Juice Flavanone Solubility and Transformation to Chalcones under Gastrointestinal Conditions. <i>Journal of Agricultural and Food Chemistry</i> , 2003, 51, 3024-3028.	2.4	65
164	MAP, product safety and nutritional quality. , 2003, , 208-230.		5
165	QUANTIFICATION OF CHLOROGENIC ACID AND POLYPHENOL OXIDASE ACTIVITY IN 'ROCHA' PEAR AFTER CA-LONG STORAGE. <i>Acta Horticulturae</i> , 2003, , 405-408.	0.1	3
166	EFFECTS OF LOW-OXYGEN AND HIGH-CARBON DIOXIDE ATMOSPHERE ON POSTHARVEST QUALITY OF ARTICHOKES. <i>Acta Horticulturae</i> , 2003, , 385-388.	0.1	1
167	Modified atmosphere packaging (MAP). , 2002, , 342-370.		10
168	Effect of Processing Techniques at Industrial Scale on Orange Juice Antioxidant and Beneficial Health Compounds. <i>Journal of Agricultural and Food Chemistry</i> , 2002, 50, 5107-5114.	2.4	171
169	Phenolic Compounds and Related Enzymes Are Not Rate-Limiting in Browning Development of Fresh-Cut Potatoes. <i>Journal of Agricultural and Food Chemistry</i> , 2002, 50, 3015-3023.	2.4	219
170	Antioxidant Capacities, Phenolic Compounds, Carotenoids, and Vitamin C Contents of Nectarine, Peach, and Plum Cultivars from California. <i>Journal of Agricultural and Food Chemistry</i> , 2002, 50, 4976-4982.	2.4	679
171	Induction of Antioxidant Flavonol Biosynthesis in Fresh-Cut Potatoes. Effect of Domestic Cooking. <i>Journal of Agricultural and Food Chemistry</i> , 2002, 50, 5925-5931.	2.4	127
172	Antioxidant Capacity and Phenolic Content of Spinach As Affected by Genetics and Growing Season. <i>Journal of Agricultural and Food Chemistry</i> , 2002, 50, 5891-5896.	2.4	142
173	Quality changes in fresh cut tomato as affected by modified atmosphere packaging. <i>Postharvest Biology and Technology</i> , 2002, 25, 199-207.	2.9	82
174	Vitamin C retention in fresh-cut potatoes. <i>Postharvest Biology and Technology</i> , 2002, 26, 75-84.	2.9	83
175	Influence of modified atmosphere packaging on quality, vitamin C and phenolic content of artichokes ( <i>Cynara scolymus</i> L.). <i>European Food Research and Technology</i> , 2002, 215, 21-27.	1.6	34
176	HPLC-ESI/MS Analysis of Phenolic Compounds in Nectarines, Peaches, and Plums. <i>Journal of Agricultural and Food Chemistry</i> , 2001, 49, 4748-4760.	2.4	594
177	The effect of storage temperatures on vitamin C and phenolics content of artichoke ( <i>Cynara scolymus</i> ) <a href="#">Tj ETQq1 1 0,784314 rgBT /Over</a>	2.7	59
178	In Vitro Availability of Flavonoids and Other Phenolics in Orange Juice. <i>Journal of Agricultural and Food Chemistry</i> , 2001, 49, 1035-1041.	2.4	239
179	EFFECT OF STORAGE TEMPERATURE ON QUALITY OF ARTICHOKES ( <i>Cynara scolymus</i> L.). <i>Acta Horticulturae</i> , 2001, , 555-557.	0.1	3
180	MODIFIED ATMOSPHERE PACKAGING OF FRESH-CUT TOMATO. <i>Acta Horticulturae</i> , 2001, , 703-704.	0.1	6

#	ARTICLE	IF	CITATIONS
181	Antioxidant phenolic metabolites from fruit and vegetables and changes during postharvest storage and processing. <i>Studies in Natural Products Chemistry</i> , 2000, 23, 739-795.	0.8	58
182	Antioxidant Activity of Pomegranate Juice and Its Relationship with Phenolic Composition and Processing. <i>Journal of Agricultural and Food Chemistry</i> , 2000, 48, 4581-4589.	2.4	1,957
183	Browning susceptibility of minimally processed Baby and Romaine lettuces. <i>European Food Research and Technology</i> , 1999, 209, 52-56.	1.6	63
184	Keeping quality of fresh-cut tomato. <i>Postharvest Biology and Technology</i> , 1999, 17, 153-162.	2.9	113
185	Effect of Postharvest Storage and Processing on the Antioxidant Constituents (Flavonoids and) Tj ETQq1 1 0.784314 rgBT /Overlock 10	2.4	340
186	Modified-atmosphere packaging of minimally processed "Lollo Rosso" ( <i>Lactuca sativa</i> ). <i>European Food Research and Technology</i> , 1998, 206, 350-354.	0.6	33
187	Improving the keeping quality of pomegranate fruit by intermittent warming. <i>European Food Research and Technology</i> , 1998, 207, 316-321.	0.6	29
188	Effect of Modified Atmosphere Packaging on the Flavonoids and Vitamin C Content of Minimally Processed Swiss Chard ( <i>Beta vulgaris</i> Subspeciescycla). <i>Journal of Agricultural and Food Chemistry</i> , 1998, 46, 2007-2012.	2.4	104
189	RevisiÃ³n: El pardeamiento enzimÃ¡tico en frutas y hortalizas mÃ¡nimamente procesadas Review: Enzymatic browning in minimally processed fruit and vegetables. <i>Food Science and Technology International</i> , 1998, 4, 377-389.	1.1	87
190	POSTHARVEST PHYSIOLOGY AND QUALITY MAINTENANCE OF FRESH-CUT PEARS. <i>Acta Horticulturae</i> , 1998, , 231-236.	0.1	54
191	Effect of Carbon Dioxide on Anthocyanins, Phenylalanine Ammonia Lyase and Glucosyltransferase in the Arils of Stored Pomegranates. <i>Journal of the American Society for Horticultural Science</i> , 1998, 123, 136-140.	0.5	91
192	Effect of Selected Browning Inhibitors on Phenolic Metabolism in Stem Tissue of Harvested Lettuce. <i>Journal of Agricultural and Food Chemistry</i> , 1997, 45, 583-589.	2.4	135
193	Phenolic Metabolites in Red Pigmented Lettuce ( <i>Lactuca sativa</i> ). Changes with Minimal Processing and Cold Storage. <i>Journal of Agricultural and Food Chemistry</i> , 1997, 45, 4249-4254.	2.4	163
194	Changes in Strawberry Anthocyanins and Other Polyphenols in Response to Carbon Dioxide Treatments. <i>Journal of Agricultural and Food Chemistry</i> , 1997, 45, 1662-1667.	2.4	256
195	Inhibition of superficial scald in apples by wounding: changes in lipids and phenolics. <i>Postharvest Biology and Technology</i> , 1997, 12, 203-212.	2.9	23
196	Organic acids as browning inhibitors on harvested "Baby" lettuce and endive. <i>European Food Research and Technology</i> , 1997, 205, 375-379.	0.6	9
197	Determination of phenolic compounds in honeys with different floral origin by capillary zone electrophoresis. <i>Food Chemistry</i> , 1997, 60, 79-84.	4.2	116
198	Anthocyanins and flavonoids from shredded red onion and changes during storage in perforated films. <i>Food Research International</i> , 1996, 29, 389-395.	2.9	60

#	ARTICLE	IF	CITATIONS
199	Minimally Processed Pomegranate Seeds. LWT - Food Science and Technology, 1996, 29, 708-713.	2.5	62
200	Quality changes in pomegranates during ripening and cold storage. Zeitschrift Fur Lebensmittel-Untersuchung Und -Forschung, 1996, 202, 481-485.	0.7	28
201	Floral nectar phenolics as biochemical markers for the botanical origin of heather honey. Zeitschrift Fur Lebensmittel-Untersuchung Und -Forschung, 1996, 202, 40-44.	0.7	91
202	Inhibition of Browning of Harvested Head Lettuce. Journal of Food Science, 1996, 61, 314-316.	1.5	48
203	Minimal Processing and Modified Atmosphere Packaging Effects on Pigmentation of Pomegranate Seeds. Journal of Food Science, 1996, 61, 161-164.	1.5	57
204	Changes in pomegranate juice pigmentation during ripening. Journal of the Science of Food and Agriculture, 1995, 68, 77-81.	1.7	143
205	Plant Phenolic Metabolites and Floral Origin of Rosemary Honey. Journal of Agricultural and Food Chemistry, 1995, 43, 2833-2838.	2.4	121
206	Analysis of phenolic compounds in Spanish red wines by capillary zone electrophoresis. Zeitschrift Fur Lebensmittel-Untersuchung Und -Forschung, 1995, 200, 278-281.	0.7	27
207	Influence of cultivar, maturity stage and geographical location on the juice pigmentation of Tunisian pomegranates. Zeitschrift Fur Lebensmittel-Untersuchung Und -Forschung, 1995, 201, 361-364.	0.7	31
208	Flavonoid patterns of French honeys with different floral origin. Apidologie, 1995, 26, 53-60.	0.9	57
209	Micellar Electrokinetic Capillary Chromatography of Methylated Flavone Aglycones. Journal of Liquid Chromatography and Related Technologies, 1995, 18, 3007-3019.	0.9	15
210	Separation of honey flavonoids by micellar electrokinetic capillary chromatography. Journal of Chromatography A, 1994, 669, 268-274.	1.8	65
211	Distribution of flavonoid aglycones and glycosides in Sideritis species from the canary islands and madeira. Phytochemistry, 1993, 34, 227-232.	1.4	26
212	Distribution of 8-Hydroxyflavone glycosides and flavonoid aglycones in some Spanish Sideritis species. Biochemical Systematics and Ecology, 1993, 21, 487-497.	0.6	12
213	Flavonoid in some Leonurus, Chaiturus and Panzerina species (Lamiaceae). Biochemical Systematics and Ecology, 1993, 21, 531-532.	0.6	7
214	Flavonoid p-coumaroylglucosides and 8-hydroxyflavone allosylglucosides in some labiatae. Phytochemistry, 1992, 31, 3097-3102.	1.4	71
215	External and vacuolar flavonoids from Nepeta transcaucasica. Biochemical Systematics and Ecology, 1992, 20, 589-590.	0.6	8
216	Correlations between flavonoid composition and infrageneric taxonomy of some european Galeopsis species. Phytochemistry, 1991, 30, 3311-3314.	1.4	12

#	ARTICLE	IF	CITATIONS
217	Flavonoids from some Yugoslavian <i>Micromeria</i> species: Chemotaxonomical aspects. <i>Biochemical Systematics and Ecology</i> , 1991, 19, 697-698.	0.6	18
218	An HPLC technique for flavonoid analysis in honey. <i>Journal of the Science of Food and Agriculture</i> , 1991, 56, 49-56.	1.7	109
219	Distribution of 6-hydroxy-, 6-methoxy- and 8-hydroxyflavone glycosides in the labiatae, the scrophulariaceae and related families. <i>Phytochemistry</i> , 1988, 27, 2631-2645.	1.4	70
220	The distribution of methylated flavones in the Lamiaceae. <i>Biochemical Systematics and Ecology</i> , 1988, 16, 43-46.	0.6	48
221	A chemotaxonomic study of flavonoids from european <i>teucrium</i> species. <i>Phytochemistry</i> , 1986, 25, 2811-2816.	1.4	90
222	Flavonoid Aglycones and Glycosides from <i>Teucrium gnaphalodes</i> . <i>Journal of Natural Products</i> , 1985, 48, 859-860.	1.5	23
223	Harvest maturity indicators of leafy vegetables. <i>Stewart Postharvest Review</i> , 0, 8, 1-9.	0.7	21
224	Water management and its effect on the postharvest quality of fresh-cut vegetables. <i>Stewart Postharvest Review</i> , 0, 9, 1-8.	0.7	3