

# Marcos Gino Fernandes

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9277491/publications.pdf>

Version: 2024-02-01

32  
papers

248  
citations

1163117

8  
h-index

1058476

14  
g-index

34  
all docs

34  
docs citations

34  
times ranked

246  
citing authors

#	ARTICLE	IF	CITATIONS
1	Distribuição espacial de <i>Alabama argillacea</i> (Hübner) (Lepidoptera: Noctuidae) em algodoeiro. <i>Neotropical Entomology</i> , 2003, 32, 107-115.	1.2	39
2	<i>Harmonia axyridis</i> (Coleoptera: Coccinellidae) Exhibits No Preference between Bt and Non-Bt Maize Fed <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae). <i>PLoS ONE</i> , 2012, 7, e44867.	2.5	24
3	Distribuição espacial de <i>Aphis gossypii</i> (Glover) (Hemiptera, Aphididae) e <i>Bemisia tabaci</i> (Gennadius) biótipo B (Hemiptera, Aleyrodidae) em algodoeiro Bt e não-Bt. <i>Revista Brasileira De Entomologia</i> , 2010, 54, 136-143.	0.4	23
4	Susceptibility of Cotton Boll Weevil (Coleoptera: Curculionidae) to Spinosyns. <i>Journal of Economic Entomology</i> , 2019, 112, 1688-1694.	1.8	13
5	Sequential sampling of <i>Aphis gossypii</i> Glover (Hemiptera: Aphididae) and <i>Frankliniella schultzei</i> Trybom (Thysanoptera: Thripidae) on cotton crop. <i>Neotropical Entomology</i> , 2011, 40, 258-263.	1.2	12
6	Spatial distribution and sampling plan of the phytophagous stink bug complex in different soybean production systems. <i>Journal of Applied Entomology</i> , 2019, 143, 236-249.	1.8	9
7	Resistance of maize landraces from Brazil to fall armyworm (Lepidoptera: Noctuidae) in the winter and summer seasons. <i>Bragantia</i> , 2020, 79, 377-386.	1.3	9
8	Spatial Distribution of <i>Euschistus heros</i> (Hemiptera: Pentatomidae) in Cotton ( <i>Gossypium hirsutum</i> ) Tj ETQq 0 0 0 rgBT /Overlock 10 Tf 5	0.8	8
9	Spatial distribution and control levels of <i>Anastrepha</i> spp. (Diptera: Tephritidae) in guava orchards. <i>Anais Da Academia Brasileira De Ciencias</i> , 2019, 91, e20180428.	0.8	8
10	Amostragem seqüencial de <i>Alabama argillacea</i> (Hübner) (Lepidoptera: Noctuidae) em algodoeiro. <i>Neotropical Entomology</i> , 2003, 32, 117-122.	1.2	7
11	Spatial Distribution of Adult <i>Anthonomus grandis</i> Boheman (Coleoptera: Curculionidae) and Damage to Cotton Flower Buds Due to Feeding and Oviposition. <i>Neotropical Entomology</i> , 2017, 46, 442-451.	1.2	7
12	First confirmed record of <i>Harmonia axyridis</i> (Pallas, 1773) (Coleoptera: Coccinellidae) in the state of Mato Grosso do Sul, Brazil. <i>Check List</i> , 2011, 7, 476.	0.4	6
13	Species Richness and Community Composition of Ants and Beetles in Bt and non-Bt Maize Fields. <i>Environmental Entomology</i> , 2019, 48, 1095-1103.	1.4	5
14	Resistance of Cowpea Genotypes to <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae) and Its Relationship to Resistance-Related Enzymes. <i>Journal of Economic Entomology</i> , 2020, 113, 2521-2529.	1.8	5
15	Sequential sampling of <i>Bemisia tuberculata</i> (Bondar, 1923) (Hemiptera: Aleyrodidae) on cassava crop. <i>Anais Da Academia Brasileira De Ciencias</i> , 2014, 86, 889-896.	0.8	5
16	Leaf Chlorophyll Content and Agronomic Performance of Bt and Non-Bt Soybean. <i>Journal of Agricultural Science</i> , 2013, 5, .	0.2	4
17	Disposição espacial e injúrias da lagarta-do-cartucho <i>Spodoptera frugiperda</i> (J. E. Smith) (Lepidoptera:) Tj ETQq 1 1 0.7843 14 rgBT (C	0.4	4
18	Sequential Sampling Plan of <i>Anthonomus grandis</i> (Coleoptera: Curculionidae) in Cotton Plants. <i>Journal of Economic Entomology</i> , 2017, 110, 763-769.	1.8	4

#	ARTICLE	IF	CITATIONS
19	Biology of <i>Trichogramma pretiosum</i> (Hymenoptera: Trichogrammatidae) Fed Transgenic Maize Pollen. Florida Entomologist, 2017, 100, 653-656.	0.5	4
20	Antibiosis Levels to <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae) in Cowpea Commercial Cultivars and Landrace Varieties. Journal of Economic Entomology, 2019, 112, 1941-1945.	1.8	4
21	Sand Fly Fauna, Spatial Distribution of <i>Lutzomyia longipalpis</i> (Diptera: Psychodidae), and Climate Factors in Dourados, Brazil. Journal of Medical Entomology, 2021, 58, 1952-1957.	1.8	3
22	Maize resistance to <i>Spodoptera frugiperda</i> and its relationship to landrace variety, plant stage, and larval origin. Entomologia Experimentalis Et Applicata, 2021, 169, 711-720.	1.4	3
23	Distribuição espacial de <i>Spodoptera frugiperda</i> (J. E. Smith) (Lepidoptera, noctuidae) em milho convencional e Bt. EntomoBrasilis, 2017, 10, 89-93.	0.2	3
24	DISTRIBUCIÓN ESPACIAL DE <i>Vatiga</i> spp. (HEMIPTERA: TINGIDAE) EN EL CULTIVO DE YUCA. Acta Biologica Colombiana, 2015, 21, .	0.4	3
25	Spatial distribution of adult <i>Anthonomus grandis</i> Boheman (Coleoptera: Curculionidae) and buds with feeding punctures on conventional and Bt cotton. Semina:Ciencias Agrarias, 2013, 34, .	0.3	2
26	Influence of Bt Maize on Diversity and Composition of Non-target Arthropod Species. Journal of Agricultural Science, 2019, 11, 201.	0.2	2
27	Spatial distribution and sequential sampling plan for <i>Diceraeus melacanthus</i> (Hemiptera: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 5	2.1	2
28	Above- and belowground resistance in Brazilian maize varieties under attack of <i>Spodoptera frugiperda</i> and <i>Diabrotica speciosa</i> . Entomologia Experimentalis Et Applicata, 0, , .	1.4	2
29	Himenópteros parasitoides asociados a ninfas de <i>Triozoida limbata</i> na cultura da goiabeira, em Vinhema, MS, Brasil. Ciencia Rural, 2015, 45, 19-21.	0.5	1
30	Sequential sampling plan for fruit fly species of the Genus <i>Anastrepha</i> (Diptera: Tephritidae) in guava orchards. Anais Da Academia Brasileira De Ciencias, 2018, 90, 3607-3614.	0.8	1
31	Spatial Distribution of Eggs of <i>Alabama argillacea</i> H&#246;bnner and <i>Heliothis virescens</i> Fabricius (Lepidoptera: Noctuidae) on Bt and non-Bt Cotton. Anais Da Academia Brasileira De Ciencias, 2015, 87, 2243-2253.	0.8	0
32	ESTUDIO DE LA ENTOMOFAUNA DE GENOTIPOS DE FRIJOL POR MEDIO DE ANÁLISIS MULTIVARIADO. Acta Biologica Colombiana, 2021, 26, 312-317.	0.4	0