

Marcos Gino Fernandes

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9277491/publications.pdf>

Version: 2024-02-01

32

papers

248

citations

1163117

8

h-index

1058476

14

g-index

34

all docs

34

docs citations

34

times ranked

246

citing authors

#	ARTICLE	IF	CITATIONS
1	Spatial distribution and sequential sampling plan for <i>Diceraeus melacanthus</i> (Hemiptera: Tj ETQq1 1 0.784314 rgBT /Overlock 2.1	0.784314 rgBT /Overlock 2.1	10 Tf 50
2	Sand Fly Fauna, Spatial Distribution of <i>Lutzomyia longipalpis</i> (Diptera: Psychodidae), and Climate Factors in Dourados, Brazil. Journal of Medical Entomology, 2021, 58, 1952-1957.	1.8	3
3	ESTUDIO DE LA ENTOMOFALINA DE GENOTIPOS DE FRIJOL POR MEDIO DE ANÁLISIS MULTIVARIADO. Acta Biologica Colombiana, 2021, 26, 312-317.	0.4	0
4	Maize resistance to <i>Spodoptera frugiperda</i> and its relationship to landrace variety, plant stage, and larval origin. Entomologia Experimentalis Et Applicata, 2021, 169, 711-720.	1.4	3
5	Resistance of Cowpea Genotypes to <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae) and Its Relationship to Resistance-Related Enzymes. Journal of Economic Entomology, 2020, 113, 2521-2529.	1.8	5
6	Resistance of maize landraces from Brazil to fall armyworm (Lepidoptera: Noctuidae) in the winter and summer seasons. Bragantia, 2020, 79, 377-386.	1.3	9
7	Spatial distribution and control levels of <i>Anastrepha</i> spp. (Diptera: Tephritidae) in guava orchards. Anais Da Academia Brasileira De Ciencias, 2019, 91, e20180428.	0.8	8
8	Species Richness and Community Composition of Ants and Beetles in Bt and non-Bt Maize Fields. Environmental Entomology, 2019, 48, 1095-1103.	1.4	5
9	Antibiosis Levels to <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae) in Cowpea Commercial Cultivars and Landrace Varieties. Journal of Economic Entomology, 2019, 112, 1941-1945.	1.8	4
10	Susceptibility of Cotton Boll Weevil (Coleoptera: Curculionidae) to Spinosyns. Journal of Economic Entomology, 2019, 112, 1688-1694.	1.8	13
11	Spatial distribution and sampling plan of the phytophagous stink bug complex in different soybean production systems. Journal of Applied Entomology, 2019, 143, 236-249.	1.8	9
12	Influence of Bt Maize on Diversity and Composition of Non-target Arthropod Species. Journal of Agricultural Science, 2019, 11, 201.	0.2	2
13	Spatial Distribution of <i>Euschistus heros</i> (Hemiptera: Pentatomidae) in Cotton (<i>Gossypium hirsutum</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 0.8	0.784314 rgBT /Overlock 0.8	10
14	Sequential sampling plan for fruit fly species of the Genus <i>Anastrepha</i> (Diptera: Tephritidae) in guava orchards. Anais Da Academia Brasileira De Ciencias, 2018, 90, 3607-3614.	0.8	1
15	Sequential Sampling Plan of <i>Anthonomus grandis</i> (Coleoptera: Curculionidae) in Cotton Plants. Journal of Economic Entomology, 2017, 110, 763-769.	1.8	4
16	Spatial Distribution of Adult <i>Anthonomus grandis</i> Boheman (Coleoptera: Curculionidae) and Damage to Cotton Flower Buds Due to Feeding and Oviposition. Neotropical Entomology, 2017, 46, 442-451.	1.2	7
17	Biology of <i>Trichogramma pretiosum</i> (Hymenoptera: Trichogrammatidae) Fed Transgenic Maize Pollen. Florida Entomologist, 2017, 100, 653-656.	0.5	4
18	Distribuição espacial de <i>Spodoptera frugiperda</i> (J. E. Smith) (Lepidoptera, noctuidae) em milho convencional e Bt. EntomoBrasilis, 2017, 10, 89-93.	0.2	3

#	ARTICLE	IF	CITATIONS
19	Spatial Distribution of Eggs of Alabama argillacea HÃ¼bner and Heliothis virescens Fabricius (Lepidoptera: Noctuidae) on Bt and non-Bt Cotton. Anais Da Academia Brasileira De Ciencias, 2015, 87, 2243-2253.	0.8	0
20	Himenópteros parasitoides associados a ninfas de Triozoida limbata na cultura da goiabeira, em Ivinhema, MS, Brasil. Ciencia Rural, 2015, 45, 19-21.	0.5	1
21	DISTRIBUCIÓN ESPACIAL DE Vatiga spp. (HEMIPTERA: TINGIDAE) EN EL CULTIVO DE YUCA. Acta Biologica Colombiana, 2015, 21, .	0.4	3
22	DisposiÃ§Ã£o espacial e injÃ³rias da lagarta-do-cartucho Spodoptera frugiperda (J. E. Smith) (Lepidoptera:) Tj ETQq0.00 rgBT ₄ /Overlock	0.4	1
23	Sequential sampling of Bemisia tuberculata (Bondar, 1923) (Hemiptera: Aleyrodidae) on cassava crop. Anais Da Academia Brasileira De Ciencias, 2014, 86, 889-896.	0.8	5
24	Spatial distribution of adult Anthonomus grandis Boheman (Coleoptera: Curculionidae) and buds with feeding punctures on conventional and Bt cotton. Semina: Ciencias Agrarias, 2013, 34, .	0.3	2
25	Leaf Chlorophyll Content and Agronomic Performance of Bt and Non-Bt Soybean. Journal of Agricultural Science, 2013, 5, .	0.2	4
26	Harmonia axyridis (Coleoptera: Coccinellidae) Exhibits No Preference between Bt and Non-Bt Maize Fed Spodoptera frugiperda (Lepidoptera: Noctuidae). PLoS ONE, 2012, 7, e44867.	2.5	24
27	Sequential sampling of Aphis gossypii Glover (Hemiptera: Aphididae) and Frankliniella schultzei Trybom (Thysanoptera: Thripidae) on cotton crop. Neotropical Entomology, 2011, 40, 258-263.	1.2	12
28	First confirmed record of Harmonia axyridis (Pallas, 1773) (Coleoptera: Coccinellidae) in the state of Mato Grosso do Sul, Brazil. Check List, 2011, 7, 476.	0.4	6
29	DistribuiÃ§Ã£o espacial de Aphis gossypii (Glover) (Hemiptera, Aphididae) e Bemisia tabaci (Gennadius) biÃ³tipo B (Hemiptera, Aleyrodidae) em algodoeiro Bt e nÃ£o-Bt. Revista Brasileira De Entomologia, 2010, 54, 136-143.	0.4	23
30	DistribuiÃ§Ã£o espacial de Alabama argillacea (HÃ¼bner) (Lepidoptera: Noctuidae) em algodoeiro. Neotropical Entomology, 2003, 32, 107-115.	1.2	39
31	Amostragem seqÃ¼encial de Alabama argillacea (HÃ¼bner) (Lepidoptera: Noctuidae) em algodoeiro. Neotropical Entomology, 2003, 32, 117-122.	1.2	7
32	Aboveâ€¢and belowground resistance in Brazilian maize varieties under attack of <i>Spodoptera frugiperda</i> and <i>Diabrotica speciosa</i> . Entomologia Experimentalis Et Applicata, 0, , .	1.4	2