Gudrun Marteinsdottir

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/927923/publications.pdf

Version: 2024-02-01

71 papers 2,620 citations

28 h-index 197818 49 g-index

71 all docs

71 docs citations

71 times ranked 1921 citing authors

#	Article	IF	CITATIONS
1	New insight into trophic niche partitioning and diet of mackerel (Scomber scombrus) and herring (Clupea harengus) in Icelandic waters. ICES Journal of Marine Science, 2021, 78, 1485-1499.	2.5	1
2	Egg size and density estimates for three gadoids in Icelandic waters and their implications for the vertical distribution of eggs along a stratified water column. Journal of Marine Systems, 2020, 204, 103290.	2.1	3
3	Isolating the influence of ontogeny helps predict island-wide variability in fish otolith chemistry. Reviews in Fish Biology and Fisheries, 2020, 30, 173-202.	4.9	14
4	Dispersal of juvenile cod (Gadus morhua L.) in Icelandic waters. Fisheries Research, 2020, 232, 105721.	1.7	2
5	Species richness in North Atlantic fish: Process concealed by pattern. Global Ecology and Biogeography, 2020, 29, 842-856.	5.8	11
6	Diet and feeding strategy of Northeast Atlantic mackerel (Scombrus scomber) in Icelandic waters. PLoS ONE, 2019, 14, e0225552.	2.5	6
7	Determination of bioactive properties of food grade extracts from Icelandic edible brown seaweed sugar kelp (Saccharina latissima) with in vitro human cell cultures (THP-1) Functional Foods in Health and Disease, 2019, 9, 1.	0.6	7
8	Can collective memories shape fish distributions? A test, linking spaceâ€time occurrence models and population demographics. Ecography, 2018, 41, 938-957.	4.5	11
9	Otolith shape differences between ecotypes of Icelandic cod (Gadus morhua) with known migratory behaviour inferred from data storage tags. Canadian Journal of Fisheries and Aquatic Sciences, 2017, 74, 2122-2130.	1.4	11
10	Bioactive effect of sulphated polysaccharides derived from orange-footed sea cucumber (Cucumaria) Tj ETQq0	0 0 rgBT /	Overlock 10 Tf
		2.7	8
11	Distribution, maturity and population structure of Meganyctiphanes norvegica and Thysanoessa inermis around Iceland in spring. PLoS ONE, 2017, 12, e0187360.	2.5	9
11	Distribution, maturity and population structure of Meganyctiphanes norvegica and Thysanoessa inermis around Iceland in spring. PLoS ONE, 2017, 12, e0187360. Evaluation of bioactivity of fucoidan from laminaria with in vitro human cell cultures (THP-1). Functional Foods in Health and Disease, 2017, 7, 688.	2.7	0
	inermis around Iceland in spring. PLoS ONE, 2017, 12, e0187360. Evaluation of bioactivity of fucoidan from laminaria with in vitro human cell cultures (THP-1).	2.5	9
12	inermis around Iceland in spring. PLoS ONE, 2017, 12, e0187360. Evaluation of bioactivity of fucoidan from laminaria with in vitro human cell cultures (THP-1). Functional Foods in Health and Disease, 2017, 7, 688. Abundance and distribution of early life stages of krill around Iceland during spring. Marine Biology	2.5	9
12	inermis around Iceland in spring. PLoS ONE, 2017, 12, e0187360. Evaluation of bioactivity of fucoidan from laminaria with in vitro human cell cultures (THP-1). Functional Foods in Health and Disease, 2017, 7, 688. Abundance and distribution of early life stages of krill around Iceland during spring. Marine Biology Research, 2016, 12, 864-873. Nile perch and the transformation of Lake Victoria. African Journal of Aquatic Science, 2016, 41,	2.5 0.6	9 6 4
12 13 14	inermis around Iceland in spring. PLoS ONE, 2017, 12, e0187360. Evaluation of bioactivity of fucoidan from laminaria with in vitro human cell cultures (THP-1). Functional Foods in Health and Disease, 2017, 7, 688. Abundance and distribution of early life stages of krill around Iceland during spring. Marine Biology Research, 2016, 12, 864-873. Nile perch and the transformation of Lake Victoria. African Journal of Aquatic Science, 2016, 41, 127-142. Environmental and climactic effects of chlorophyll-a variability around Iceland using reconstructed	2.5 0.6 0.7	9 6 4 37
12 13 14 15	inermis around Iceland in spring. PLoS ONE, 2017, 12, e0187360. Evaluation of bioactivity of fucoidan from laminaria with in vitro human cell cultures (THP-1). Functional Foods in Health and Disease, 2017, 7, 688. Abundance and distribution of early life stages of krill around Iceland during spring. Marine Biology Research, 2016, 12, 864-873. Nile perch and the transformation of Lake Victoria. African Journal of Aquatic Science, 2016, 41, 127-142. Environmental and climactic effects of chlorophyll-a variability around Iceland using reconstructed satellite data fields. Journal of Marine Systems, 2016, 163, 31-42. The cross-shore distribution of plankton and particles southwest of Iceland observed with a Video	2.5 0.6 0.7 1.1	9 6 4 37

#	Article	IF	CITATIONS
19	Evidence from data storage tags for the presence of lunar and semi-lunar behavioral cycles in spawning Atlantic cod. Environmental Biology of Fishes, 2015, 98, 1767-1776.	1.0	8
20	Live holding of Nephrops norvegicus (Linnaeus, 1758) in land-based facilities: Health and condition effects. Marine Biology Research, 2015, 11, 603-612.	0.7	2
21	Spatial and temporal variation in the distribution and density of pelagic fish species in Lake Victoria, East Africa. Aquatic Ecosystem Health and Management, 2014, 17, 52-61.	0.6	24
22	To glue or not to glue? Reassembling broken otoliths for population discrimination. Journal of Fish Biology, 2014, 84, 1626-1633.	1.6	4
23	Long-term changes of euphausiids in shelf and oceanic habitats southwest, south and southeast of Iceland. Journal of Plankton Research, 2014, 36, 1262-1278.	1.8	12
24	Spawning behavior in Atlantic cod: analysis by use of data storage tags. Marine Ecology - Progress Series, 2014, 506, 279-290.	1.9	5
25	Spatio-temporal effects of stray hatchery-reared Atlantic salmon Salmo salar on population genetic structure within a 21Âkm-long Icelandic river system. Conservation Genetics, 2013, 14, 1217-1231.	1.5	6
26	Distribution and exploitation of Nile perch Lates niloticus in relation to stratification in Lake Victoria, East Africa. Journal of Great Lakes Research, 2013, 39, 466-475.	1.9	24
27	The circulation of Icelandic waters – a modelling study. Ocean Science, 2013, 9, 931-955.	3.4	58
28	Identification of stock components using morphological markers. Journal of Fish Biology, 2012, 81, 1447-1462.	1.6	13
29	The composition of adult overwintering and juvenile aggregations of Atlantic cod (<i>) Gadus) Tj ETQq1 1 0.78431 Journal of Fisheries and Aquatic Sciences, 2012, 69, 307-320.</i>	.4 rgBT /O 1.4	
30	Characterization of Atlantic Cod Spawning Habitat and Behavior in Icelandic Coastal Waters. PLoS ONE, 2012, 7, e51321.	2.5	19
31	Evidence of Segregated Spawning in a Single Marine Fish Stock: Sympatric Divergence of Ecotypes in Icelandic Cod?. PLoS ONE, 2011, 6, e17528.	2.5	63
32	The use of otolith chemistry to determine the juvenile source of spawning cod in Icelandic waters. ICES Journal of Marine Science, 2011, 68, 98-106.	2.5	26
33	Groundfish species diversity and assemblage structure in Icelandic waters during recent years of warming. Fisheries Oceanography, 2010, 19, 42-62.	1.7	44
34	Fecundity and growth of Atlantic cod (Gadus morhua L.) along a latitudinal gradient. Fisheries Research, 2010, 104, 45-55.	1.7	49
35	Temporal trends in probabilistic maturation reaction norms and growth of Atlantic cod (Gadus) Tj ETQq $1\ 1\ 0.784$	314 rgBT 1.4	/Oyerlock 10
36	Evidence of phenotypic plasticity and local adaption in metabolic rates between components of the Icelandic cod (Gadus morhua L.) stock. Environmental Biology of Fishes, 2009, 86, 361-370.	1.0	19

#	Article	IF	Citations
37	Contrasting trends in two condition indices: bathymetric and spatial variation in autumn condition of Icelandic cod <i>Gadus morhua</i> . Journal of Fish Biology, 2009, 75, 282-289.	1.6	18
38	Abundance and growth of larval and early juvenile cod (Gadus morhua) in relation to variable environmental conditions west of Iceland. Deep-Sea Research Part II: Topical Studies in Oceanography, 2009, 56, 1992-2000.	1.4	10
39	The evaluation of reference points and stock productivity in the context of alternative indices of stock reproductive potential. Canadian Journal of Fisheries and Aquatic Sciences, 2009, 66, 404-414.	1.4	43
40	Are Vertical Behaviour Patterns Related to the Pantophysin Locus in the Atlantic Cod (Gadus morhua) Tj ETQq0 (0 0 rgBT /0	Overlock 10 Tf
41	Growth, maturity and fecundity of female spotted wolffish <i>Anarhichas minor </i> in Icelandic waters. Journal of Fish Biology, 2008, 73, 1393-1406.	1.6	9
42	Spatial and temporal trends in condition of Atlantic cod Gadus morhua on the Icelandic shelf. Marine Ecology - Progress Series, 2008, 362, 261-277.	1.9	38
43	Distribution and migration of saithe (Pollachius virens) around Iceland inferred from mark-recapture studies. ICES Journal of Marine Science, 2007, 64, 1006-1016.	2.5	24
44	Drift probabilities for Icelandic cod larvae. ICES Journal of Marine Science, 2007, 64, 49-59.	2.5	22
45	Contribution of different spawning components to the mixed stock fishery for cod in Icelandic waters. ICES Journal of Marine Science, 2007, 64, 1749-1759.	2.5	22
46	Collapse of the fishery for Iceland scallop (Chlamys islandica) in Breidafjordur, West Iceland. ICES Journal of Marine Science, 2007, 64, 298-308.	2.5	27
47	Optimized biophysical model for Icelandic cod (<i>Gadus morhua</i>) larvae. Fisheries Oceanography, 2007, 16, 448-458.	1.7	5
48	Formulation and application of an efficient optimized biophysical model. Marine Ecology - Progress Series, 2007, 347, 275-284.	1.9	5
49	Discrimination between Icelandic cod (Gadus morhua L.) populations from adjacent spawning areas based on otolith growth and shape. Fisheries Research, 2006, 80, 182-189.	1.7	59
50	The genetic structure of Atlantic cod (Gadus morhua) around Iceland: insight from microsatellites, the Pan I locus, and tagging experiments. Canadian Journal of Fisheries and Aquatic Sciences, 2006, 63, 2660-2674.	1.4	116
51	Stock structure of Icelandic cod Gadus morhua L. based on otolith chemistry. Journal of Fish Biology, 2006, 69, 136-150.	1.6	35
52	Growth, maturity and fecundity of wolffish Anarhichas lupus L. in Icelandic waters. Journal of Fish Biology, 2006, 68, 1158-1176.	1.6	16
53	Effects of population size/age structure, condition and temporal dynamics of spawning on reproductive output in Atlantic cod (Gadus morhua). Ecological Modelling, 2006, 191, 383-415.	2.5	100
54	Otolith shape and temporal stability of spawning groups of Icelandic cod (Gadus morhua L.). ICES Journal of Marine Science, 2006, 63, 1501-1512.	2.5	58

#	Article	IF	CITATIONS
55	Temperature tolerance of Iceland scallop, Chlamys islandica (O.F. Muller) under controlled experimental conditions. Aquaculture Research, 2004, 35, 1405-1414.	1.8	12
56	Spatial partitioning of relative fishing mortality and spawning stock biomass of Icelandic cod. Fisheries Research, 2003, 59, 343-362.	1.7	24
57	Using Environmental and Biological Indices as Proxies for Egg and Larval Production of Marine Fish. Journal of Northwest Atlantic Fishery Science, 2003, 33, 115-159.	1.4	94
58	Developing Alternative Indices of Reproductive Potential for Use in Fisheries Management: Case Studies for Stocks Spanning an Information Gradient. Journal of Northwest Atlantic Fishery Science, 2003, 33, 161-190.	1.4	117
59	Genetic heterogeneity and growth properties of different genotypes of Atlantic cod (Gadus morhua) Tj $$ ETQq $$ 1 $$ 1 $$	0.784314 1.7	rgBT /Overloo
60	Environmental and stock effects on spatial distribution and abundance of mature cod Gadus morhua. Marine Ecology - Progress Series, 2002, 229, 245-262.	1.9	49
61	Environmental and stock effects on spawning origins and recruitment of cod Gadus morhua. Marine Ecology - Progress Series, 2002, 229, 263-277.	1.9	40
62	Essential relationships incorporating the influence of age, size and condition on variables required for estimation of reproductive potential in Atlantic cod Gadus morhua. Marine Ecology - Progress Series, 2002, 235, 235-256.	1.9	257
63	Spatial variation in hatch date distributions and origin of pelagic juvenile cod in Icelandic waters. ICES Journal of Marine Science, 2000, 57, 1182-1195.	2.5	46
64	Spatial variation in abundance, size composition and viable egg production of spawning cod (Gadus) Tj ETQq0 0	0 rgBT /O	verlock 10 Tf
65	Spawning origins of pelagic juvenile cod Gadus morhua inferred from spatially explicit age distributions: potential influences on year-class strength and recruitment. Marine Ecology - Progress Series, 2000, 202, 193-217.	1.9	46
66	Potential effects of maternal factors on spawning stock–recruitment relationships under varying fishing pressure. Canadian Journal of Fisheries and Aquatic Sciences, 1999, 56, 1882-1890.	1.4	82
67	Potential effects of maternal factors on spawning stock-recruitment relationships under varying fishing pressure. Canadian Journal of Fisheries and Aquatic Sciences, 1999, 56, 1882-1890.	1.4	70
68	Improving the stock-recruitment relationship in Icelandic cod (Gadus morhua) by including age diversity of spawners. Canadian Journal of Fisheries and Aquatic Sciences, 1998, 55, 1372-1377.	1.4	188
69	Maternal influence on the size and viability of Iceland codGadus morhuaeggs and larvae. Journal of Fish Biology, 1998, 52, 1241-1258.	1.6	185
70	Influence of egg size on embryos and larvae of Fundulus heteroclitus (L.). Journal of Fish Biology, 1992, 41, 883-896.	1.6	35
71	Retrospective assessment of premenstrual changes in Icelandic women. Health Care for Women International, 1991, 12, 303-315.	1.1	8