Young-Hwan Jo

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/928186/young-hwan-jo-publications-by-year.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

2,602 24 37 39 h-index g-index citations papers 4.88 8.3 2,909 39 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
37	Optogenetic stimulation of the liver-projecting melanocortinergic pathway promotes hepatic glucose production. <i>Nature Communications</i> , 2020 , 11, 6295	17.4	7
36	Hydrocarboxylic acid receptor 1 in BAT regulates glucose uptake in mice fed a high-fat diet. <i>PLoS ONE</i> , 2020 , 15, e0228320	3.7	1
35	Activation of the ARC-MeA Projection Reduces Food Intake. Frontiers in Neural Circuits, 2020, 14, 59578	33 3.5	9
34	A gut-brain axis regulating glucose metabolism mediated by bile acids and competitive fibroblast growth factor actions at the hypothalamus. <i>Molecular Metabolism</i> , 2018 , 8, 37-50	8.8	34
33	Intracellular glycolysis in brown adipose tissue is essential for optogenetically induced nonshivering thermogenesis in mice. <i>Scientific Reports</i> , 2018 , 8, 6672	4.9	34
32	Activation of temperature-sensitive TRPV1-like receptors in ARC POMC neurons reduces food intake. <i>PLoS Biology</i> , 2018 , 16, e2004399	9.7	35
31	Cholinergic neurons in the dorsomedial hypothalamus regulate food intake. <i>Molecular Metabolism</i> , 2017 , 6, 306-312	8.8	49
30	Single-Cell Gene Expression Analysis of Cholinergic Neurons in the Arcuate Nucleus of the Hypothalamus. <i>PLoS ONE</i> , 2016 , 11, e0162839	3.7	18
29	Cholinergic neurons in the dorsomedial hypothalamus regulate mouse brown adipose tissue metabolism. <i>Molecular Metabolism</i> , 2015 , 4, 483-92	8.8	36
28	Interplay between glucose and leptin signalling determines the strength of GABAergic synapses at POMC neurons. <i>Nature Communications</i> , 2015 , 6, 6618	17.4	24
27	Apelin-13 enhances arcuate POMC neuron activity via inhibiting M-current. <i>PLoS ONE</i> , 2015 , 10, e01194	15 <i>3</i> 7.7	12
26	Central action of FGF19 reduces hypothalamic AGRP/NPY neuron activity and improves glucose metabolism. <i>Molecular Metabolism</i> , 2014 , 3, 19-28	8.8	92
25	Why leptin keeps you warm. <i>Molecular Metabolism</i> , 2014 , 3, 779-80	8.8	4
24	The brain-liver connection between BDNF and glucose control. <i>Diabetes</i> , 2013 , 62, 1367-8	0.9	8
23	Clusterin and LRP2 are critical components of the hypothalamic feeding regulatory pathway. <i>Nature Communications</i> , 2013 , 4, 1862	17.4	43
22	pRb is an obesity suppressor in hypothalamus and high-fat diet inhibits pRb in this location. <i>EMBO Journal</i> , 2013 , 32, 844-57	13	16
21	Overnight fasting regulates inhibitory tone to cholinergic neurons of the dorsomedial nucleus of the hypothalamus. <i>PLoS ONE</i> , 2013 , 8, e60828	3.7	13

20	Effects of leptin and melanocortin signaling interactions on pubertal development and reproduction. <i>Endocrinology</i> , 2012 , 153, 2408-19	4.8	77
19	TXNIP in Agrp neurons regulates adiposity, energy expenditure, and central leptin sensitivity. <i>Journal of Neuroscience</i> , 2012 , 32, 9870-7	6.6	33
18	Endogenous BDNF regulates inhibitory synaptic transmission in the ventromedial nucleus of the hypothalamus. <i>Journal of Neurophysiology</i> , 2012 , 107, 42-9	3.2	14
17	Leptin action via neurotensin neurons controls orexin, the mesolimbic dopamine system and energy balance. <i>Cell Metabolism</i> , 2011 , 14, 313-23	24.6	241
16	Interplay between ionotropic receptors modulates inhibitory synaptic strength. <i>Communicative and Integrative Biology</i> , 2011 , 4, 706-9	1.7	4
15	Cross-talk between P2X4 and gamma-aminobutyric acid, type A receptors determines synaptic efficacy at a central synapse. <i>Journal of Biological Chemistry</i> , 2011 , 286, 19993-20004	5.4	44
14	Mediobasal hypothalamic leucine sensing regulates food intake through activation of a hypothalamus-brainstem circuit. <i>Journal of Neuroscience</i> , 2009 , 29, 8302-11	6.6	170
13	Direct innervation of GnRH neurons by metabolic- and sexual odorant-sensing leptin receptor neurons in the hypothalamic ventral premammillary nucleus. <i>Journal of Neuroscience</i> , 2009 , 29, 3138-47	, 6.6	124
12	Oleic acid directly regulates POMC neuron excitability in the hypothalamus. <i>Journal of Neurophysiology</i> , 2009 , 101, 2305-16	3.2	79
11	Transcription factors in the development of medial hypothalamic structures. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2009 , 297, E563-7	6	17
11		24.6	17 315
	Physiology - Endocrinology and Metabolism, 2009, 297, E563-7 Leptin acts via leptin receptor-expressing lateral hypothalamic neurons to modulate the		
10	Physiology - Endocrinology and Metabolism, 2009, 297, E563-7 Leptin acts via leptin receptor-expressing lateral hypothalamic neurons to modulate the mesolimbic dopamine system and suppress feeding. Cell Metabolism, 2009, 10, 89-98 Steroidogenic factor 1 regulates expression of the cannabinoid receptor 1 in the ventromedial		315
10	Physiology - Endocrinology and Metabolism, 2009, 297, E563-7 Leptin acts via leptin receptor-expressing lateral hypothalamic neurons to modulate the mesolimbic dopamine system and suppress feeding. Cell Metabolism, 2009, 10, 89-98 Steroidogenic factor 1 regulates expression of the cannabinoid receptor 1 in the ventromedial hypothalamic nucleus. Molecular Endocrinology, 2008, 22, 1950-61 Integration of endocannabinoid and leptin signaling in an appetite-related neural circuit. Neuron,	24.6	315
10 9 8	Physiology - Endocrinology and Metabolism, 2009, 297, E563-7 Leptin acts via leptin receptor-expressing lateral hypothalamic neurons to modulate the mesolimbic dopamine system and suppress feeding. Cell Metabolism, 2009, 10, 89-98 Steroidogenic factor 1 regulates expression of the cannabinoid receptor 1 in the ventromedial hypothalamic nucleus. Molecular Endocrinology, 2008, 22, 1950-61 Integration of endocannabinoid and leptin signaling in an appetite-related neural circuit. Neuron, 2005, 48, 1055-66 Cholinergic modulation of appetite-related synapses in mouse lateral hypothalamic slice. Journal of	24.6	315 28 190
10 9 8	Physiology - Endocrinology and Metabolism, 2009, 297, E563-7 Leptin acts via leptin receptor-expressing lateral hypothalamic neurons to modulate the mesolimbic dopamine system and suppress feeding. Cell Metabolism, 2009, 10, 89-98 Steroidogenic factor 1 regulates expression of the cannabinoid receptor 1 in the ventromedial hypothalamic nucleus. Molecular Endocrinology, 2008, 22, 1950-61 Integration of endocannabinoid and leptin signaling in an appetite-related neural circuit. Neuron, 2005, 48, 1055-66 Cholinergic modulation of appetite-related synapses in mouse lateral hypothalamic slice. Journal of Neuroscience, 2005, 25, 11133-44 Cholinergic modulation of purinergic and GABAergic co-transmission at in vitro hypothalamic	24.6 13.9 6.6	315 28 190 45
10 9 8 7 6	Physiology - Endocrinology and Metabolism, 2009, 297, E563-7 Leptin acts via leptin receptor-expressing lateral hypothalamic neurons to modulate the mesolimbic dopamine system and suppress feeding. Cell Metabolism, 2009, 10, 89-98 Steroidogenic factor 1 regulates expression of the cannabinoid receptor 1 in the ventromedial hypothalamic nucleus. Molecular Endocrinology, 2008, 22, 1950-61 Integration of endocannabinoid and leptin signaling in an appetite-related neural circuit. Neuron, 2005, 48, 1055-66 Cholinergic modulation of appetite-related synapses in mouse lateral hypothalamic slice. Journal of Neuroscience, 2005, 25, 11133-44 Cholinergic modulation of purinergic and GABAergic co-transmission at in vitro hypothalamic synapses. Journal of Neurophysiology, 2002, 88, 2501-8	24.6 13.9 6.6	315 28 190 45

Oxytocin modulates glutamatergic synaptic transmission between cultured neonatal spinal cord dorsal horn neurons. *Journal of Neuroscience*, **1998**, 18, 2377-86

6.6 81

Electrophysiological characterization of non-NMDA glutamate receptors on cultured intermediate lobe cells of the rat pituitary. *Neuroendocrinology*, **1996**, 64, 162-8

5.6 4