

Marcin Czepiel

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9289248/publications.pdf>

Version: 2024-02-01

19
papers

570
citations

840776

11
h-index

940533

16
g-index

19
all docs

19
docs citations

19
times ranked

971
citing authors

#	ARTICLE	IF	CITATIONS
1	Angiotensin II receptor 1 controls profibrotic Wnt/ β -catenin signalling in experimental autoimmune myocarditis. <i>Cardiovascular Research</i> , 2022, 118, 573-584.	3.8	13
2	Human and mouse PD-L1: similar molecular structure, but different druggability profiles. <i>IScience</i> , 2021, 24, 101960.	4.1	45
3	Expression of VEGFA-mRNA in classical and MSX2-mRNA in non-classical monocytes in patients with spondyloarthritis is associated with peripheral arthritis. <i>Scientific Reports</i> , 2021, 11, 9693.	3.3	0
4	Downregulation of Dkk1 in Platelets of Patients With Axial Spondyloarthritis. <i>Arthritis and Rheumatology</i> , 2021, 73, 1831-1834.	5.6	5
5	WNT/ β -Catenin Signaling Promotes TGF- β -Mediated Activation of Human Cardiac Fibroblasts by Enhancing IL-11 Production. <i>International Journal of Molecular Sciences</i> , 2021, 22, 10072.	4.1	31
6	Activated myofibroblasts promote cardiac hypertrophy and systolic dysfunction independently of cardiac fibrosis in experimental autoimmune myocarditis. <i>European Heart Journal</i> , 2021, 42, .	2.2	0
7	Abstract 121: Interleukin 11 Mediates Wnt/ β -catenin-dependent Fibrotic Response Of Human Cardiac Fibroblasts. <i>Circulation Research</i> , 2021, 129, .	4.5	0
8	Heart non-specific effector CD4+ T cells protect from postinflammatory fibrosis and cardiac dysfunction in experimental autoimmune myocarditis. <i>Basic Research in Cardiology</i> , 2020, 115, 6.	5.9	17
9	Haploinsufficient <i>Rock1</i> ^{+/-} and <i>Rock2</i> ^{+/-} Mice Are Not Protected from Cardiac Inflammation and Postinflammatory Fibrosis in Experimental Autoimmune Myocarditis. <i>Cells</i> , 2020, 9, 700.	4.1	5
10	Identification and Isolation of Cardiac Fibroblasts From the Adult Mouse Heart Using Two-Color Flow Cytometry. <i>Frontiers in Cardiovascular Medicine</i> , 2019, 6, 105.	2.4	23
11	WNT3a and WNT5a Transported by Exosomes Activate WNT Signaling Pathways in Human Cardiac Fibroblasts. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1436.	4.1	54
12	Sera of patients with axial spondyloarthritis (axSpA) enhance osteoclastogenic potential of monocytes isolated from healthy individuals. <i>BMC Musculoskeletal Disorders</i> , 2018, 19, 434.	1.9	6
13	Characterization and comparison of osteoblasts derived from mouse embryonic stem cells and induced pluripotent stem cells. <i>Journal of Bone and Mineral Metabolism</i> , 2017, 35, 21-30.	2.7	17
14	Survival and Functionality of Human Induced Pluripotent Stem Cell-Derived Oligodendrocytes in a Nonhuman Primate Model for Multiple Sclerosis. <i>Stem Cells Translational Medicine</i> , 2016, 5, 1550-1561.	3.3	57
15	Human oligodendrocytes in remyelination research. <i>Glia</i> , 2015, 63, 513-530.	4.9	55
16	Overexpression of Polysialylated Neural Cell Adhesion Molecule Improves the Migration Capacity of Induced Pluripotent Stem Cell-Derived Oligodendrocyte Precursors. <i>Stem Cells Translational Medicine</i> , 2014, 3, 1100-1109.	3.3	19
17	Generation of Induced Pluripotent Stem Cells from Hair Follicle Bulge Neural Crest Stem Cells. <i>Cellular Reprogramming</i> , 2014, 16, 307-313.	0.9	3
18	Differentiation of induced pluripotent stem cells into functional oligodendrocytes. <i>Glia</i> , 2011, 59, 882-892.	4.9	118

#	ARTICLE	IF	CITATIONS
19	Continuous hypoxic culturing maintains activation of Notch and allows long-term propagation of human embryonic stem cells without spontaneous differentiation. <i>Cell Proliferation</i> , 2009, 42, 63-74.	5.3	102