

Katia Boniface

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9301685/publications.pdf>

Version: 2024-02-01

65
papers

6,983
citations

230014

27
h-index

162838

57
g-index

66
all docs

66
docs citations

66
times ranked

10484
citing authors

#	ARTICLE	IF	CITATIONS
1	Vitiligo Skin T Cells Are Prone to Produce Type 1 and Type 2 Cytokines to Induce Melanocyte Dysfunction and Epidermal Inflammatory Response Through Jak Signaling. <i>Journal of Investigative Dermatology</i> , 2022, 142, 1194-1205.e7.	0.3	23
2	Assessment of Vitiligo Area Scoring Index (VASI), Facial-VASI and Vitiligo Extent Score using standardized photography of patients with vitiligo. <i>British Journal of Dermatology</i> , 2022, 187, 422-424.	1.4	6
3	Cytokine-Mediated Crosstalk Between Keratinocytes and T Cells in Atopic Dermatitis. <i>Frontiers in Immunology</i> , 2022, 13, 801579.	2.2	23
4	Alopecia areata: Recent advances and emerging therapies. <i>Annales De Dermatologie Et De Venereologie</i> , 2022, 149, 222-227.	0.5	5
5	An update on Vitiligo pathogenesis. <i>Pigment Cell and Melanoma Research</i> , 2021, 34, 236-243.	1.5	61
6	Targeting Innate Immunity to Combat Cutaneous Stress: The Vitiligo Perspective. <i>Frontiers in Immunology</i> , 2021, 12, 613056.	2.2	19
7	Demographic and clinical characteristics of patients with both psoriasis and vitiligo in a cohort of vitiligo patients: a cross-sectional study. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2021, 35, e676-e679.	1.3	9
8	Editorial: Immunology of Vitiligo. <i>Frontiers in Immunology</i> , 2021, 12, 711080.	2.2	4
9	Complete response in a patient with advanced melanoma following anti-PD-1 therapy is associated with a high frequency of melanoma-infiltrating CXCR3 ⁺ resident memory CD8 ⁺ T cells and multiple chemokine pathways. <i>British Journal of Dermatology</i> , 2021, 185, 663-666.	1.4	1
10	Analysis of tumor response and clinical factors associated with vitiligo in patients receiving anti-programmed cell death-1 therapies for melanoma: A cross-sectional study. <i>JAAD International</i> , 2021, 5, 112-120.	1.1	10
11	NKG2D Defines a Subset of Skin Effector Memory CD8 T Cells with Proinflammatory Functions in Vitiligo. <i>Journal of Investigative Dermatology</i> , 2020, 140, 1143-1153.e5.	0.3	32
12	Vitiligo, From Physiopathology to Emerging Treatments: A Review. <i>Dermatology and Therapy</i> , 2020, 10, 1185-1198.	1.4	29
13	Characteristics of postinflammatory hyper- and hypopigmentation in patients with psoriasis: A survey study. <i>Journal of the American Academy of Dermatology</i> , 2020, 83, 1188-1191.	0.6	8
14	Phenotype and function of circulating memory T cells in human vitiligo*. <i>British Journal of Dermatology</i> , 2020, 183, 899-908.	1.4	14
15	Type-1 cytokines regulate matrix metalloprotease-9 production and E-cadherin disruption to promote melanocyte loss in vitiligo. <i>JCI Insight</i> , 2020, 5, .	2.3	31
16	An unusual presentation of Sutton's phenomenon presenting as halo cherry angioma. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2019, 33, e464-e465.	1.3	0
17	Vitiligo-Like Lesions in Patients with Metastatic Melanoma Receiving Immunotherapies. , 2019, , 163-167.		0
18	Cytokines, Growth Factors, and POMC Peptides. , 2019, , 303-312.		0

#	ARTICLE	IF	CITATIONS
19	Imbalance of peripheral follicular helper T lymphocyte subsets in active vitiligo. <i>Pigment Cell and Melanoma Research</i> , 2019, 32, 588-592.	1.5	4
20	Vitiligo as a skin memory disease: The need for early intervention with immunomodulating agents and a maintenance therapy to target resident memory T cells. <i>Experimental Dermatology</i> , 2019, 28, 656-661.	1.4	26
21	Vitiligo-like lesions occurring in patients receiving anti-programmed cell death-1 therapies. <i>Giornale Italiano Di Dermatologia E Venereologia</i> , 2019, 154, 435-443.	0.8	5
22	Cell delivery using microneedle devices: a new approach to treat depigmenting disorders. <i>British Journal of Dermatology</i> , 2018, 178, 588-589.	1.4	2
23	Vitiligo-like lesions in patients receiving anti-programmed cell death-1 therapies are distinct from spontaneously occurring active vitiligo. <i>Journal of the American Academy of Dermatology</i> , 2018, 78, e17-e18.	0.6	13
24	Elevated total serum IgE in vitiligo might be protective for other autoimmune diseases. <i>British Journal of Dermatology</i> , 2018, 179, 987-988.	1.4	2
25	Dual efficacy of dupilumab in a patient with concomitant atopic dermatitis and alopecia areata. <i>British Journal of Dermatology</i> , 2018, 179, 534-536.	1.4	40
26	Vitiligo: Focus on Clinical Aspects, Immunopathogenesis, and Therapy. <i>Clinical Reviews in Allergy and Immunology</i> , 2018, 54, 52-67.	2.9	155
27	Vitiligo Skin Is Imprinted with Resident Memory CD8 T Cells Expressing CXCR3. <i>Journal of Investigative Dermatology</i> , 2018, 138, 355-364.	0.3	168
28	Inflammatory skin eruptions induced by anti-tumour necrosis factor- α therapy differ undeniably from psoriasis or eczema. <i>British Journal of Dermatology</i> , 2018, 178, 1007-1008.	1.4	0
29	Inflammasome Activation Characterizes Lesional Skin of Folliculitis Decalvans. <i>Acta Dermato-Venereologica</i> , 2018, 98, 570-575.	0.6	11
30	In vitro models of vitiligo. , 2018, , 129-149.		0
31	Vitiligo therapy: restoring immune privilege?. <i>Experimental Dermatology</i> , 2017, 26, 635-636.	1.4	3
32	Vitiligo-like lesions occurring in patients receiving anti-programmed cell death-1 therapies are clinically and biologically distinct from vitiligo. <i>Journal of the American Academy of Dermatology</i> , 2017, 76, 863-870.	0.6	128
33	Meeting report: Vitiligo Global Issues Consensus Conference Workshop "Outcome measurement instruments" and Vitiligo International Symposium, Rome, Nov 30-Dec 3rd. <i>Pigment Cell and Melanoma Research</i> , 2017, 30, 436-443.	1.5	14
34	MicroRNA-211 Regulates Oxidative Phosphorylation and Energy Metabolism in Human Vitiligo. <i>Journal of Investigative Dermatology</i> , 2017, 137, 1965-1974.	0.3	55
35	Heat shock protein 70 potentiates interferon alpha production by plasmacytoid dendritic cells: relevance for cutaneous lupus and vitiligo pathogenesis. <i>British Journal of Dermatology</i> , 2017, 177, 1367-1375.	1.4	75
36	A Score with a VESTed Interest in Vitiligo. <i>Journal of Investigative Dermatology</i> , 2016, 136, 902-904.	0.3	0

#	ARTICLE	IF	CITATIONS
37	New insights into immune mechanisms of vitiligo. <i>Giornale Italiano Di Dermatologia E Venereologia</i> , 2016, 151, 44-54.	0.8	13
38	Comment: the mystery of melanocyte demise in vitiligo. <i>Experimental Dermatology</i> , 2015, 24, 260-261.	1.4	3
39	A Th2 Cytokine Interleukin-31 Signature in a Case of Sporadic Lichen Amyloidosis. <i>Acta Dermato-Venereologica</i> , 2015, 95, 223-224.	0.6	9
40	Type I interferon signature in the initiation of the immune response in vitiligo. <i>Pigment Cell and Melanoma Research</i> , 2014, 27, 398-407.	1.5	118
41	Expression of interleukin-1 alpha in amicrobial pustulosis of the skin folds with complete response to anakinra. <i>Journal of the American Academy of Dermatology</i> , 2014, 71, e53-e56.	0.6	25
42	Accelerating bleaching in vitiligo: balancing benefits versus risks. <i>Experimental Dermatology</i> , 2014, 23, 879-880.	1.4	4
43	Malvidin-3-O- β glucoside, major grape anthocyanin, inhibits human macrophage-derived inflammatory mediators and decreases clinical scores in arthritic rats. <i>Biochemical Pharmacology</i> , 2013, 86, 1461-1467.	2.0	68
44	Development and validation of the VSCOR for scoring Koebner's phenomenon in vitiligo/non-segmental vitiligo. <i>Pigment Cell and Melanoma Research</i> , 2013, 26, 402-407.	1.5	26
45	Autoimmune Memory T Helper 17 Cell Function and Expansion Are Dependent on Interleukin-23. <i>Cell Reports</i> , 2013, 3, 1378-1388.	2.9	72
46	IL-17 in the Rheumatologist's Line of Sight. <i>BioMed Research International</i> , 2013, 2013, 1-18.	0.9	24
47	Role of Th17 cells in the pathogenesis of rheumatoid arthritis. <i>World Journal of Rheumatology</i> , 2013, 3, 25.	0.5	5
48	Biomarkers of Therapeutic Response in the IL-23 Pathway in Inflammatory Bowel Disease. <i>Clinical and Translational Gastroenterology</i> , 2012, 3, e10.	1.3	47
49	Pre- vs. post-pubertal onset of vitiligo: multivariate analysis indicates atopic diathesis association in pre-pubertal onset vitiligo. <i>British Journal of Dermatology</i> , 2012, 167, 490-495.	1.4	55
50	T helper type 1 and 17 cells determine efficacy of interferon- β in multiple sclerosis and experimental encephalomyelitis. <i>Nature Medicine</i> , 2010, 16, 406-412.	15.2	509
51	Skin Inflammation Induced by the Synergistic Action of IL-17A, IL-22, Oncostatin M, IL-1 β , and TNF- α Recapitulates Some Features of Psoriasis. <i>Journal of Immunology</i> , 2010, 184, 5263-5270.	0.4	274
52	Human Th17 Cells Comprise Heterogeneous Subsets Including IFN- γ -Producing Cells with Distinct Properties from the Th1 Lineage. <i>Journal of Immunology</i> , 2010, 185, 679-687.	0.4	163
53	Circulating and gut-resident human Th17 cells express CD161 and promote intestinal inflammation. <i>Journal of Experimental Medicine</i> , 2009, 206, 525-534.	4.2	430
54	Prostaglandin E2 regulates Th17 cell differentiation and function through cyclic AMP and EP2/EP4 receptor signaling. <i>Journal of Experimental Medicine</i> , 2009, 206, 535-548.	4.2	426

#	ARTICLE	IF	CITATIONS
55	IL-27 Blocks ROR γ c Expression to Inhibit Lineage Commitment of Th17 Cells. <i>Journal of Immunology</i> , 2009, 182, 5748-5756.	0.4	302
56	Development and function of TH17 cells in health and disease. <i>Journal of Allergy and Clinical Immunology</i> , 2009, 123, 1004-1011.	1.5	223
57	Prostaglandin E2 regulates Th17 cell differentiation and function through cyclic AMP and EP2/EP4 receptor signaling. <i>Journal of Cell Biology</i> , 2009, 184, i16-i16.	2.3	1
58	From interleukin-23 to T γ H17 cells: human T γ H17 cell differentiation revisited. <i>Immunological Reviews</i> , 2008, 226, 132-146.	2.8	194
59	Oncostatin M Secreted by Skin Infiltrating T Lymphocytes Is a Potent Keratinocyte Activator Involved in Skin Inflammation. <i>Journal of Immunology</i> , 2007, 178, 4615-4622.	0.4	160
60	104 A role for Th17-Derived IL-22 in Psoriatic Skin Inflammation. <i>Cytokine</i> , 2007, 39, 28-29.	1.4	0
61	Development, cytokine profile and function of human interleukin 17 γ -producing helper T cells. <i>Nature Immunology</i> , 2007, 8, 950-957.	7.0	1,795
62	A role for T cell-derived interleukin 22 in psoriatic skin inflammation. <i>Clinical and Experimental Immunology</i> , 2007, 150, 407-415.	1.1	253
63	IL-22 Inhibits Epidermal Differentiation and Induces Proinflammatory Gene Expression and Migration of Human Keratinocytes. <i>Journal of Immunology</i> , 2005, 174, 3695-3702.	0.4	726
64	Keratinocytes as targets for interleukin-10-related cytokines: a putative role in the pathogenesis of psoriasis. <i>European Cytokine Network</i> , 2005, 16, 309-19.	1.1	42
65	IL-22, in contrast to IL-10, does not induce Ig production, due to absence of a functional IL-22 receptor on activated human B cells. <i>International Immunology</i> , 2002, 14, 1351-1356.	1.8	40