L Bnger

List of Publications by Citations

Source: https://exaly.com/author-pdf/9315834/l-bunger-publications-by-citations.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

132 2,658 27 44 g-index

140 3,014 3.3 4.55 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
132	Lack of myostatin results in excessive muscle growth but impaired force generation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 1835-40	11.5	293
131	Prediction of sensory characteristics of lamb meat samples by near infrared reflectance spectroscopy. <i>Meat Science</i> , 2007 , 76, 509-16	6.4	110
130	Resting metabolic rate and morphology in mice (Mus musculus) selected for high and low food intake. <i>Journal of Experimental Biology</i> , 2001 , 204, 777-784	3	104
129	Non-invasive methods for the determination of body and carcass composition in livestock: dual-energy X-ray absorptiometry, computed tomography, magnetic resonance imaging and ultrasound: invited review. <i>Animal</i> , 2015 , 9, 1250-64	3.1	87
128	Resting metabolic rate and morphology in mice (Mus musculus) selected for high and low food intake. <i>Journal of Experimental Biology</i> , 2001 , 204, 777-84	3	80
127	Mapping of obesity QTLs in a cross between mouse lines divergently selected on fat content. <i>Mammalian Genome</i> , 2000 , 11, 2-7	3.2	68
126	Parallel selection mapping using artificially selected mice reveals body weight control loci. <i>Current Biology</i> , 2012 , 22, 794-800	6.3	64
125	A polygenic model of the metabolic syndrome with reduced circulating and intra-adipose glucocorticoid action. <i>Diabetes</i> , 2005 , 54, 3371-8	0.9	57
124	Inbred lines of mice derived from long-term growth selected lines: unique resources for mapping growth genes. <i>Mammalian Genome</i> , 2001 , 12, 678-86	3.2	53
123	Effects of low protein diets on pigs with a lean genotype. 1. Carcass composition measured by dissection and muscle fatty acid composition. <i>Meat Science</i> , 2013 , 95, 123-8	6.4	46
122	A genetic investigation of various growth models to describe growth of lambs of two contrasting breeds. <i>Journal of Animal Science</i> , 2006 , 84, 2642-54	0.7	43
121	A stratified transcriptomics analysis of polygenic fat and lean mouse adipose tissues identifies novel candidate obesity genes. <i>PLoS ONE</i> , 2011 , 6, e23944	3.7	42
120	Effects of the compact mutant myostatin allele Mstn (Cmpt-dl1Abc) introgressed into a high growth mouse line on skeletal muscle cellularity. <i>Journal of Muscle Research and Cell Motility</i> , 2005 , 26, 103-12	3.5	39
119	A review of the development and use of video image analysis (VIA) for beef carcass evaluation as an alternative to the current EUROP system and other subjective systems. <i>Meat Science</i> , 2012 , 92, 307-7	18.4	37
118	The use of various live animal measurements to predict carcass and meat quality in two divergent lamb breeds. <i>Meat Science</i> , 2008 , 80, 1138-49	6.4	36
117	Genetic complexity of an obesity QTL (Fob3) revealed by detailed genetic mapping. <i>Mammalian Genome</i> , 2004 , 15, 472-81	3.2	35
116	Evaluation of Video Image Analysis (VIA) technology to predict meat yield of sheep carcasses on-line under UK abattoir conditions. <i>Meat Science</i> , 2009 , 82, 94-100	6.4	34

(2012-2005)

1	15	Microarray gene expression analysis of the Fob3b obesity QTL identifies positional candidate gene Sqle and perturbed cholesterol and glycolysis pathways. <i>Physiological Genomics</i> , 2005 , 20, 224-32	3.6	33	
1	14	Inbred lines of mice derived from long-term divergent selection on fat content and body weight. Mammalian Genome, 1999 , 10, 645-8	3.2	33	
1	13	Factors affecting dystocia and offspring vigour in different sheep genotypes. <i>Preventive Veterinary Medicine</i> , 2012 , 103, 257-64	3.1	29	
1	12	Comparison of the RichardsonLucy method and a classical approach for spectrometer bandpass correction. <i>Metrologia</i> , 2013 , 50, 107-118	2.1	29	
1	11	Meta-analysis of effects of gender in combination with carcass weight and breed on pork quality. Journal of Animal Science, 2013, 91, 1480-92	0.7	29	
1	10	Breeding for resistance to mastitis in United Kingdom sheep, a review and economic appraisal. Veterinary Record, 2008, 162, 369-76	0.9	29	
1	09	Thermoregulatory responses of two mouse Mus musculus strains selectively bred for high and low food intake. <i>Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology,</i> 2001 , 171, 661-8	2.2	29	
1	08	Analysis of response to 20 generations of selection for body composition in mice: fit to infinitesimal model assumptions. <i>Genetics Selection Evolution</i> , 2000 , 32, 3-21	4.9	29	
1	07	Relationships between quantitative and reproductive fitness traits in animals. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2005 , 360, 1489-502	5.8	28	
1	06	Muscularity and eating quality of lambs: Effects of breed, sex and selection of sires using muscularity measurements by computed tomography. <i>Meat Science</i> , 2008 , 79, 105-12	6.4	27	
1	05	Accuracy of in vivo muscularity indices measured by computed tomography and their association with carcass quality in lambs. <i>Meat Science</i> , 2007 , 75, 533-42	6.4	26	
1	04	Breeding for resistance to footrotthe use of hoof lesion scoring to quantify footrot in sheep. Veterinary Research Communications, 2008, 32, 583-9	2.9	25	
1	03	Marker-assisted introgression of the Compact mutant myostatin allele MstnCmpt-dl1Abc into a mouse line with extreme growth effects on body composition and muscularity. <i>Genetical Research</i> , 2004 , 84, 161-73	1.1	25	
1	02	Prediction of intramuscular fat content using CT scanning of packaged lamb cuts and relationships with meat eating quality. <i>Meat Science</i> , 2017 , 123, 112-119	6.4	24	
1	01	Genetic and phenotypic aspects of foot lesion scores in sheep of different breeds and ages. <i>Animal</i> , 2008 , 2, 1289-96	3.1	23	
1	00	In vivo measurements of muscle volume by automatic image analysis of spiral computed tomography scans. <i>Animal Science</i> , 2006 , 82, 545-553		22	
9	9	Polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) assay for the mouse leptin receptor (Lepr(db)) mutation. <i>Laboratory Animals</i> , 1999 , 33, 380-4	2.6	22	
9	8	Genetic parameters for fitness and neonatal behavior traits in sheep. <i>Behavior Genetics</i> , 2012 , 42, 899-91	3 1 2	21	

97	Prediction of lamb carcass composition and meat quality using combinations of post-mortem measurements. <i>Meat Science</i> , 2009 , 81, 711-9	6.4	21
96	Divergent physical activity and novel alternative responses to high fat feeding in polygenic fat and lean mice. <i>Behavior Genetics</i> , 2008 , 38, 292-300	3.2	21
95	Long-term selection for protein amount over 70 generations in mice. <i>Genetical Research</i> , 1998 , 72, 93-	109.1	21
94	Effects of feed allowance and indispensable amino acid reduction on feed intake, growth performance and carcass characteristics of growing pigs. <i>PLoS ONE</i> , 2018 , 13, e0195645	3.7	20
93	Effects of low protein diets on pigs with a lean genotype 2. Compositional traits measured with computed tomography (CT). <i>Meat Science</i> , 2013 , 95, 129-36	6.4	20
92	Development and validation of on-farm behavioural scoring systems to assess birth assistance and lamb vigour. <i>Animal</i> , 2011 , 5, 776-83	3.1	20
91	Bayesian meta-analysis of the effect of fasting, transport and lairage times on four attributes of pork meat quality. <i>Meat Science</i> , 2012 , 90, 584-98	6.4	19
90	Effects of selection on food intake in the adult mouse. <i>Journal of Animal Breeding and Genetics</i> , 1997 , 114, 419-34	2.9	19
89	Testing selection indices for sustainable hill sheep production [lamb growth and carcass traits. <i>Animal Science</i> , 2006 , 82, 445-453		19
88	Meta-analysis of the effects of dietary vitamin E supplementation on £ocopherol concentration and lipid oxidation in pork. <i>Meat Science</i> , 2011 , 87, 305-14	6.4	18
87	Evaluating the effects of a single copy of a mutation in the myostatin gene (c.*1232G>A) on carcass traits in crossbred lambs. <i>Meat Science</i> , 2011 , 87, 412-8	6.4	18
86	Genetic-statistical analysis of growth in selected and unselected mouse lines. <i>Journal of Experimental Animal Science</i> , 2003 , 42, 218-232		18
85	Mice with low metabolic rates are not susceptible to weight gain when fed a high-fat diet. <i>Obesity</i> , 2005 , 13, 556-66		18
84	Prediction of intramuscular fat levels in Texel lamb loins using X-ray computed tomography scanning. <i>Meat Science</i> , 2014 , 98, 263-71	6.4	17
83	Changes in carcass traits during growth in lambs of two contrasting breeds, measured using computer tomography. <i>Livestock Science</i> , 2007 , 107, 37-52	1.7	17
82	Long-term divergent selection on fatness in mice indicates a regulation system independent of leptin production and reception. <i>FASEB Journal</i> , 2003 , 17, 85-7	0.9	17
81	Genetic relationship between longevity and objectively or subjectively assessed performance traits in sheep using linear censored models. <i>Journal of Animal Science</i> , 2009 , 87, 3482-9	0.7	16
80	Effects of leptin administration on long-term selected fat mice. <i>Genetical Research</i> , 1997 , 69, 215-25	1.1	16

(2012-2006)

79	In vivo prediction of internal fat weight in Scottish Blackface lambs, using computer tomography. Journal of Animal Breeding and Genetics, 2006 , 123, 105-13	2.9	16	
78	A paternally imprinted QTL for mature body mass on mouse chromosome 8. <i>Mammalian Genome</i> , 2005 , 16, 567-77	3.2	16	
77	Between- and within-breed variations of spine characteristics in sheep. <i>Journal of Animal Science</i> , 2013 , 91, 995-1004	0.7	15	
76	Associations of polymorphisms of the ovine prion protein gene with growth, carcass, and computerized tomography traits in Scottish Blackface lambs. <i>Journal of Animal Science</i> , 2007 , 85, 632-4	10 ^{0.7}	15	
75	Intrinsic properties of muscle satellite cells are changed in response to long-term selection of mice for different growth traits. <i>Cell and Tissue Research</i> , 2002 , 310, 339-48	4.2	15	
74	Genome-wide association study of footrot in Texel sheep. <i>Genetics Selection Evolution</i> , 2015 , 47, 35	4.9	14	
73	The prediction of carcass composition and tissue distribution in beef cattle using ultrasound scanning at the start and/or end of the finishing period. <i>Livestock Science</i> , 2010 , 131, 193-202	1.7	13	
7 ²	Prediction of lamb meat eating quality in two divergent breeds using various live animal and carcass measurements. <i>Meat Science</i> , 2009 , 83, 366-75	6.4	13	
71	Genetic growth potential interacts with nutrition on the ability of mice to cope with Heligmosomoides bakeri infection. <i>Parasitology</i> , 2009 , 136, 1043-55	2.7	13	
70	Characterisation of white line degeneration in sheep and evidence for genetic influences on its occurrence. <i>Veterinary Research Communications</i> , 2010 , 34, 481-9	2.9	13	
69	Growth selection in mice reveals conserved and redundant expression patterns of the insulin-like growth factor system. <i>General and Comparative Endocrinology</i> , 2004 , 136, 248-59	3	13	
68	Genetic and genomic analyses of musculoskeletal differences between BEH and BEL strains. <i>Physiological Genomics</i> , 2013 , 45, 940-7	3.6	12	
67	Effects of the Texel muscling quantitative trait locus on carcass traits in crossbred lambs. <i>Animal</i> , 2009 , 3, 189-99	3.1	12	
66	The effects of selection indices for sustainable hill sheep production on carcass composition and muscularity of lambs, measured using X-ray computed tomography. <i>Animal</i> , 2008 , 2, 27-35	3.1	12	
65	Inbred lines of mice derived from long-term growth selected lines: unique resources for mapping growth genes 2001 , 12, 678		12	
64	Analyses of muscle spindles in the soleus of six inbred mouse strains. <i>Journal of Anatomy</i> , 2013 , 223, 289-96	2.9	11	
63	Genotypic effects of the Texel Muscling QTL (TM-QTL) on meat quality in purebred Texel lambs. <i>Meat Science</i> , 2011 , 89, 125-32	6.4	11	
62	The effect of sex on some carcass and meat quality traits in Texel ewe and ram lambs. <i>Animal Production Science</i> , 2012 , 52, 601	1.4	11	

61	Oxidative costs of reproduction in mouse strains selected for different levels of food intake and which differ in reproductive performance. <i>Scientific Reports</i> , 2016 , 6, 36353	4.9	10
60	Evaluating the effects of the c.*1232G > A mutation and TM-QTL in TexelWelsh Mountain lambs using ultrasound and video image analyses. <i>Small Ruminant Research</i> , 2011 , 99, 99-109	1.7	10
59	Muscle fibre characteristics of two contrasting sheep breeds: Scottish Blackface and Texel. <i>Meat Science</i> , 2009 , 81, 372-81	6.4	10
58	The effects of a loin muscling quantitative trait locus (LoinMAXII) on carcass and VIA-based traits in crossbred lambs. <i>Animal</i> , 2010 , 4, 407-16	3.1	10
57	Leptin levels in lines of mice developed by long-term divergent selection on fat content. <i>Genetical Research</i> , 1999 , 73, 37-44	1.1	10
56	Role of growth hormone in the genetic change of mice divergently selected for body weight and fatness. <i>Genetical Research</i> , 1999 , 74, 351-60	1.1	10
55	Analysis of a long-term selection experiment with an exponential model. <i>Journal of Animal Breeding and Genetics</i> , 1994 , 111, 1-13	2.9	10
54	The relationship between video image analysis (VIA), visual classification, and saleable meat yield of sirloin and fillet cuts of beef carcasses differing in breed and gender. <i>Livestock Science</i> , 2013 , 158, 169-178	1.7	9
53	Genetic parameters for carcass dimensional measurements from Video Image Analysis and their association with conformation and fat class scores. <i>Livestock Science</i> , 2010 , 128, 92-100	1.7	9
52	The effect of the Texel muscling QTL (TM-QTL) on meat quality traits in crossbred lambs. <i>Meat Science</i> , 2010 , 85, 684-90	6.4	9
51	Effects of a quantitative trait locus for increased muscularity on carcass traits measured by subjective conformation and fat class scores and video image analysis in crossbred lambs. <i>Animal</i> , 2009 , 3, 1532-43	3.1	9
50	Relationships between lamb carcass quality traits measured by X-ray computed tomography and current UK hill sheep breeding goals. <i>Animal</i> , 2008 , 2, 36-43	3.1	9
49	Characterization of a major X-linked quantitative trait locus influencing body weight of mice. <i>Journal of Heredity</i> , 2001 , 92, 355-7	2.4	9
48	Traceability of a CCD-Camera System for High-Temperature Measurements. <i>International Journal of Thermophysics</i> , 2015 , 36, 1784-1802	2.1	8
47	Investigation into the presence of genotype by environment interactions (GE) in Scottish Blackface lamb traits. <i>Small Ruminant Research</i> , 2012 , 105, 46-52	1.7	8
46	Interactive effects of protein nutrition, genetic growth potential and Heligmosomoides bakeri infection pressure on resilience and resistance in mice. <i>Parasitology</i> , 2011 , 138, 1305-15	2.7	7
45	Quantitative trait Loci for regional adiposity in mouse lines divergently selected for food intake. <i>Obesity</i> , 2007 , 15, 2994-3004	8	7
44	Effects of thyroid hormone deficiency on mice selected for increased and decreased body weight and fatness. <i>Genetical Research</i> , 1998 , 72, 39-53	1.1	7

(2010-2020)

43	Predicting the shear value and intramuscular fat in meat from Nellore cattle using Vis-NIR spectroscopy. <i>Meat Science</i> , 2020 , 163, 108077	6.4	7
42	Myostatin dysfunction is associated with reduction in overload induced hypertrophy of soleus muscle in mice. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2016 , 26, 894-901	4.6	7
41	Absolute Radiation Thermometry in the NIR. International Journal of Thermophysics, 2017, 38, 1	2.1	6
40	Prediction of intramuscular fat content and shear force in Texel lamb loins using combinations of different X-ray computed tomography (CT) scanning techniques. <i>Meat Science</i> , 2018 , 140, 78-85	6.4	6
39	Genetic parameters for carcass composition and performance data in crossbred lambs measured by Video Image Analysis. <i>Meat Science</i> , 2009 , 81, 619-25	6.4	6
38	Effects of low protein diets on performance of pigs with a lean genotype between 40 and 115 kg liveweight. <i>Animal Production Science</i> , 2015 , 55, 461	1.4	5
37	Analysis of single nucleotide polymorphisms variation associated with important economic and computed tomography measured traits in Texel sheep. <i>Animal</i> , 2018 , 12, 915-922	3.1	5
36	Genetic improvement of hill sheep Impacts on profitability and greenhouse gas emissions. <i>Small Ruminant Research</i> , 2014 , 120, 27-34	1.7	5
35	Index selection in terminal sires improves lamb performance at finishing. <i>Journal of Animal Science</i> , 2013 , 91, 38-43	0.7	5
34	Use of X-Ray Computed Tomography (CT) in UK Sheep Production and Breeding 2011 ,		5
33	The effect of conditioning period on loin muscle tenderness in crossbred lambs with or without the Texel muscling QTL (TM-QTL). <i>Meat Science</i> , 2010 , 85, 715-20	6.4	5
32	Index selection in terminal sires improves early lamb growth. Journal of Animal Science, 2012, 90, 142-5	10.7	5
31	Neuropeptide Y gene expression in lines of mice subjected to long-term divergent selection on fat content. <i>Journal of Molecular Endocrinology</i> , 1999 , 23, 77-83	4.5	5
30	Limits to sustained energy intake. XXII. Reproductive performance of two selected mouse lines with different thermal conductance. <i>Journal of Experimental Biology</i> , 2014 , 217, 3718-32	3	4
29	Effect of the Texel muscling QTL (TM-QTL) on spine characteristics in purebred Texel lambs. <i>Small Ruminant Research</i> , 2014 , 117, 34-40	1.7	4
28	The effects of different farm environments on the performance of Texel sheep. <i>Animal</i> , 2015 , 9, 1624-3	43.1	4
27	Characterisation of terminal sire sheep farm systems, based on a range of environmental factors: a case study in the context of genotype by environment interactions using Charollais lambs. <i>Animal</i> , 2014 , 8, 867-76	3.1	4
26	Comparison of repeatability and multiple trait threshold models for litter size in sheep using observed and simulated data in Bayesian analyses. <i>Journal of Animal Breeding and Genetics</i> , 2010 , 127, 261-71	2.9	4

25	Baseline Muscle Mass Is a Poor Predictor of Functional Overload-Induced Gain in the Mouse Model. <i>Frontiers in Physiology</i> , 2016 , 7, 534	4.6	4
24	Prediction of intramuscular fat in lamb by visible and near-infrared spectroscopy in an abattoir environment. <i>Meat Science</i> , 2021 , 171, 108286	6.4	4
23	Myostatin dysfunction impairs force generation in extensor digitorum longus muscle and increases exercise-induced protein efflux from extensor digitorum longus and soleus muscles. <i>Applied Physiology, Nutrition and Metabolism</i> , 2015 , 40, 817-21	3	3
22	Interactive effects of selection for growth and protein supply on the consequences of gastrointestinal parasitism on growth performance in mice. <i>Proceedings of the British Society of Animal Science</i> , 2007 , 2007, 92-92		3
21	Fine mapping of mouse QTLs for fatness using SNP data. <i>OMICS A Journal of Integrative Biology</i> , 2007 , 11, 341-50	3.8	3
20	GENETICS OF BODY COMPOSITION AND METABOLIC RATE 2005 , 131-160		3
19	Inferences on the Genetics of Quantitative Traits from Long-Term Selection in Laboratory and Domestic Animals169-210		3
18	Heterogeneous variances and genetics by environment interactions in genetic evaluation of crossbred lambs. <i>Animal</i> , 2015 , 9, 380-7	3.1	2
17	The effect of the Texel Muscling QTL on live and carcass weight in Texel lambs. <i>Small Ruminant Research</i> , 2012 , 105, 117-121	1.7	2
16	Evaluation of Video Image Analysis (VIA) technology to predict meat yield of sheep carcasses online under abattoir conditions. <i>Proceedings of the British Society of Animal Science</i> , 2007 , 2007, 108-108		2
15	Identification and reciprocal introgression of a QTL affecting body mass in mice. <i>Genetics Selection Evolution</i> , 2004 , 36, 577-91	4.9	2
14	Zur SelektionswEdigkeit von Merkmalen der Muskelstruktur IModellversuch mit LabormIIsen. Journal of Animal Breeding and Genetics, 1989 , 106, 208-216	2.9	2
13	Evaluating invasive and non-invasive methods to determine fat content in the laboratory mouse. <i>Open Life Sciences</i> , 2015 , 10,	1.2	1
12	Effect and mode of action of the Texel muscling QTL (TM-QTL) on carcass traits in purebred Texel lambs. <i>Animal</i> , 2014 , 8, 1053-61	3.1	1
11	Use of meat quality information in breeding programmes 2009 , 264-291		1
10	The effect of gestational undernutrition on maternal weight change and fetal weight in lines of mice selected for different growth characteristics. <i>British Journal of Nutrition</i> , 2011 , 105, 539-48	3.6	1
9	Prenatal growth in lines of mice selected for body weight. <i>Journal of Animal Breeding and Genetics</i> , 1996 , 113, 535-543	2.9	1
8	Selection for litter weight on the 21st day after long-term selection for first litter performance in laboratory mice. <i>Journal of Animal Breeding and Genetics</i> , 1990 , 107, 161-168	2.9	1

LIST OF PUBLICATIONS

7	Genetic evaluation of days to harvest in crossbred lambs. <i>Journal of Animal Science</i> , 2013 , 91, 5153-60	0.7
6	Investigation into the presence of genotype by environment (GE) interactions in Scottish Blackface lamb weaning traits 2012 , 23-31	
5	Effects of leptin administration on lines of mice selected long-term for fatness. <i>Genetical Research</i> , 1997 , 70, 79-89	1.1
4	The effects of three muscling Quantitative Trait Loci on growth patterns of crossbred lambs. <i>Proceedings of the British Society of Animal Science</i> , 2009 , 2009, 40-40	
3	Phenotypic characterisation of extreme growth-selected mouse lines: An important prerequisite for future QTL analysis. <i>Open Life Sciences</i> , 2006 , 1, 345-375	1.2
2	Effects of thyroid hormone deficiency on mice selected for increased and decreased body weight and fatness. <i>Genetical Research</i> , 1998 , 72, 59-72	1.1
1	Kreuzungswirkungen bei Merkmalen der Fruchtbarkeit und des Wachstums nach Langzeitselektion	2.9