Stein Tore Nilsen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/932680/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Preeclampsia and fetal growth. Obstetrics and Gynecology, 2000, 96, 950-955.	2.4	245
2	Insulin-like Growth Factor I and Leptin in Umbilical Cord Plasma and Infant Birth Size at Term. Pediatrics, 2002, 109, 1131-1135.	2.1	138
3	Size at Birth and Gestational Age as Predictors of Adult Height and Weight. Epidemiology, 2005, 16, 175-181.	2.7	121
4	Serum Urate as a Predictor of Fetal Outcome in Severe Pre-Eclampsia. Acta Obstetricia Et Gynecologica Scandinavica, 1984, 63, 71-75.	2.8	62
5	Maternal Hemoglobin Concentration is Closely Related to Birth Weight in Normal Pregnancies. Acta Obstetricia Et Gynecologica Scandinavica, 1984, 63, 245-248.	2.8	46
6	Males with Low Birthweight Examined at 18 Years of Age. Acta Paediatrica, International Journal of Paediatrics, 1984, 73, 168-175.	1.5	40
7	Boys Born by Forceps and Vacuum Extraction Examined at 18 Years of Age. Acta Obstetricia Et Gynecologica Scandinavica, 1984, 63, 549-554.	2.8	38
8	Males with Neonatal Hyperbilirubinemia Examined at 18 Years of Age. Acta Paediatrica, International Journal of Paediatrics, 1984, 73, 176-180.	1.5	38
9	Relationship of insulin-like growth factor-I and insulin-like growth factor binding proteins in umbilical cord plasma to preeclampsia and infant birth weight. Obstetrics and Gynecology, 2002, 99, 85-90.	2.4	38
10	Breech Delivery and Intelligence: A Population-Based Study of 8,738 Breech Infants. Obstetrics and Gynecology, 2005, 105, 4-11.	2.4	38
11	Preeclampsia and Fetal Growth. Obstetrics and Gynecology, 2000, 96, 950-955.	2.4	37
12	Maternal and Fetal Variants of Genetic Thrombophilias and the Risk of Preeclampsia. Epidemiology, 2004, 15, 317-322.	2.7	35
13	Umbilical cord plasma leptin is increased in preeclampsia. American Journal of Obstetrics and Gynecology, 2002, 186, 427-432.	1.3	30
14	Cervical intraepithelial neoplasia grade 3 lesions can regress. Apmis, 2007, 115, 1409-1414.	2.0	30
15	Serum 25(OH)D Is a 2-Year Predictor of All-Cause Mortality, Cardiac Death and Sudden Cardiac Death in Chest Pain Patients from Northern Argentina. PLoS ONE, 2012, 7, e43228.	2.5	27
16	Umbilical cord plasma interleukin-6 and fetal growth restriction in preeclampsia: a prospective study in Norway. Obstetrics and Gynecology, 2001, 98, 289-294.	2.4	19
17	Knowledge, attitudes and management strategies in Scandinavia concerning hormone replacement therapy. Maturitas, 2001, 39, 83-90.	2.4	16
18	Impact of recent studies on attitudes and use of hormone therapy among Scandinavian gynaecologists. Acta Obstetricia Et Gynecologica Scandinavica, 2007, 86, 1490-1495.	2.8	16

STEIN TORE NILSEN

#	Article	IF	CITATIONS
19	Male twins at birth and 18 years later. BJOG: an International Journal of Obstetrics and Gynaecology, 1984, 91, 122-127.	2.3	15
20	Patient safety challenges in a case study hospital – Of relevance for transfusion processes?. Transfusion and Apheresis Science, 2008, 39, 167-172.	1.0	14
21	Males Born in Breech Presentation 18 Years After Birth. Acta Obstetricia Et Gynecologica Scandinavica, 1985, 64, 323-325.	2.8	11
22	B-type natriuretic peptide and high sensitive C-reactive protein predict 2-year all cause mortality in chest pain patients: a prospective observational study from Salta, Argentina. BMC Cardiovascular Disorders, 2011, 11, 57.	1.7	11
23	In utero preeclampsia exposure, milk intake and pubertal development. Reproductive Toxicology, 2015, 54, 19-25.	2.9	11
24	Umbilical Cord Plasma Interleukin-6 and Fetal Growth Restriction in Preeclampsia. Obstetrics and Gynecology, 2001, 98, 289-294.	2.4	10
25	Vitamin D Predicts All-Cause and Cardiac Mortality in Females with Suspected Acute Coronary Syndrome: A Comparison with Brain Natriuretic Peptide and High-Sensitivity C-Reactive Protein. Cardiology Research and Practice, 2013, 2013, 1-8.	1.1	10
26	Preeclampsia and Adiponectin in Cord Blood. Hormone Research in Paediatrics, 2010, 74, 92-97.	1.8	9
27	The Predictive Value of Total Estriol; HPL and HB on Perinatal Outcome in Severe Pre-Eclampsia. Acta Obstetricia Et Gynecologica Scandinavica, 1984, 63, 603-608.	2.8	8
28	Suggested Cut-Off Values for Vitamin D as a Risk Marker for Total and Cardiac Death in Patients with Suspected Acute Coronary Syndrome. Frontiers in Cardiovascular Medicine, 2016, 3, 4.	2.4	7
29	Variations in serum concentrations of selected organochlorines among delivering women in Argentina. The EMASAR study. Environmental Sciences: Processes and Impacts, 2017, 19, 1542-1553.	3.5	6
30	Omega-3 index and prognosis in acute coronary chest pain patients with a low dietary intake of omega-3. Scandinavian Cardiovascular Journal, 2013, 47, 69-79.	1.2	4
31	Borderline Values of Troponin-T and High Sensitivity C-Reactive Protein Did Not Predict 2-Year Mortality in TnT Positive Chest-Pain Patients, Whereas Brain Natriuretic Peptide Did. Frontiers in Cardiovascular Medicine, 2015, 2, 16.	2.4	3
32	Socioeconomic Assessment and Impact of Social Security on Outcome in Patients Admitted with Suspected Coronary Chest Pain in the City of Salta, Argentina. Cardiology Research and Practice, 2013, 2013, 1-9.	1.1	2
33	Males born by cesarean section examined 18 years after delivery. Acta Obstetricia Et Gynecologica Scandinavica, 1985, 64, 237-240.	2.8	1
34	Relationship of Insulin-Like Growth Factor-I and Insulin-Like Growth Factor Binding Proteins in Umbilical Cord Plasma to Preeclampsia and Infant Birth Weight. Obstetrics and Gynecology, 2002, 99, 85-90.	2.4	1