Pere Mestre

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9330536/publications.pdf

Version: 2024-02-01

567144 940416 1,244 16 15 16 citations h-index g-index papers 18 18 18 1194 citing authors docs citations times ranked all docs

#	Article	IF	CITATIONS
1	Identification of the First Oomycete Mating-type Locus Sequence in the Grapevine Downy Mildew Pathogen, Plasmopara viticola. Current Biology, 2020, 30, 3897-3907.e4.	1.8	23
2	A secreted WY-domain-containing protein present in European isolates of the oomycete Plasmopara viticola induces cell death in grapevine and tobacco species. PLoS ONE, 2019, 14, e0220184.	1.1	25
3	Overexpression of the WSWEET4 Transporter in Grapevine Hairy Roots Increases Sugar Transport and Contents and Enhances Resistance to Pythium irregulare, a Soilborne Pathogen. Frontiers in Plant Science, 2019, 10, 884.	1.7	22
4	A High-Quality Grapevine Downy Mildew Genome Assembly Reveals Rapidly Evolving and Lineage-Specific Putative Host Adaptation Genes. Genome Biology and Evolution, 2019, 11, 954-969.	1.1	61
5	Introgression reshapes recombination distribution in grapevine interspecific hybrids. Theoretical and Applied Genetics, 2019, 132, 1073-1087.	1.8	19
6	Breeding for durable resistance to downy and powdery mildew in grapevine. Oeno One, 2018, 52, 203-209.	0.7	86
7	ldentification of a <i>Vitis vinifera</i> endoâ€ β ‶,3â€glucanase with antimicrobial activity against <i>Plasmopara viticola</i> Molecular Plant Pathology, 2017, 18, 708-719.	2.0	28
8	Draft Genome Sequence of <i>Plasmopara viticola</i> , the Grapevine Downy Mildew Pathogen. Genome Announcements, 2016, 4, .	0.8	29
9	The SWEET family of sugar transporters in grapevine: VvSWEET4 is involved in the interaction with Botrytis cinerea. Journal of Experimental Botany, 2014, 65, 6589-6601.	2.4	214
10	Geographic Distribution of Cryptic Species of <i>Plasmopara viticola</i> Causing Downy Mildew on Wild and Cultivated Grape in Eastern North America. Phytopathology, 2014, 104, 692-701.	1.1	51
11	Genetic dissection of a <scp>TIR</scp> â€ <scp>NB</scp> â€ <scp>LRR</scp> locus from the wild <scp>N</scp> orth <scp>A</scp> merican grapevine species <i><scp>M</scp>uscadinia rotundifolia</i> identifies paralogous genes conferring resistance to major fungal and oomycete pathogens in cultivated grapevine. Plant Journal, 2013, 76, 661-674.	2.8	152
12	Phylogenetic and experimental evidence for hostâ€specialized cryptic species in a biotrophic oomycete. New Phytologist, 2013, 197, 251-263.	3.5	110
13	A reference genetic map of Muscadinia rotundifolia and identification of Ren5, a new major locus for resistance to grapevine powdery mildew. Theoretical and Applied Genetics, 2012, 125, 1663-1675.	1.8	74
14	Identification of effector genes from the phytopathogenic Oomycete Plasmopara viticola through the analysis of gene expression in germinated zoospores. Fungal Biology, 2012, 116, 825-835.	1.1	52
15	Construction of a reference linkage map of Vitis amurensis and genetic mapping of Rpv8, a locus conferring resistance to grapevine downy mildew. Theoretical and Applied Genetics, 2011, 123, 43-53.	1.8	132
16	Breakdown of resistance to grapevine downy mildew upon limited deployment of a resistant variety. BMC Plant Biology, 2010, 10, 147.	1.6	162