

Angelique Stalmach

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9331991/publications.pdf>

Version: 2024-02-01

32
papers

2,815
citations

218381

26
h-index

433756

31
g-index

35
all docs

35
docs citations

35
times ranked

3867
citing authors

#	ARTICLE	IF	CITATIONS
1	Urinary proteomics predict onset of microalbuminuria in normoalbuminuric type 2 diabetic patients, a sub-study of the DIRECT-Protect 2 study. <i>Nephrology Dialysis Transplantation</i> , 2017, 32, gfw292.	0.4	66
2	Acute kidney injury prediction in cardiac surgery patients by a urinary peptide pattern: a case-control validation study. <i>Critical Care</i> , 2016, 20, 157.	2.5	24
3	Urinary proteomic biomarkers to predict cardiovascular events. <i>Proteomics - Clinical Applications</i> , 2015, 9, 610-617.	0.8	33
4	<i>In vitro</i> studies on the stability in the proximal gastrointestinal tract and bioaccessibility in Caco-2 cells of chlorogenic acids from spent coffee grounds. <i>International Journal of Food Sciences and Nutrition</i> , 2015, 66, 657-664.	1.3	34
5	Methods in Capillary Electrophoresis Coupled to Mass Spectrometry for the Identification of Clinical Proteomic/Peptidomic Biomarkers in Biofluids. <i>Methods in Molecular Biology</i> , 2015, 1243, 187-205.	0.4	20
6	Identification of Urinary Peptide Biomarkers Associated with Rheumatoid Arthritis. <i>PLoS ONE</i> , 2014, 9, e104625.	1.1	32
7	Dark chocolate consumption improves leukocyte adhesion factors and vascular function in overweight men. <i>FASEB Journal</i> , 2014, 28, 1464-1473.	0.2	53
8	Bioavailability of Dietary Anthocyanins and Hydroxycinnamic Acids. , 2014, , 561-576.		7
9	Classical MALDI-MS versus CE-based ESI-MS proteomic profiling in urine for clinical applications. <i>Bioanalysis</i> , 2014, 6, 247-266.	0.6	20
10	Impact of dose on the bioavailability of coffee chlorogenic acids in humans. <i>Food and Function</i> , 2014, 5, 1727-1737.	2.1	91
11	Recent advances in capillary electrophoresis coupled to mass spectrometry for clinical proteomic applications. <i>Electrophoresis</i> , 2013, 34, 1452-1464.	1.3	103
12	Fetal Urinary Peptides to Predict Postnatal Outcome of Renal Disease in Fetuses with Posterior Urethral Valves (PUV). <i>Science Translational Medicine</i> , 2013, 5, 198ra106.	5.8	86
13	Improving peptide relative quantification in MALDI-TOF MS for biomarker assessment. <i>Proteomics</i> , 2013, 13, 2967-2975.	1.3	21
14	Colonic catabolism of dietary phenolic and polyphenolic compounds from Concord grape juice. <i>Food and Function</i> , 2013, 4, 52-62.	2.1	70
15	Polyphenolic and Hydroxycinnamate Contents of Whole Coffee Fruits from China, India, and Mexico. <i>Journal of Agricultural and Food Chemistry</i> , 2013, 61, 5298-5309.	2.4	64
16	Absorption, Disposition, Metabolism, and Excretion of [¹⁴ C]Caffeic Acid in Rats. <i>Journal of Agricultural and Food Chemistry</i> , 2012, 60, 5205-5214.	2.4	40
17	Espresso coffees, caffeine and chlorogenic acid intake: potential health implications. <i>Food and Function</i> , 2012, 3, 30-33.	2.1	142
18	Gastrointestinal stability and bioavailability of (poly)phenolic compounds following ingestion of Concord grape juice by humans. <i>Molecular Nutrition and Food Research</i> , 2012, 56, 497-509.	1.5	106

#	ARTICLE	IF	CITATIONS
19	Identification of (Poly)phenolic Compounds in Concord Grape Juice and Their Metabolites in Human Plasma and Urine after Juice Consumption. <i>Journal of Agricultural and Food Chemistry</i> , 2011, 59, 9512-9522.	2.4	95
20	The Antioxidant and Chlorogenic Acid Profiles of Whole Coffee Fruits Are Influenced by the Extraction Procedures. <i>Journal of Agricultural and Food Chemistry</i> , 2011, 59, 3754-3762.	2.4	87
21	Reduction of monocyte chemoattractant protein 1 and macrophage migration inhibitory factor by a polyphenol-rich extract in subjects with clustered cardiometabolic risk factors. <i>British Journal of Nutrition</i> , 2011, 106, 1416-1422.	1.2	17
22	First synthesis, characterization, and evidence for the presence of hydroxycinnamic acid sulfate and glucuronide conjugates in human biological fluids as a result of coffee consumption. <i>Organic and Biomolecular Chemistry</i> , 2010, 8, 5199.	1.5	53
23	Absorption, metabolism, and excretion of green tea flavanols in humans with an ileostomy. <i>Molecular Nutrition and Food Research</i> , 2010, 54, 323-334.	1.5	178
24	In vitro and in vivo conjugation of dietary hydroxycinnamic acids by UDP-glucuronosyltransferases and sulfotransferases in humans. <i>Journal of Nutritional Biochemistry</i> , 2010, 21, 1060-1068.	1.9	61
25	Unfermented and fermented rooibos teas (<i>Aspalathus linearis</i>) increase plasma total antioxidant capacity in healthy humans. <i>Food Chemistry</i> , 2010, 123, 679-683.	4.2	40
26	Bioavailability of Coffee Chlorogenic Acids and Green Tea Flavan-3-ols. <i>Nutrients</i> , 2010, 2, 820-833.	1.7	98
27	Green Tea Flavan-3-ols: Colonic Degradation and Urinary Excretion of Catabolites by Humans. <i>Journal of Agricultural and Food Chemistry</i> , 2010, 58, 1296-1304.	2.4	229
28	Bioavailability of chlorogenic acids following acute ingestion of coffee by humans with an ileostomy. <i>Archives of Biochemistry and Biophysics</i> , 2010, 501, 98-105.	1.4	217
29	Absorption, metabolism and excretion of Choldi green tea flavanols by humans. <i>Molecular Nutrition and Food Research</i> , 2009, 53, S44-53.	1.5	190
30	Metabolite Profiling of Hydroxycinnamate Derivatives in Plasma and Urine after the Ingestion of Coffee by Humans: Identification of Biomarkers of Coffee Consumption. <i>Drug Metabolism and Disposition</i> , 2009, 37, 1749-1758.	1.7	343
31	Bioavailability of C-Linked Dihydrochalcone and Flavanone Glucosides in Humans Following Ingestion of Unfermented and Fermented Rooibos Teas. <i>Journal of Agricultural and Food Chemistry</i> , 2009, 57, 7104-7111.	2.4	86
32	On-line HPLC analysis of the antioxidant activity of phenolic compounds in brewed, paper-filtered coffee. <i>Brazilian Journal of Plant Physiology</i> , 2006, 18, 253-262.	0.5	94