

Hideyuki Tamai

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9341391/publications.pdf>

Version: 2024-02-01

47
papers

783
citations

567281

15
h-index

526287

27
g-index

47
all docs

47
docs citations

47
times ranked

1310
citing authors

#	ARTICLE	IF	CITATIONS
1	Feeding artery ablation before radiofrequency ablation for hepatocellular carcinoma may reduce critical recurrence. <i>JGH Open</i> , 2021, 5, 478-485.	1.6	4
2	Measurements of Serum Mac-2-Binding Protein Glycosylation Isomer and Shear Wave Velocity in Health Checkups Are Useful in Screening for Non-Alcoholic Steatohepatitis. <i>Healthcare (Switzerland)</i> , 2021, 9, 523.	2.0	0
3	New next-generation microwave thermosphere ablation for small hepatocellular carcinoma. <i>Clinical and Molecular Hepatology</i> , 2021, 27, 564-574.	8.9	15
4	Prognostic value of 18F-fluorodeoxyglucose positron emission tomography in patients with small hepatocellular carcinoma treated by radiofrequency ablation. <i>Cancer Imaging</i> , 2020, 20, 74.	2.8	2
5	Combination with portosystemic shunt occlusion and antiviral therapy improves prognosis of decompensated cirrhosis. <i>JGH Open</i> , 2020, 4, 670-676.	1.6	1
6	Sofosbuvir plus ribavirin is tolerable and effective even in elderly patients 75-years-old and over. <i>World Journal of Hepatology</i> , 2020, 12, 672-684.	2.0	0
7	Trends and Efficacy of Interferon-Free Anti-hepatitis C Virus Therapy in the Region of High Prevalence of Elderly Patients, Cirrhosis, and Hepatocellular Carcinoma: A Real-World, Nationwide, Multicenter Study of 10 688 Patients in Japan. <i>Open Forum Infectious Diseases</i> , 2019, 6, ofz185.	0.9	18
8	Real-world safety and efficacy of sofosbuvir and ledipasvir for elderly patients. <i>JGH Open</i> , 2018, 2, 300-306.	1.6	5
9	Real-world virological efficacy and safety of elbasvir and grazoprevir in patients with chronic hepatitis C virus genotype 1 infection in Japan. <i>Journal of Gastroenterology</i> , 2018, 53, 1276-1284.	5.1	25
10	The Real-World Safety and Efficacy of Daclatasvir and Asunaprevir for Elderly Patients. <i>Gut and Liver</i> , 2018, 12, 86-93.	2.9	6
11	Irregular Defects in Hepatocellular Carcinomas During the Kupffer Phase of Contrast-Enhanced Ultrasonography with Perfluorobutane Microbubbles: Pathological Features and Metastatic Recurrence After Surgical Resection. <i>Ultrasound in Medicine and Biology</i> , 2017, 43, 1829-1836.	1.5	7
12	Add-on Pegylated Interferon Alpha-2a Therapy in Chronic Hepatitis B Japanese Patients Treated with Entecavir. <i>Hepatitis Research and Treatment</i> , 2017, 2017, 1-7.	2.0	2
13	Simeprevir-Based Triple Therapy with Reduced Doses of Pegylated Interferon α -2a Plus Ribavirin for Interferon Ineligible Patients with Genotype 1b Hepatitis C Virus. <i>Gut and Liver</i> , 2017, 11, 551-558.	2.9	3
14	Rapid On-Site Evaluation by Endosonographers during Endoscopic Ultrasonography-Guided Fine-Needle Aspiration for Diagnosis of Gastrointestinal Stromal Tumors. <i>Clinical Endoscopy</i> , 2017, 50, 372-378.	1.5	18
15	The Wire-Grasping Method as a New Technique for Forceps Biopsy of Biliary Strictures: A Prospective Randomized Controlled Study of Effectiveness. <i>Gut and Liver</i> , 2016, 10, 642-648.	2.9	6
16	Endoscopic features associated with development of metachronous gastric cancer in patients who underwent endoscopic resection followed by <i>Helicobacter pylori</i> eradication. <i>Digestive Endoscopy</i> , 2016, 28, 434-442.	2.3	29
17	Serum pepsinogen levels can quantify the risk of development of metachronous gastric cancer after endoscopic resection. <i>International Journal of Cancer</i> , 2016, 139, 1150-1156.	5.1	12
18	Hypointense hepatocellular carcinomas on apparent diffusion coefficient mapping: Pathological features and metastatic recurrence after hepatectomy. <i>Hepatology Research</i> , 2016, 46, 634-641.	3.4	8

#	ARTICLE	IF	CITATIONS
19	Contribution of Contrast-Enhanced Sonography With Perfluorobutane Microbubbles for Diagnosis of Recurrent Hepatocellular Carcinoma. <i>Journal of Ultrasound in Medicine</i> , 2016, 35, 1383-1391.	1.7	8
20	The arterial tumor enhancement pattern on contrast-enhanced computed tomography is associated with primary cancer death after radiofrequency ablation for small hepatocellular carcinoma. <i>Hepatology International</i> , 2016, 10, 328-339.	4.2	14
21	Kupffer Imaging by Contrast-Enhanced Sonography With Perfluorobutane Microbubbles Is Associated With Outcomes After Radiofrequency Ablation of Hepatocellular Carcinoma. <i>Journal of Ultrasound in Medicine</i> , 2016, 35, 359-371.	1.7	7
22	Low-Dose Pegylated Interferon \pm 2b Plus Ribavirin for Elderly and/or Cirrhotic Patients with Genotype 2 Hepatitis C Virus. <i>Gut and Liver</i> , 2016, 10, 617-623.	2.9	5
23	Video capsule endoscopy as the initial examination for overt obscure gastrointestinal bleeding can efficiently identify patients who require double-balloon enteroscopy. <i>BMC Gastroenterology</i> , 2015, 15, 132.	2.0	15
24	Contrast-Enhanced Endoscopic Ultrasonography for Pancreatic Tumors. <i>BioMed Research International</i> , 2015, 2015, 1-8.	1.9	36
25	Prognostic impact of surgery and radiofrequency ablation on single nodular HCC \leq 2 cm: Cohort study based on serum HCC markers. <i>Journal of Hepatology</i> , 2015, 63, 1352-1359.	3.7	57
26	Prognostic effect of response to interferon therapy after laparoscopic splenectomy among patients with marked thrombocytopenia and hepatitis C virus-related cirrhosis. <i>Hepatology International</i> , 2015, 9, 67-75.	4.2	4
27	Japanese apricot improves symptoms of gastrointestinal dysmotility associated with gastroesophageal reflux disease. <i>World Journal of Gastroenterology</i> , 2015, 21, 8170.	3.3	8
28	<i>Helicobacter pylori</i> -related chronic gastritis as a risk factor for colonic neoplasms. <i>World Journal of Gastroenterology</i> , 2014, 20, 1485.	3.3	24
29	N-Butyl Cyanoacrylate Embolization with Blood Flow Control of an Arterioportal Shunt That Developed after Radiofrequency Ablation of Hepatocellular Carcinoma. <i>Korean Journal of Radiology</i> , 2014, 15, 250.	3.4	15
30	Prediction of Sustained Virological Response to Telaprevir-Based Triple Therapy Using Viral Response within 2 Weeks. <i>Hepatitis Research and Treatment</i> , 2014, 2014, 1-8.	2.0	4
31	An open-label phase I/II study of tamibarotene in patients with advanced hepatocellular carcinoma. <i>Hepatology International</i> , 2014, 8, 94-103.	4.2	11
32	Prediction of a Null Response to Pegylated Interferon \pm 2b Plus Ribavirin in Patients with High Viral Load Genotype 1b Hepatitis C. <i>Gut and Liver</i> , 2014, 8, 421-427.	2.9	3
33	Tumor Vessel Depiction With Contrast-Enhanced Endoscopic Ultrasonography Predicts Efficacy of Chemotherapy in Pancreatic Cancer. <i>Pancreas</i> , 2013, 42, 990-995.	1.1	31
34	Low-dose pegylated interferon- α 2a monotherapy in elderly and/or cirrhotic patients infected with hepatitis C virus genotype 2 or genotype 1 low level infection. <i>Hepatology Research</i> , 2013, 43, 702-708.	3.4	1
35	Serological and histological indices of hepatocellular carcinoma and tumor volume doubling time. <i>Molecular and Clinical Oncology</i> , 2013, 1, 977-981.	1.0	18
36	Low-dose pegylated interferon- α 2a plus ribavirin therapy for elderly and/or cirrhotic patients with hepatitis C virus genotype-1b and high viral load. <i>Antiviral Therapy</i> , 2013, 19, 107-115.	1.0	4

#	ARTICLE	IF	CITATIONS
37	Usefulness of Contrast-Enhanced Endoscopic Sonography for Discriminating Mural Nodules From Mucous Clots in Intraductal Papillary Mucinous Neoplasms. <i>Journal of Ultrasound in Medicine</i> , 2013, 32, 61-68.	1.7	87
38	Elevated risk of recurrent colorectal neoplasia with <i>Helicobacter pylori</i> -associated chronic atrophic gastritis: A follow-up study of patients with endoscopically resected colorectal neoplasia. <i>Molecular and Clinical Oncology</i> , 2013, 1, 75-82.	1.0	7
39	Common Bile Duct Dilatation With Stones Indicates Requirement for Early Drainage in Patients With or Without Cholangitis. <i>Gastroenterology Research</i> , 2013, 6, 219-226.	1.3	4
40	Clinical features and risk factors of extrahepatic seeding after percutaneous radiofrequency ablation for hepatocellular carcinoma. <i>Hepatology Research</i> , 2011, 41, 738-745.	3.4	39
41	Prediction of sustained response to low-dose pegylated interferon alpha-2b plus ribavirin in patients with genotype 1b and high hepatitis C virus level using viral reduction within 2 weeks after therapy initiation. <i>Hepatology Research</i> , 2011, 41, 1137-1144.	3.4	11
42	<i>A Laparoscopic Splenectomy Allows the Induction of Antiviral Therapy for Patients with Cirrhosis Associated with Hepatitis C Virus</i> . <i>American Surgeon</i> , 2011, 77, 174-179.	0.8	11
43	Diagnosis of pancreatic tuberculosis by combined, contrast-enhanced sonography and endoscopic ultrasound-guided fine-needle aspiration. <i>Clinical Journal of Gastroenterology</i> , 2010, 3, 159-164.	0.8	6
44	Diffuse intrahepatic recurrence after percutaneous radiofrequency ablation for solitary and small hepatocellular carcinoma. <i>Hepatology International</i> , 2009, 3, 509-515.	4.2	42
45	Multifocal Nodular Fatty Infiltration of the Liver Mimicking Metastatic Liver Tumors. <i>Journal of Ultrasound in Medicine</i> , 2006, 25, 403-406.	1.7	7
46	Contrast-Enhanced Ultrasonography in the Diagnosis of Solid Renal Tumors. <i>Journal of Ultrasound in Medicine</i> , 2005, 24, 1635-1640.	1.7	139
47	Contrast Harmonic Sonographically Guided Radio Frequency Ablation for Spontaneous Ruptured Hepatocellular Carcinoma. <i>Journal of Ultrasound in Medicine</i> , 2005, 24, 1021-1026.	1.7	4