

Mizuko Mamura

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9350820/mizuko-mamura-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

48
papers

3,722
citations

28
h-index

57
g-index

57
ext. papers

3,971
ext. citations

7.9
avg, IF

4.13
L-index

#	Paper	IF	Citations
48	Repression of Smad3 by Stat3 and c-Ski/SnoN induces gefitinib resistance in lung adenocarcinoma. <i>Biochemical and Biophysical Research Communications</i> , 2017 , 484, 269-277	3.4	16
47	Phosphorylation status determines the opposing functions of Smad2/Smad3 as STAT3 cofactors in TH17 differentiation. <i>Nature Communications</i> , 2015 , 6, 7600	17.4	68
46	Inhibition of lethal inflammatory responses through the targeting of membrane-associated Toll-like receptor 4 signaling complexes with a Smad6-derived peptide. <i>EMBO Molecular Medicine</i> , 2015 , 7, 577-92 ¹²		22
45	Activin receptor-like kinase5 inhibition suppresses mouse melanoma by ubiquitin degradation of Smad4, thereby derepressing eomesodermin in cytotoxic T lymphocytes. <i>EMBO Molecular Medicine</i> , 2014 , 6, 703-703	12	78
44	Proangiogenic TIE2(+)/CD31 (+) macrophages are the predominant population of tumor-associated macrophages infiltrating metastatic lymph nodes. <i>Molecules and Cells</i> , 2013 , 36, 432-8	3.5	24
43	Activin receptor-like kinase5 inhibition suppresses mouse melanoma by ubiquitin degradation of Smad4, thereby derepressing eomesodermin in cytotoxic T lymphocytes. <i>EMBO Molecular Medicine</i> , 2013 , 5, 1720-39	12	45
42	Association of variants in PPAR α , IGF2BP2, and KCNQ1 with a susceptibility to gestational diabetes mellitus in a Korean population. <i>Yonsei Medical Journal</i> , 2013 , 54, 352-7	3	30
41	Inhibition of erythropoiesis by Smad6 in human cord blood hematopoietic stem cells. <i>Biochemical and Biophysical Research Communications</i> , 2012 , 423, 750-6	3.4	15
40	Regulation of tumor immune surveillance and tumor immune subversion by tgf-Beta. <i>Immune Network</i> , 2009 , 9, 122-6	6.1	13
39	IN-1130, a novel transforming growth factor-beta type I receptor kinase (activin receptor-like kinase 5) inhibitor, promotes regression of fibrotic plaque and corrects penile curvature in a rat model of Peyronie's disease. <i>Journal of Sexual Medicine</i> , 2009 , 6, 1284-96	1.1	33
38	Role of increased penile expression of transforming growth factor-beta1 and activation of the Smad signaling pathway in erectile dysfunction in streptozotocin-induced diabetic rats. <i>Journal of Sexual Medicine</i> , 2008 , 5, 2318-29	1.1	45
37	Absence of Smad3 induces neutrophil migration after cutaneous irradiation: possible contribution to subsequent radioprotection. <i>American Journal of Pathology</i> , 2008 , 173, 68-76	5.8	14
36	An anti-transforming growth factor beta antibody suppresses metastasis via cooperative effects on multiple cell compartments. <i>Cancer Research</i> , 2008 , 68, 3835-43	10.1	188
35	Interleukin-17 gene expression in patients with rheumatoid arthritis. <i>Modern Rheumatology</i> , 2008 , 18, 15-22	3.3	28
34	Intravascular large B-cell lymphoma with acute abdomen as a presenting symptom in a patient with systemic lupus erythematosus. <i>Journal of Clinical Oncology</i> , 2008 , 26, 1553-5	2.2	9
33	Transforming growth factor beta subverts the immune system into directly promoting tumor growth through interleukin-17. <i>Cancer Research</i> , 2008 , 68, 3915-23	10.1	203
32	Crucial role of the interleukin-6/interleukin-17 cytokine axis in the induction of arthritis by glucose-6-phosphate isomerase. <i>Arthritis and Rheumatism</i> , 2008 , 58, 754-63		105

31	Interleukin-17 gene expression in patients with rheumatoid arthritis. <i>Modern Rheumatology</i> , 2008 , 18, 15-22	3.3	17
30	Effects of infliximab therapy on gene expression levels of tumor necrosis factor alpha, tristetraprolin, T cell intracellular antigen 1, and Hu antigen R in patients with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2007 , 56, 2160-9		14
29	Transforming growth factor-beta can suppress tumorigenesis through effects on the putative cancer stem or early progenitor cell and committed progeny in a breast cancer xenograft model. <i>Cancer Research</i> , 2007 , 67, 8643-52	10.1	78
28	Successful treatment of refractory thrombotic thrombocytopenic purpura with cyclosporine and corticosteroids in a patient with systemic lupus erythematosus and antibodies to ADAMTS13. <i>Internal Medicine</i> , 2007 , 46, 1033-7	1.1	9
27	Anti-interleukin-6 receptor antibody (tocilizumab) treatment of multicentric Castleman's disease. <i>Internal Medicine</i> , 2007 , 46, 771-4	1.1	70
26	Sarcoid myositis with muscle weakness as a presenting symptom. <i>Modern Rheumatology</i> , 2007 , 17, 243-246	3.3	5
25	Sarcoid myositis with muscle weakness as a presenting symptom. <i>Modern Rheumatology</i> , 2007 , 17, 243-6	3.3	1
24	Overexpression of phosphorylated STAT-1alpha in the labial salivary glands of patients with Sjögren's syndrome. <i>Arthritis and Rheumatism</i> , 2006 , 54, 3476-84		28
23	Significance of antiprothrombin antibodies in patients with systemic lupus erythematosus: clinical evaluation of the antiprothrombin assay and the antiphosphatidylserine/prothrombin assay, and comparison with other antiphospholipid antibody assays. <i>Modern Rheumatology</i> , 2006 , 16, 158-164	3.3	28
22	Diagnosis of chlamydia-induced reactive arthritis. <i>Internal Medicine</i> , 2006 , 45, 37	1.1	
21	Expression of TNF- α , tristetraprolin, T-cell intracellular antigen-1 and Hu antigen R genes in synovium of patients with rheumatoid arthritis. <i>International Journal of Molecular Medicine</i> , 2006 , 18, 273	4.4	3
20	Smad4 signalling in T cells is required for suppression of gastrointestinal cancer. <i>Nature</i> , 2006 , 441, 1015-9	5.4	264
19	Significance of antiprothrombin antibodies in patients with systemic lupus erythematosus: clinical evaluation of the antiprothrombin assay and the antiphosphatidylserine/prothrombin assay, and comparison with other antiphospholipid antibody assays. <i>Modern Rheumatology</i> , 2006 , 16, 158-64	3.3	14
18	Expression of TNF-alpha, tristetraprolin, T-cell intracellular antigen-1 and Hu antigen R genes in synovium of patients with rheumatoid arthritis. <i>International Journal of Molecular Medicine</i> , 2006 , 18, 273-8	4.4	17
17	Identification of three novel peptides that inhibit CD40-CD154 interaction. <i>Modern Rheumatology</i> , 2005 , 15, 423-426	3.3	11
16	TGF-beta1 production by CD4+ CD25+ regulatory T cells is not essential for suppression of intestinal inflammation. <i>European Journal of Immunology</i> , 2005 , 35, 2886-95	6.1	105
15	Identification of three novel peptides that inhibit CD40-CD154 interaction. <i>Modern Rheumatology</i> , 2005 , 15, 423-6	3.3	8
14	Loss of Smad3 in acute T-cell lymphoblastic leukemia. <i>New England Journal of Medicine</i> , 2004 , 351, 552-9	9.2	130

13	CD28 disruption exacerbates inflammation in Tgf-beta1-/- mice: in vivo suppression by CD4+CD25+ regulatory T cells independent of autocrine TGF-beta1. <i>Blood</i> , 2004 , 103, 4594-601	2.2	68
12	Transforming growth factor-beta production and myeloid cells are an effector mechanism through which CD1d-restricted T cells block cytotoxic T lymphocyte-mediated tumor immunosurveillance: abrogation prevents tumor recurrence. <i>Journal of Experimental Medicine</i> , 2003 , 198, 1741-52	16.6	472
11	CD4(+)CD25(+) regulatory T cells can mediate suppressor function in the absence of transforming growth factor beta1 production and responsiveness. <i>Journal of Experimental Medicine</i> , 2002 , 196, 237-46	16.6	515
10	Transforming growth factor beta 1 induces apoptosis through cleavage of BAD in a Smad3-dependent mechanism in FaO hepatoma cells. <i>Molecular and Cellular Biology</i> , 2002 , 22, 1369-78	4.8	96
9	Raloxifene, a mixed estrogen agonist/antagonist, induces apoptosis through cleavage of BAD in TSU-PR1 human cancer cells. <i>Journal of Biological Chemistry</i> , 2002 , 277, 32510-5	5.4	47
8	Tumor necrosis factor induces apoptosis in hepatoma cells by increasing Ca(2+) release from the endoplasmic reticulum and suppressing Bcl-2 expression. <i>Journal of Biological Chemistry</i> , 2002 , 277, 31381-9	5.4	46
7	Lifetime exposure to a soluble TGF-beta antagonist protects mice against metastasis without adverse side effects. <i>Journal of Clinical Investigation</i> , 2002 , 109, 1607-15	15.9	157
6	Lifetime exposure to a soluble TGF-β antagonist protects mice against metastasis without adverse side effects. <i>Journal of Clinical Investigation</i> , 2002 , 109, 1607-1615	15.9	313
5	Constitutive phosphorylation and nuclear localization of Smad3 are correlated with increased collagen gene transcription in activated hepatic stellate cells. <i>Journal of Cellular Physiology</i> , 2001 , 187, 117-23	7	104
4	Blockade of TGF-beta signaling in T cells prevents the development of experimental glomerulonephritis. <i>Journal of Immunology</i> , 2001 , 166, 2818-23	5.3	48
3	Ligation of the T cell receptor complex results in phosphorylation of Smad2 in T lymphocytes. <i>Biochemical and Biophysical Research Communications</i> , 2000 , 268, 124-7	3.4	6
2	Tumor necrosis factor-alpha mediates antiapoptotic signals partially via p38 MAP kinase activation in human eosinophils. <i>International Archives of Allergy and Immunology</i> , 1999 , 120 Suppl 1, 54-9	3.7	25
1	SMAD3 Determines Conventional versus Plasmacytoid Dendritic Cell Fates		1