

# Tetsuya Ohira

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9351643/publications.pdf>

Version: 2024-02-01

192  
papers

5,504  
citations

116194

36  
h-index

116156

66  
g-index

203  
all docs

203  
docs citations

203  
times ranked

6210  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Dietary fiber intake and risk of incident disabling dementia: the Circulatory Risk in Communities Study. <i>Nutritional Neuroscience</i> , 2023, 26, 148-155.   | 1.5 | 13        |
| 2  | Impact of Major Cardiovascular Risk Factors on the Incidence of Cardiovascular Disease among Overweight and Non-Overweight Individuals: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 422-437.                                | 0.9 | 6         |
| 3  | Association Between Lifestyle Habits and the Prevalence of Abdominal Obesity After the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>Journal of Epidemiology</i> , 2022, 32, 496-501.   | 1.1 | 4         |
| 4  | Associations Between Occupational Status, Support at Work, and Salivary Cortisol Levels. <i>International Journal of Behavioral Medicine</i> , 2022, 29, 299-307.   | 0.8 | 3         |
| 5  | Accurate endoscopic identification of the afferent limb at the Y anastomosis using the fold disruption sign after gastric resection with Roux-Y reconstruction. <i>Digestive Endoscopy</i> , 2022, 34, 238-243.   | 1.3 | 5         |
| 6  | The Association between Parenting Confidence and Later Child Mental Health in the Area Affected by the Fukushima Nuclear Disaster: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 476.                            | 1.2 | 0         |
| 7  | Does laughing with others lower the risk of functional disability among older Japanese adults? The JAGES prospective cohort study. <i>Preventive Medicine</i> , 2022, 155, 106945.  | 1.6 | 6         |
| 8  | Lifestyle Factors Associated with Undernutrition in Older People after the Great East Japan Earthquake: A Prospective Study in the Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 3399.                               | 1.2 | 1         |
| 9  | Effects of a laughter program on body weight and mental health among Japanese people with metabolic syndrome risk factors: a randomized controlled trial. <i>BMC Geriatrics</i> , 2022, 22, 361.  | 1.1 | 3         |
| 10 | The Associations between Evacuation Status and Lifestyle-Related Diseases in Fukushima after the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 5661.                                | 1.2 | 1         |
| 11 | Lifestyle-related diseases caused by evacuation: Results of the comprehensive health check in the Fukushima health management survey. , 2022, , 99-121.   |     | 0         |
| 12 | Development of an Application for Sustainable Support of Returning Residents Displaced by the Fukushima Nuclear Accident. <i>Studies in Health Technology and Informatics</i> , 2022, , .   | 0.2 | 0         |
| 13 | Changes in the proportion of anemia among young women after the Great East Japan Earthquake: the Fukushima health management survey. <i>Scientific Reports</i> , 2022, 12, .  | 1.6 | 2         |
| 14 | Impact of lifestyle and psychosocial factors on the onset of hypertension after the Great East Japan earthquake: a 7-year follow-up of the Fukushima Health Management Survey. <i>Hypertension Research</i> , 2022, 45, 1609-1621.  | 1.5 | 2         |
| 15 | Does Laughter Predict Onset of Functional Disability and Mortality Among Older Japanese Adults? The JAGES Prospective Cohort Study. <i>Journal of Epidemiology</i> , 2021, 31, 301-307.   | 1.1 | 20        |
| 16 | Dietary pattern changes in Fukushima residents after the Great East Japan Earthquake: the Fukushima Health Management Survey 2011-2013. <i>Public Health Nutrition</i> , 2021, 24, 2195-2204.   | 1.1 | 6         |
| 17 | Longitudinal Trends in Blood Pressure Associated With the Frequency of Laughter: The Circulatory Risk in Communities Study (CIRCS), a Longitudinal Study of the Japanese General Population. <i>Journal of Epidemiology</i> , 2021, 31, 125-131.  | 1.1 | 6         |
| 18 | Association of Admission Hyperglycemia with Clinical Outcomes in Japanese Patients with Acute Large Vessel Occlusion Stroke: A post hoc Analysis of the Recovery by Endovascular Salvage for Cerebral Ultra-Acute Embolism Japan Registry 2. <i>Cerebrovascular Diseases</i> , 2021, 50, 12-19. | 0.8 | 8         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Higher psychological distress experienced by evacuees relocating outside Fukushima after the nuclear accident: The Fukushima Health Management Survey. <i>International Journal of Disaster Risk Reduction</i> , 2021, 52, 101962.   | 1.8 | 9         |
| 20 | Salt taste perception and blood pressure levels in population-based samples: the Circulatory Risk in Communities Study (CIRCS). <i>British Journal of Nutrition</i> , 2021, 125, 203-211.  | 1.2 | 2         |
| 21 | Impact of Perceived Social Support on the Association Between Anger Expression and the Risk of Stroke: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Epidemiology</i> , 2021, , .   | 1.1 | 2         |
| 22 | Psychosocial support for the examinees and their families during the secondary confirmatory examination: Analyses of support records at first visit. <i>Fukushima Journal of Medical Sciences</i> , 2021, 67, 53-63.   | 0.1 | 1         |
| 23 | Socioeconomic status, damage-related conditions, and PTSD following the Fukushima-daiichi nuclear power plant accident: The Fukushima Health Management Survey. <i>Fukushima Journal of Medical Sciences</i> , 2021, 67, 71-82.  | 0.1 | 4         |
| 24 | Dietary Patterns and Progression of Impaired Kidney Function in Japanese Adults: A Longitudinal Analysis for the Fukushima Health Management Survey, 2011â€”2015. <i>Nutrients</i> , 2021, 13, 168.  | 1.7 | 7         |
| 25 | Relationship Between Risk of Hyper-Low-density Lipoprotein Cholesterolemia and Evacuation After the Great East Japan Earthquake. <i>Journal of Epidemiology</i> , 2021, , .  | 1.1 | 4         |
| 26 | Serum High-Sensitivity C-Reactive Protein Levels and the Risk of Atrial Fibrillation in Japanese Population: the Circulatory Risk in Communities Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, 28, 194-202.  | 0.9 | 11        |
| 27 | Serum Albumin and Risks of Stroke and Its Subtypesâ€” The Circulatory Risk in Communities Study (CIRCS) â€”. <i>Circulation Journal</i> , 2021, 85, 385-392.   | 0.7 | 8         |
| 28 | The association between body mass index and recovery from post-traumatic stress disorder after the nuclear accident in Fukushima. <i>Scientific Reports</i> , 2021, 11, 5330.  | 1.6 | 2         |
| 29 | Response to the Letter to the Editor: â€”Absorbed radiation doses in the thyroid as estimated by UNSCEAR and subsequent risk of childhood thyroid cancer following the Great East Japan Earthquakeâ€”™, by Ohira et al.. <i>Journal of Radiation Research</i> , 2021, 62, 425-426.             | 0.8 | 1         |
| 30 | Association of habitual exercise with adultsâ€™™ mental health following the Fukushima Daiichi nuclear power plant accident: the Fukushima Health Management Survey. <i>Mental Health and Physical Activity</i> , 2021, 20, 100388.  | 0.9 | 1         |
| 31 | Evacuation after the Great East Japan Earthquake is an independent factor associated with hyperuricemia: The Fukushima Health Management Survey. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 1177-1188.   | 1.1 | 9         |
| 32 | Trajectories of peer relationship problems and emotional symptoms in children 5Â”years after a nuclear disaster: Fukushima Health Management Survey. <i>Journal of Radiation Research</i> , 2021, 62, i114-i121.   | 0.8 | 2         |
| 33 | Factors Hindering Social Participation among Older Residents from Evacuation Zones after the Nuclear Power Plant Accident in Fukushima: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 4426.                     | 1.2 | 3         |
| 34 | Relationship between physical activity/exercise habits and the frequency of new onset of lifestyle-related diseases after the Great East Japan Earthquake among residents in Fukushima: the Fukushima Health Management Survey. <i>Journal of Radiation Research</i> , 2021, 62, i129-i139.    | 0.8 | 5         |
| 35 | Needs Survey for Health Support Application Development Project for Residents Returning from Evacuation After the Fukushima Nuclear Accident. <i>Studies in Health Technology and Informatics</i> , 2021, 281, 1091-1092.  | 0.2 | 0         |
| 36 | Association between Psychosocial Factors and Oral Symptoms among Residents in Fukushima after the Great East Japan Earthquake: A Cross-Sectional Study from the Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 6054. | 1.2 | 1         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Relationship between Endothelial Dysfunction and Prevalence of Chronic Kidney Disease: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, 28, 622-629.  | 0.9 | 8         |
| 38 | Associations of the COVID-19 pandemic with the economic status and mental health of people affected by the Fukushima disaster using the difference-in-differences method: The Fukushima Health Management Survey. <i>SSM - Population Health</i> , 2021, 14, 100801.                     | 1.3 | 6         |
| 39 | Synergistic Effect of History of Cardiovascular Disease and Mental Distress on Post-Traumatic Stress Disorder after the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 10283. | 1.2 | 0         |
| 40 | Association between post-traumatic stress disorder symptoms and bone fractures after the Great East Japan Earthquake in older adults: a prospective cohort study from the Fukushima Health Management Survey. <i>BMC Geriatrics</i> , 2021, 21, 18.                                      | 1.1 | 7         |
| 41 | Association between frequency of laughter and oral health among community-dwelling older adults: a population-based cross-sectional study in Japan. <i>Quality of Life Research</i> , 2021, 30, 1561-1569.   | 1.5 | 6         |
| 42 | Fine Needle Aspiration Cytology Implementation and Malignancy Rates in Children and Adolescents Based on Japanese Guidelines: The Fukushima Health Management Survey. <i>Thyroid</i> , 2021, 31, 1683-1692.  | 2.4 | 4         |
| 43 | Association between Laughter and Lifestyle Diseases after the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 12699.   | 1.2 | 1         |
| 44 | Factors Influencing the Proportion of Non-examinees in the Fukushima Health Management Survey for Childhood and Adolescent Thyroid Cancer: Results From the Baseline Survey. <i>Journal of Epidemiology</i> , 2020, 30, 301-308.   | 1.1 | 3         |
| 45 | Communication between Health Professionals and Community Residents in Fukushima: A Focus on the Feedback Loop. <i>Health Communication</i> , 2020, 35, 1274-1282.  | 1.8 | 17        |
| 46 | Changes in the mental health status of adolescents following the Fukushima Daiichi nuclear accident and related factors: Fukushima Health Management Survey. <i>Journal of Affective Disorders</i> , 2020, 260, 432-439.   | 2.0 | 12        |
| 47 | The association between self-reported sleep dissatisfaction after the Great East Japan Earthquake, and a deteriorated socioeconomic status in the evacuation area: the Fukushima Health Management Survey. <i>Sleep Medicine</i> , 2020, 68, 63-70.                                      | 0.8 | 11        |
| 48 | AGE DEPENDENCE OF INDIVIDUAL EXTERNAL DOSES IN AN EARLY STAGE AFTER THE FUKUSHIMA NUCLEAR ACCIDENT. <i>Radiation Protection Dosimetry</i> , 2020, 188, 238-245.  | 0.4 | 4         |
| 49 | Associations between Dietary Patterns and Cardiometabolic Risks in Japan: A Cross-Sectional Study from the Fukushima Health Management Survey, 2011-2015. <i>Nutrients</i> , 2020, 12, 129.  | 1.7 | 13        |
| 50 | Blood concentration of tacrolimus and age predict tacrolimus-induced left ventricular dysfunction after bone marrow transplantation in adults. <i>Journal of Medical Ultrasonics (2001)</i> , 2020, 47, 97-105.  | 0.6 | 2         |
| 51 | Anger Expression and the Risk of Cardiovascular Disease Among Urban and Rural Japanese Residents: The Circulatory Risk in Communities Study. <i>Psychosomatic Medicine</i> , 2020, 82, 215-223.  | 1.3 | 10        |
| 52 | The Impact of Valvuloarterial Impedance on Left Ventricular Geometrical Change after Transcatheter Aortic Valve Replacement: A Comparison between Valvuloarterial Impedance and Mean Pressure Gradient. <i>Journal of Clinical Medicine</i> , 2020, 9, 3143.                             | 1.0 | 0         |
| 53 | Nested matched case control study for the Japan Fukushima Health Management Survey's first full-scale (second-round) thyroid examination. <i>Medicine (United States)</i> , 2020, 99, e20440.  | 0.4 | 6         |
| 54 | Influence of post-disaster evacuation on incidence of hyperuricemia in residents of Fukushima Prefecture: the Fukushima Health Management Survey. <i>Clinical and Experimental Nephrology</i> , 2020, 24, 1025-1032.   | 0.7 | 7         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | All-Cause Mortality After the Great East Japan Earthquake in Fukushima Prefecture: Trends From 2009 to 2016 and Variation by Displacement. <i>Disaster Medicine and Public Health Preparedness</i> , 2020, , 1-4.   | 0.7 | 1         |
| 56 | Association between Psychological Factors and Evacuation Status and the Incidence of Cardiovascular Diseases after the Great East Japan Earthquake: A Prospective Study of the Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 7832. | 1.2 | 6         |
| 57 | Joint impact of muscle mass and waist circumference on type 2 diabetes in Japanese middle-aged adults: The Circulatory Risk in Communities Study (<scp> CIRCS</scp>). <i>Journal of Diabetes</i> , 2020, 12, 677-685.   | 0.8 | 5         |
| 58 | Lifestyle factors associated with prevalent and exacerbated musculoskeletal pain after the Great East Japan Earthquake: a cross-sectional study from the Fukushima Health Management Survey. <i>BMC Public Health</i> , 2020, 20, 677.  | 1.2 | 7         |
| 59 | Urinary 8-Hydroxy-2'-Deoxyguanosine Levels and Cardiovascular Disease Incidence in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 1086-1096.   | 0.9 | 4         |
| 60 | Relationship between the prevalence of polycythemia and factors observed in the mental health and lifestyle survey after the Great East Japan Earthquake. <i>Medicine (United States)</i> , 2020, 99, e18486.   | 0.4 | 6         |
| 61 | Cardiovascular reactivity to acute stress associated with sickness absence among Japanese men and women: A prospective study. <i>Brain and Behavior</i> , 2020, 10, e01541.   | 1.0 | 3         |
| 62 | Influence of post-disaster evacuation on childhood obesity and hyperlipidemia. <i>Pediatrics International</i> , 2020, 62, 669-676.   | 0.2 | 12        |
| 63 | Serum uric acid and risk of stroke and its types: the Circulatory Risk in Communities Study (CIRCS). <i>Hypertension Research</i> , 2020, 43, 313-321.  | 1.5 | 30        |
| 64 | Absorbed radiation doses in the thyroid as estimated by UNSCEAR and subsequent risk of childhood thyroid cancer following the Great East Japan Earthquake. <i>Journal of Radiation Research</i> , 2020, 61, 243-248.  | 0.8 | 29        |
| 65 | Effects of Psychological and Lifestyle Factors on Metabolic Syndrome Following the Fukushima Daiichi Nuclear Power Plant Accident: The Fukushima Health Management Survey. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 1010-1018.  | 0.9 | 11        |
| 66 | Changes in drinking behavior among evacuees after the Fukushima Daiichi Nuclear Power Plant accident: the Fukushima Health Management Survey. <i>Fukushima Journal of Medical Sciences</i> , 2020, 66, 133-142.   | 0.1 | 4         |
| 67 | Analysis of direction of association between radiation risk perception and relocation using a random-intercept and cross lagged panel model: The Fukushima Health Management Survey. <i>SSM - Population Health</i> , 2020, 12, 100706.   | 1.3 | 5         |
| 68 | Job stress factors measured by Brief Job Stress Questionnaire and sickness absence among Japanese workers: A longitudinal study. <i>Fukushima Journal of Medical Sciences</i> , 2020, 66, 88-96.  | 0.1 | 5         |
| 69 | Cytological examination of the thyroid in children and adolescents after the Fukushima Nuclear Power Plant accident: the Fukushima Health Management Survey. <i>Endocrine Journal</i> , 2020, 67, 1233-1238.  | 0.7 | 1         |
| 70 | Parental Recognition of Bullying and Associated Factors Among Children After the Fukushima Nuclear Disaster: A 3-Year Follow-Up Study From the Fukushima Health Management Survey. <i>Frontiers in Psychiatry</i> , 2019, 10, 283.  | 1.3 | 6         |
| 71 | Effects of laughter therapy on quality of life in patients with cancer: An open-label, randomized controlled trial. <i>PLoS ONE</i> , 2019, 14, e0219065.   | 1.1 | 29        |
| 72 | Risk Factors for Problem Drinking among Evacuees in Fukushima following the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>Tohoku Journal of Experimental Medicine</i> , 2019, 248, 239-252.   | 0.5 | 14        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | Unreliable Information as a Risk Factor for Worse Mental Fatigue among Residents in Fukushima after the Nuclear Power Station Accident. <i>Tohoku Journal of Experimental Medicine</i> , 2019, 248, 261-272.   | 0.5 | 7         |
| 74 | Who needs care? - The long-term trends and geographical distribution of deaths due to acute myocardial infarction in Fukushima Prefecture following the Great East Japan Earthquake. <i>International Journal of Disaster Risk Reduction</i> , 2019, 41, 101318.                                     | 1.8 | 3         |
| 75 | Development and Implementation of an Internet Survey to Assess Community Health in the Face of a Health Crisis: Data from the Pregnancy and Birth Survey of the Fukushima Health Management Survey, 2016. <i>International Journal of Environmental Research and Public Health</i> , 2019, 16, 1946. | 1.2 | 2         |
| 76 | Trends in the Incidence of Sudden Deaths and Heart Diseases in Fukushima After the Great East Japan Earthquake. <i>International Heart Journal</i> , 2019, 60, 1253-1258.  | 0.5 | 7         |
| 77 | The latest update on individual external doses in an early stage after the Fukushima nuclear accident. <i>Radiation Protection Dosimetry</i> , 2019, 187, 402-406.   | 0.4 | 7         |
| 78 | Common carotid artery kinking is a predictor of cardiovascular events: A long-term follow-up study using carotid ultrasonography. <i>Echocardiography</i> , 2019, 36, 2227-2233.   | 0.3 | 5         |
| 79 | External Radiation Dose, Obesity, and Risk of Childhood Thyroid Cancer After the Fukushima Daiichi Nuclear Power Plant Accident: The Fukushima Health Management Survey. <i>Epidemiology</i> , 2019, 30, 853-860.  | 1.2 | 22        |
| 80 | The Authors Respond. <i>Epidemiology</i> , 2019, 30, e11.  | 1.2 | 2         |
| 81 | Psychometric evaluation of the simplified Japanese version of the Athens Insomnia Scale: The Fukushima Health Management Survey. <i>Journal of Sleep Research</i> , 2019, 28, e12771.  | 1.7 | 17        |
| 82 | Obesity and mental health improvement following nutritional education focusing on gut microbiota composition in Japanese women: a randomised controlled trial. <i>European Journal of Nutrition</i> , 2019, 58, 3291-3302.   | 1.8 | 31        |
| 83 | Incidence of Thyroid Cancer Among Children and Young Adults in Fukushima, Japan, Screened With 2 Rounds of Ultrasonography Within 5 Years of the 2011 Fukushima Daiichi Nuclear Power Station Accident. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2019, 145, 4.                           | 1.2 | 44        |
| 84 | Mental health promotion at work associated with utilization of the "Guide of Mental Health Professionals for Workers in Osaka". <i>Journal of Occupational Safety and Health</i> , 2019, 12, 145-151.  | 0.0 | 0         |
| 85 | Findings of Thyroid Ultrasound Examination Within 3 Years After the Fukushima Nuclear Power Plant Accident: The Fukushima Health Management Survey. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 861-869.  | 1.8 | 28        |
| 86 | Impact of evacuation on trends in the prevalence, treatment, and control of hypertension before and after a disaster. <i>Journal of Hypertension</i> , 2018, 36, 924-932.  | 0.3 | 15        |
| 87 | Associations of Tobacco Smoking with Impaired Endothelial Function: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 836-845.   | 0.9 | 20        |
| 88 | Association between poor psychosocial conditions and diabetic nephropathy in Japanese type 2 diabetes patients: A cross-sectional study. <i>Journal of Diabetes Investigation</i> , 2018, 9, 162-172.  | 1.1 | 9         |
| 89 | Lifestyle factors and social ties associated with the frequency of laughter after the Great East Japan Earthquake: Fukushima Health Management Survey. <i>Quality of Life Research</i> , 2018, 27, 639-650.  | 1.5 | 18        |
| 90 | The Authors Respond. <i>Epidemiology</i> , 2018, 29, e57-e58.  | 1.2 | 0         |



| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 91  | Associations of disaster-related and psychosocial factors with changes in smoking status after a disaster: a cross-sectional survey after the Great East Japan Earthquake. <i>BMJ Open</i> , 2018, 8, e018943.   | 0.8 | 14        |
| 92  | Effects of lifestyle on hepatobiliary enzyme abnormalities following the Fukushima Daiichi nuclear power plant accident. <i>Medicine (United States)</i> , 2018, 97, e12890.   | 0.4 | 7         |
| 93  | Spatial analysis of the geographical distribution of thyroid cancer cases from the first-round thyroid ultrasound examination in Fukushima Prefecture. <i>Scientific Reports</i> , 2018, 8, 17661.   | 1.6 | 13        |
| 94  | Nonfasting Glucose and Incident Stroke and Its Types—The Circulatory Risk in Communities Study (CIRCS) —. <i>Circulation Journal</i> , 2018, 82, 1598-1604.  | 0.7 | 8         |
| 95  | Reduction of radiation-related anxiety promoted wellbeing after the 2011 disaster: —Fukushima Health Management Survey—™. <i>Journal of Radiological Protection</i> , 2018, 38, 1428-1440.   | 0.6 | 19        |
| 96  | Changes in Risk Perception of the Health Effects of Radiation and Mental Health Status: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 1219.   | 1.2 | 39        |
| 97  | Serum Fatty Acid and Risk of Coronary Artery Disease—The Circulatory Risk in Communities Study (CIRCS) —. <i>Circulation Journal</i> , 2018, 82, 3013-3020.  | 0.7 | 26        |
| 98  | The Relationship between Sleep Time and Mental Health Problems According to the Strengths and Difficulties Questionnaire in Children after an Earthquake Disaster: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 633. | 1.2 | 12        |
| 99  | Association between Serum Uric Acid Level and Ventricular Tachyarrhythmia in Heart Failure Patients with Implantable Cardioverter-Defibrillator. <i>Cardiology</i> , 2018, 140, 47-51.   | 0.6 | 7         |
| 100 | Trajectories of Emotional Symptoms and Peer Relationship Problems in Children after Nuclear Disaster: Evidence from the Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 82.   | 1.2 | 15        |
| 101 | Effect of Radiological Countermeasures on Subjective Well-Being and Radiation Anxiety after the 2011 Disaster: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 124.   | 1.2 | 20        |
| 102 | Impact of social relationships on income—laughter relationships among older people: the JAGES cross-sectional study. <i>BMJ Open</i> , 2018, 8, e019104.   | 0.8 | 7         |
| 103 | Associations Between Childhood Thyroid Cancer and External Radiation Dose After the Fukushima Daiichi Nuclear Power Plant Accident. <i>Epidemiology</i> , 2018, 29, e32-e34.   | 1.2 | 23        |
| 104 | External Dose Estimation in an Early Stage after the Fukushima Daiichi Nuclear Power Plant Accident. <i>Japanese Journal of Health Physics</i> , 2018, 53, 100-110.  | 0.1 | 4         |
| 105 | Serum $\hat{\pm}$ -linolenic and other $\hat{\%}$ -3 fatty acids, and risk of disabling dementia: Community-based nested case—control study. <i>Clinical Nutrition</i> , 2017, 36, 793-797.  | 2.3 | 30        |
| 106 | Effect of evacuation on liver function after the Fukushima Daiichi Nuclear Power Plant accident: The Fukushima Health Management Survey. <i>Journal of Epidemiology</i> , 2017, 27, 180-185.   | 1.1 | 25        |
| 107 | Persistent prevalence of polycythemia among evacuees 4 years after the Great East Japan Earthquake: A follow-up study. <i>Preventive Medicine Reports</i> , 2017, 5, 251-256.  | 0.8 | 8         |
| 108 | Associations of central aortic pressure and brachial blood pressure with flow mediated dilatation in apparently healthy Japanese men: The Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2017, 259, 46-50.  | 0.4 | 8         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 109 | Changes of Posttraumatic Stress Responses in Evacuated Residents and Their Related Factors. <i>Asia-Pacific Journal of Public Health</i> , 2017, 29, 182S-192S.   | 0.4 | 21        |
| 110 | Exercise Habits Are Important for the Mental Health of Children in Fukushima After the Fukushima Daiichi Disaster. <i>Asia-Pacific Journal of Public Health</i> , 2017, 29, 171S-181S.  | 0.4 | 17        |
| 111 | Changes in Cardiovascular Risk Factors After the Great East Japan Earthquake. <i>Asia-Pacific Journal of Public Health</i> , 2017, 29, 47S-55S.   | 0.4 | 64        |
| 112 | The impact of evacuation on the incidence of chronic kidney disease after the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>Clinical and Experimental Nephrology</i> , 2017, 21, 995-1002.  | 0.7 | 26        |
| 113 | Effects of socioeconomic factors on cardiovascular-related symptoms among residents in Fukushima after the Great East Japan Earthquake: a cross-sectional study using data from the Fukushima Health Management Survey. <i>BMJ Open</i> , 2017, 7, e014077.                             | 0.8 | 13        |
| 114 | Comparative Analysis of the Growth Pattern of Thyroid Cancer in Young Patients Screened by Ultrasonography in Japan After a Nuclear Accident. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2017, 144, 57-63.  | 1.2 | 17        |
| 115 | Matched-pair analysis of a multi-institutional cohort reveals that epidermal growth factor receptor mutation is not a risk factor for postoperative recurrence of lung adenocarcinoma. <i>Lung Cancer</i> , 2017, 114, 23-30.   | 0.9 | 12        |
| 116 | Association between markers of arterial stiffness and atrial fibrillation in the Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2017, 263, 244-248.  | 0.4 | 14        |
| 117 | Changes in Hepatobiliary Enzyme Abnormality After the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>Scientific Reports</i> , 2017, 7, 710.  | 1.6 | 8         |
| 118 | Lifestyle-related factors that explain disaster-induced changes in socioeconomic status and poor subjective health: a cross-sectional study from the Fukushima health management survey. <i>BMC Public Health</i> , 2017, 17, 340.  | 1.2 | 19        |
| 119 | Prevalence and Characterization of Thyroid Hemiagenesis in Japan: The Fukushima Health Management Survey. <i>Thyroid</i> , 2017, 27, 1011-1016.   | 2.4 | 21        |
| 120 | Evacuation after the Great East Japan Earthquake was associated with poor dietary intake: The Fukushima Health Management Survey. <i>Journal of Epidemiology</i> , 2017, 27, 14-23.   | 1.1 | 47        |
| 121 | Simulation of expected childhood and adolescent thyroid cancer cases in Japan using a cancer-progression model based on the National Cancer Registry. <i>Medicine (United States)</i> , 2017, 96, e8631.  | 0.4 | 19        |
| 122 | Influence of Post-disaster Evacuation on Incidence of Metabolic Syndrome. <i>Journal of Atherosclerosis and Thrombosis</i> , 2017, 24, 327-337.   | 0.9 | 48        |
| 123 | Perception of Radiation Risk as a Predictor of Mid-Term Mental Health after a Nuclear Disaster: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2017, 14, 1067.  | 1.2 | 32        |
| 124 | The Relationship between Starting to Drink and Psychological Distress, Sleep Disturbance after the Great East Japan Earthquake and Nuclear Disaster: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2017, 14, 1281. | 1.2 | 32        |
| 125 | Fifty-year Time Trends in Blood Pressures, Body Mass Index and their Relations in a Japanese Community: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2017, 24, 518-529.  | 0.9 | 15        |
| 126 | Effects of Changes in Living Place and Communities on Health and Welfare After the Great East Japan Earthquake: A Lesson from a Case of Fukushima. <i>Trends in the Sciences</i> , 2017, 22, 6_10-6_17.   | 0.0 | 0         |



| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 127 | Three-year trend survey of psychological distress, post-traumatic stress, and problem drinking among residents in the evacuation zone after the Fukushima Daiichi Nuclear Power Plant accident [The Fukushima Health Management Survey]. <i>Psychiatry and Clinical Neurosciences</i> , 2016, 70, 245-252. | 1.0 | 74        |
| 128 | Explanatory Meetings on Thyroid Examination for the "Fukushima Health Management Survey" after the Great East Japan Earthquake: Reduction of Anxiety and Improvement of Comprehension. <i>Tohoku Journal of Experimental Medicine</i> , 2016, 239, 333-343.  | 0.5 | 26        |
| 129 | Prevalence of Renal Dysfunction among Evacuees and Non-evacuees after the Great East Earthquake: Results from the Fukushima Health Management Survey. <i>Internal Medicine</i> , 2016, 55, 2563-2569.  | 0.3 | 12        |
| 130 | Hypo-high-density Lipoprotein Cholesterolemia Caused by Evacuation after the Fukushima Daiichi Nuclear Power Plant Accident: Results from the Fukushima Health Management Survey. <i>Internal Medicine</i> , 2016, 55, 1967-1976.  | 0.3 | 34        |
| 131 | Association between psychological distress and dietary intake among evacuees after the Great East Japan Earthquake in a cross-sectional study: the Fukushima Health Management Survey. <i>BMJ Open</i> , 2016, 6, e011534.   | 0.8 | 37        |
| 132 | Comparison of childhood thyroid cancer prevalence among 3 areas based on external radiation dose after the Fukushima Daiichi nuclear power plant accident. <i>Medicine (United States)</i> , 2016, 95, e4472.  | 0.4 | 46        |
| 133 | Inappropriate Suppression of Thyrotropin Concentrations in Young Patients with Thyroid Nodules Including Thyroid Cancer: The Fukushima Health Management Survey. <i>Thyroid</i> , 2016, 26, 717-725.   | 2.4 | 8         |
| 134 | Relationship between HbA1c and risk of retinal hemorrhage in the Japanese general population: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 834-838.   | 1.2 | 2         |
| 135 | Occupational status and job stress in relation to cardiovascular stress reactivity in Japanese workers. <i>Preventive Medicine Reports</i> , 2016, 4, 61-67.   | 0.8 | 19        |
| 136 | Comprehensive Survey Results of Childhood Thyroid Ultrasound Examinations in Fukushima in the First Four Years After the Fukushima Daiichi Nuclear Power Plant Accident. <i>Thyroid</i> , 2016, 26, 843-851.   | 2.4 | 65        |
| 137 | Dehydroepiandrosterone-sulfate is associated with cardiovascular reactivity to stress in women. <i>Psychoneuroendocrinology</i> , 2016, 69, 116-122.   | 1.3 | 9         |
| 138 | Evacuation and Risk of Hypertension After the Great East Japan Earthquake. <i>Hypertension</i> , 2016, 68, 558-564.  | 1.3 | 83        |
| 139 | Drinking Behavior and Mental Illness Among Evacuees in Fukushima Following the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>Alcoholism: Clinical and Experimental Research</i> , 2016, 40, 623-630.   | 1.4 | 33        |
| 140 | Laughter is the Best Medicine? A Cross-Sectional Study of Cardiovascular Disease Among Older Japanese Adults. <i>Journal of Epidemiology</i> , 2016, 26, 546-552.  | 1.1 | 95        |
| 141 | Predictors of severe psychological distress trajectory after nuclear disaster: evidence from the Fukushima Health Management Survey. <i>BMJ Open</i> , 2016, 6, e013400.   | 0.8 | 42        |
| 142 | Epidermal growth factor receptor gene mutation as risk factor for recurrence in patients with surgically resected lung adenocarcinoma: a matched-pair analysis. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2016, 23, 216-222.  | 0.5 | 13        |
| 143 | Psychometric Evaluation of the Japanese Version of the Posttraumatic Stress Disorder Checklist in Community Dwellers Following the Fukushima Daiichi Nuclear Power Plant Incident. <i>SAGE Open</i> , 2016, 6, 215824401665244.  | 0.8 | 24        |
| 144 | The relationship between sodium concentrations in spot urine and blood pressure increases: a prospective study of Japanese general population: the Circulatory Risk in Communities Study (CIRCS). <i>BMC Cardiovascular Disorders</i> , 2016, 16, 55.  | 0.7 | 10        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 145 | Effect of Evacuation on Body Weight After the Great East Japan Earthquake. <i>American Journal of Preventive Medicine</i> , 2016, 50, 553-560.  | 1.6 | 73        |
| 146 | A survey on psychiatric services for an occupational mental health program in Osaka prefecture. <i>Journal of Occupational Safety and Health</i> , 2016, 9, 9-15.   | 0.0 | 0         |
| 147 | Laughter and Subjective Health Among Community-Dwelling Older People in Japan. <i>Journal of Nervous and Mental Disease</i> , 2015, 203, 934-942.   | 0.5 | 25        |
| 148 | Evacuation after the Fukushima Daiichi Nuclear Power Plant Accident Is a Cause of Diabetes: Results from the Fukushima Health Management Survey. <i>Journal of Diabetes Research</i> , 2015, 2015, 1-9.                           | 1.0 | 63        |
| 149 | Health effects of radiation and other health problems in the aftermath of nuclear accidents, with an emphasis on Fukushima. <i>Lancet, The</i> , 2015, 386, 479-488.  | 6.3 | 195       |
| 150 | Increased prevalence of atrial fibrillation after the Great East Japan Earthquake: Results from the Fukushima Health Management Survey. <i>International Journal of Cardiology</i> , 2015, 198, 102-105.                          | 0.8 | 44        |
| 151 | Effects of a traditional herbal medicine on peripheral blood flow in women experiencing peripheral coldness: a randomized controlled trial. <i>BMC Complementary and Alternative Medicine</i> , 2015, 15, 105.                    | 3.7 | 23        |
| 152 | Psychological distress and the perception of radiation risks: the Fukushima health management survey. <i>Bulletin of the World Health Organization</i> , 2015, 93, 598-605.   | 1.5 | 160       |
| 153 | HOW LIFESTYLE AFFECTS HEALTH—CHANGES IN HEALTH STATUS BEFORE AND AFTER THE EARTHQUAKE. <i>Fukushima Journal of Medical Sciences</i> , 2014, 60, 211-212.  | 0.1 | 10        |
| 154 | Life as an evacuee after the Fukushima Daiichi nuclear power plant accident is a cause of polycythemia: the Fukushima Health Management Survey. <i>BMC Public Health</i> , 2014, 14, 1318.  | 1.2 | 30        |
| 155 | Menopausal Status in Relation to Cardiovascular Stress Reactivity in Healthy Japanese Participants. <i>Psychosomatic Medicine</i> , 2014, 76, 701-708.  | 1.3 | 11        |
| 156 | An association between central aortic pressure and subclinical organ damage of the heart among a general Japanese cohort: Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2014, 232, 94-98.               | 0.4 | 9         |
| 157 | Adult Height and Body Mass Index in Relation to Risk of Total Stroke and its Subtypes: The Circulatory Risk in Communities Study. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2014, 23, 667-674.                      | 0.7 | 29        |
| 158 | Effects of a laughter and exercise program on physiological and psychological health among community-dwelling elderly in Japan: Randomized controlled trial. <i>Geriatrics and Gerontology International</i> , 2013, 13, 152-160. | 0.7 | 50        |
| 159 | Alkaline Phosphatase and Risk of Stroke Among Japanese: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2013, 22, 1046-1055.   | 0.7 | 36        |
| 160 | Total and high molecular weight adiponectin levels and risk of cardiovascular disease in individuals with high blood glucose levels. <i>Atherosclerosis</i> , 2013, 229, 222-227.   | 0.4 | 21        |
| 161 | Cardiovascular Disease Epidemiology in Asia. <i>Circulation Journal</i> , 2013, 77, 1646-1652.  | 0.7 | 217       |
| 162 | Insulin secretion capacity in normal glucose tolerance, impaired glucose tolerance, and diabetes in obese and non-obese Japanese patients. <i>Journal of Diabetes Investigation</i> , 2012, 3, 271-275.                           | 1.1 | 24        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 163 | Associations of Anger, Anxiety, and Depressive Symptoms With Carotid Arterial Wall Thickness. <i>Psychosomatic Medicine</i> , 2012, 74, 517-525.  | 1.3 | 30        |
| 164 | Risk factors for sudden cardiac death among Japanese. <i>Journal of Hypertension</i> , 2012, 30, 1137-1143.   | 0.3 | 19        |
| 165 | Non-fasting blood glucose and risk of incident coronary heart disease in middle-aged general population: The Circulatory Risk in Communities Study (CIRCS). <i>Preventive Medicine</i> , 2012, 55, 603-607. | 1.6 | 22        |
| 166 | Associations between alcohol consumption and sleep-disordered breathing among Japanese women. <i>Respiratory Medicine</i> , 2011, 105, 796-800.   | 1.3 | 9         |
| 167 | C-reactive protein levels and risk of stroke and its subtype in Japanese: The Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2011, 217, 187-193.                                   | 0.4 | 35        |
| 168 | Carotid Artery Wall Thickness and Risk of Stroke Subtypes. <i>Stroke</i> , 2011, 42, 397-403.   | 1.0 | 44        |
| 169 | Chronic Kidney Disease and Drinking Status in Relation to Risks of Stroke and Its Subtypes. <i>Stroke</i> , 2011, 42, 2531-2537.  | 1.0 | 53        |
| 170 | Psychological Distress and Cardiovascular Disease: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Epidemiology</i> , 2010, 20, 185-191.   | 1.1 | 38        |
| 171 | Reproductive history, hormone replacement, and incidence of venous thromboembolism: the Longitudinal Investigation of Thromboembolism Etiology. <i>British Journal of Haematology</i> , 2010, 149, 606-612. | 1.2 | 18        |
| 172 | Î³-Glutamyltranspeptidase and Incident Stroke Among Japanese Men and Women. <i>Stroke</i> , 2010, 41, 385-388.  | 1.0 | 40        |
| 173 | Effects of Habitual Alcohol Intake on Ambulatory Blood Pressure, Heart Rate, and Its Variability Among Japanese Men. <i>Hypertension</i> , 2009, 53, 13-19.   | 1.3 | 97        |
| 174 | Serum and Dietary Magnesium and Risk of Ischemic Stroke: The Atherosclerosis Risk in Communities Study. <i>American Journal of Epidemiology</i> , 2009, 169, 1437-1444.                                     | 1.6 | 118       |
| 175 | Perceived Level of Life Enjoyment and Risks of Cardiovascular Disease Incidence and Mortality. <i>Circulation</i> , 2009, 120, 956-963.   | 1.6 | 72        |
| 176 | Trends for Blood Pressure and Its Contribution to Stroke Incidence in the Middle-Aged Japanese Population. <i>Stroke</i> , 2009, 40, 1571-1577.   | 1.0 | 104       |
| 177 | Trends in the Incidence of Coronary Heart Disease and Stroke and Their Risk Factors in Japan, 1964 to 2003. <i>Journal of the American College of Cardiology</i> , 2008, 52, 71-79.                         | 1.2 | 152       |
| 178 | The joint impact on being overweight of self reported behaviours of eating quickly and eating until full : cross sectional survey. <i>BMJ: British Medical Journal</i> , 2008, 337, a2002-a2002.            | 2.4 | 196       |
| 179 | Associations of Psychosocial Factors With Heart Rate and Its Short-Term Variability: Multi-Ethnic Study of Atherosclerosis. <i>Psychosomatic Medicine</i> , 2008, 70, 141-146.                              | 1.3 | 44        |
| 180 | Longitudinal Association of Serum Carotenoids and Tocopherols with Hostility: The CARDIA Study. <i>American Journal of Epidemiology</i> , 2007, 167, 42-50.   | 1.6 | 10        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 181 | Effects of weight training on quality of life in recent breast cancer survivors. <i>Cancer</i> , 2006, 106, 2076-2083.  | 2.0 | 179       |
| 182 | Lipoprotein(a) and Incident Ischemic Stroke. <i>Stroke</i> , 2006, 37, 1407-1412.   | 1.0 | 96        |
| 183 | Risk Factors for Ischemic Stroke Subtypes. <i>Stroke</i> , 2006, 37, 2493-2498.   | 1.0 | 329       |
| 184 | Plasma Fibrinogen Concentrations and Risk of Stroke and Its Subtypes Among Japanese Men and Women. <i>Stroke</i> , 2006, 37, 2488-2492.   | 1.0 | 19        |
| 185 | Carotid Intima-Media Thickness and Plaque Characteristics as a Risk Factor for Stroke in Japanese Elderly Men. <i>Stroke</i> , 2004, 35, 2788-2794.   | 1.0 | 262       |
| 186 | Serum Total Homocysteine Concentrations and Risk of Stroke and Its Subtypes in Japanese. <i>Circulation</i> , 2004, 109, 2766-2772.   | 1.6 | 133       |
| 187 | Prospective Study of Major and Minor ST-T Abnormalities and Risk of Stroke Among Japanese. <i>Stroke</i> , 2003, 34, e250-3.  | 1.0 | 29        |
| 188 | Smoking Raises the Risk of Total and Ischemic Strokes in Hypertensive Men.. <i>Hypertension Research</i> , 2003, 26, 209-217.   | 1.5 | 61        |
| 189 | The relation of anger expression with blood pressure levels and hypertension in rural and urban Japanese communities. <i>Journal of Hypertension</i> , 2002, 20, 21-27.   | 0.3 | 32        |
| 190 | Prospective Study of Depressive Symptoms and Risk of Stroke Among Japanese. <i>Stroke</i> , 2001, 32, 903-908.  | 1.0 | 133       |
| 191 | Validity and Reliability of the Japanese Version of the Selected Anger Expression Scale and Age, Sex, Occupation and Regional Differences in Anger Expression Among Japanese. <i>Journal of Epidemiology</i> , 2000, 10, 118-123. | 1.1 | 8         |
| 192 | Impact of anger expression on blood pressure levels in white-color workers with low-coping behavior. <i>Environmental Health and Preventive Medicine</i> , 2000, 5, 37-42.  | 1.4 | 8         |